

PACKAGE LEAFLET: INFORMATION FOR THE USER

Istin™ 10mg Tablets amlodipine (as besilate)

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Istin is and what it is used for
2. Before you take Istin
3. How to take Istin
4. Possible side effects
5. How to store Istin
6. Further information

1. WHAT ISTIN IS AND WHAT IT IS USED FOR

Istin contains the active substance amlodipine which belongs to a group of medicines called calcium antagonists.

Istin is used to treat high blood pressure (hypertension) or a certain type of chest pain called angina, a rare form of which is Prinzmetal's or variant angina.

In patients with high blood pressure your medicine works by relaxing blood vessels, so that blood passes through them more easily. In patients with angina Istin works by improving blood supply to the heart muscle which then receives more oxygen and as a result chest pain is prevented. Your medicine does not provide immediate relief of chest pain from angina.

2. BEFORE YOU TAKE ISTIN

Do not take Istin

- If you are allergic (hypersensitive) to amlodipine, or any of the other ingredients of your medicine listed in section 6, or to any other calcium antagonists. This may be itching, reddening of the skin or difficulty in breathing.
- If you have severe low blood pressure (hypotension).
- If you have narrowing of the aortic heart valve (aortic stenosis), or cardiogenic shock (a condition where your heart is unable to supply enough blood to the body).
- If you suffer from heart failure after a heart attack.

Take special care with Istin

You should inform your doctor if you have or have had any of the following conditions:

- Recent heart attack
- Heart failure
- Severe increase in blood pressure (Hypertensive crisis)
- Liver disease
- You are elderly and your dose needs to be increased

Use in children and adolescents

Safety and effectiveness have been studied in 6-17 year old boys and in girls. Istin has not been studied in children under the age of 6 years. Istin should only be used for hypertension in children and adolescents from 6 years to 17 years of age (see section 3). For more information, talk to your doctor.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Istin may affect or be affected by other medicines, such as:

- ketoconazole, itraconazole (anti-fungal medicines)
- ritonavir, indinavir, nelfinavir (so called protease inhibitors used to treat HIV)
- rifampicin, erythromycin, clarithromycin (antibiotics)
- hypericum perforatum (St. John's Wort)
- verapamil, diltiazem (heart medicines)
- dantrolene (infusion for severe body temperature abnormalities)
- simvastatin (a cholesterol lowering medicine)

Istin may lower your blood pressure even more if you are already taking other medicines to treat your high blood pressure.

Taking Istin with food and drink

Grapefruit juice and grapefruit should not be consumed by people who are taking Istin. This is because grapefruit and grapefruit juice can lead to an increase in the blood levels of the active ingredient amlodipine, which can cause an unpredictable increase in the blood pressure lowering effect of Istin.

Pregnancy

The safety of amlodipine in human pregnancy has not been established.

If you think you might be pregnant, or are planning to get pregnant, you must tell your doctor before you take Istin.

Breast-feeding

It is not known whether amlodipine is passed into breast milk. If you are breast-feeding or about to start breast-feeding you must tell your doctor before taking Istin.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Istin may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Important information about some of the ingredients of Istin

This medicine contains less than 1mmol of sodium (23mg) per tablet, which means it is essentially 'sodium free'.

3. HOW TO TAKE ISTIN

Always take your medicine exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

The usual initial dose is Istin 5mg once daily. The dose can be increased to Istin 10mg once daily.

Your medicine can be used before or after food and drinks. You should take your medicine at the same time each day with a drink of water. Do not take Istin with grapefruit juice.

Use in children and adolescents

For children and adolescents (6-17 years old), the recommended usual starting dose is 2.5mg a day. The maximum recommended dose is 5mg a day. *For capsules or tablets without a breakline:* Amlodipine 2.5mg is not currently available and the 2.5mg dose cannot be obtained with Istin 5mg tablets as these tablets are not manufactured to break into two equal halves.

It is important to keep taking the tablets. Do not wait until your tablets are finished before seeing your doctor.

If you take more Istin than you should

Taking too many tablets may cause your blood pressure to become low or even dangerously low. You may feel dizzy, lightheaded, faint or weak. If blood pressure drop is severe enough shock can occur. Your skin could feel cool and clammy and you could lose consciousness. Seek immediate medical attention if you take too many Istin tablets.

If you forget to take Istin

Do not worry. If you forget to take a tablet, leave out that dose completely. Take your next dose at the right time. Do not take a double dose to make up for a missed dose.

If you stop taking Istin

Your doctor will advise you how long to take your medicine. Your condition may return if you stop using your medicine before you are advised.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Istin can cause side effects, although not everybody gets them.

Visit your doctor **immediately** if you experience any of the following very rare, severe side effects after taking this medicine.

- Sudden wheeziness, chest pain, shortness of breath or difficulty in breathing
- Swelling of eyelids, face or lips
- Swelling of the tongue and throat which causes great difficulty breathing
- Severe skin reactions including intense skin rash, hives, reddening of the skin over your whole body, severe itching, blistering, peeling and swelling of the skin, inflammation of mucous membranes (Stevens Johnson Syndrome) or other allergic reactions
- Heart attack, abnormal heart beat
- Inflamed pancreas which may cause severe abdominal and back pain accompanied with feeling very unwell

The following **common side effects** have been reported. If any of these cause you problems or if they **last for more than one week**, you should **contact your doctor**.

- Common:** affects 1 to 10 users in 100
- Headache, dizziness, sleepiness (especially at the beginning of treatment)
 - Palpitations (awareness of your heart beat), flushing
 - Abdominal pain, feeling sick (nausea)
 - Ankle swelling (oedema), tiredness

Other side effects that have been reported include the following list. If any of these get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

- Uncommon:** affects 1 to 10 users in 1,000
- Mood changes, anxiety, depression, sleeplessness
 - Trembling, taste abnormalities, fainting, weakness
 - Numbness or tingling sensation in your limbs, loss of pain sensation
 - Visual disturbances, double vision, ringing in the ears
 - Low blood pressure
 - Sneezing/running nose caused by inflammation of the lining of the nose (rhinitis)
 - Altered bowel habits, diarrhoea, constipation, indigestion, dry mouth, vomiting (being sick)
 - Hair loss, increased sweating, itchy skin, red patches on skin, skin discolouration
 - Disorder in passing urine, increased need to urinate at night, increased number of times of passing urine
 - Inability to obtain an erection; discomfort or enlargement of the breasts in men
 - Weakness, pain, feeling unwell
 - Joint or muscle pain, muscle cramps, back pain
 - Weight increase or decrease

Rare: affects 1 to 10 users in 10,000

- Confusion

Very rare: affects less than 1 user in 10,000

- Decreased numbers of white blood cells, decrease in blood platelets which may result in unusual bruising or easy bleeding (red blood cell damage)
- Excess sugar in blood (hyperglycaemia)

- A disorder of the nerves which can cause weakness, tingling or numbness
- Cough, swelling of the gums
- Abdominal bloating (gastritis)
- Abnormal liver function, inflammation of the liver (hepatitis), yellowing of the skin (jaundice), liver enzyme increase which may have an effect on some medical tests
- Increased muscle tension
- Inflammation of blood vessels, often with skin rash
- Sensitivity to light
- Disorders combining rigidity, tremor, and/or movement disorders

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

5. HOW TO STORE ISTIN

This treatment is for **YOU**. DO NOT give it to others. It may not suit them.

- Keep out of the sight and reach of children.
- Do not use Istin after the expiry date (EXP) which is stated on the carton and the blister. The expiry date refers to the last day of that month.
- If your tablets appear to be discoloured, damaged or show any other signs of deterioration, please return these to your pharmacist who will advise you further.
- Keep your medicine in a dry place.
- Do not store above 25°C.
- Medicines should not be disposed of via waste water or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

What Istin contains

The active substance is amlodipine (as amlodipine besilate). Each tablet contains 10mg amlodipine (as besilate). Each tablet also contains other ingredients: anhydrous calcium hydrogen phosphate, magnesium stearate, microcrystalline cellulose and sodium starch glycolate.

What Istin looks like and contents of the pack

Istin 10mg tablets are white, emerald shaped and marked AML-10 on one side and the Pfizer logo on the other.

Istin comes in packs of 28 tablets.

Manufacturer

Your tablets are manufactured by: Pfizer Manufacturing Deutschland GmbH, Heinrich-Mack-Strasse 35, 89257 Illertissen, Germany.
Procured from within the EU by the Parallel Product Authorisation holder: Imbat Ltd, Unit L2, North Ring Business Park, Santry, Dublin 9.
The product is repackaged by: Doncaster Pharmaceuticals Group Ltd, Kirk Sandall, Doncaster, South Yorkshire, DN3 1QR, UK.
Distributed by: Eurodrug Ltd, Santry, Dublin 9.
PPA No: 1151/10/2

This medicinal product is authorised in the Member States of the EEA under the following names:

Tablets

Austria, Bulgaria, Denmark, Estonia, Finland, Germany, Hungary, Italy, Latvia, Lithuania, Netherlands, Poland, Portugal, Romania, Slovakia, Slovenia, Sweden: Norvasc
Czech Republic: Zorem
Denmark: Amlodipine Pfizer
Ireland, Malta, United Kingdom: Istin
Ireland: Amlodipine Pfizer 10mg tablets
United Kingdom : Amlodipine 10mg tablets
Italy: Monopina
Spain : Norvas 10mg comprimidos
Spain: Amlodipino Pharmacia 10mg comprimidos
United Kingdom : Amlodipine 10mg tablets

Leaflet revision and issue date (ref): 15.07.13

Istin™ is a trademark of Pfizer Ltd.