

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Nexium 40 mg Gastro-resistant Tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each gastro-resistant tablet contains 44.5 mg esomeprazole magnesium trihydrate equivalent to 40 mg esomeprazole.
Excipient(s) with known effect:
Each gastro resistant tablet contains sucrose.
For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Gastro-resistant tablet.

Product imported from the UK, Greece and Italy:
Pink, oblong, biconvex, film-coated tablet engraved ‘40 MG’ on one side and ‘A/EI’ on the other side.

4 CLINICAL PARTICULARS

As per PA0970/027/002

5 PHARMACOLOGICAL PROPERTIES

As per PA0970/027/002

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Glycerol monostearate 40-55
Hyprolose
Hypromellose
Iron oxide (reddish-brown) (E 172)
Magnesium stearate
Methacrylic acid ethyl acrylate copolymer (1:1) dispersion 30 per cent
Microcrystalline cellulose
Synthetic paraffin
Macrogol
Polysorbate 80
Crospovidone
Sodium stearyl fumarate
Sugar spheres (sucrose and maize starch)
Talc
Titanium dioxide (E171)
Triethyl citrate

6.2 Incompatibilities

Not applicable

6.3 Shelf life

The shelf life expiry date of this product is the date shown on the blister strips and outer carton of the product as marketed in the country of origin.

6.4 Special precautions for storage

Do not store above 30°C.

Store in the original package in order to protect from moisture.

6.5 Nature and contents of container

Over-labelled outer carton containing blister package. Pack size 28

Blister package reboxed into cartons. Pack size 28

6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product

Administration through gastric tube

1. Put the tablet into an appropriate syringe and fill the syringe with approximately 25 mL water and approximately 5 mL air. For some tubes, dispersion in 50 mL water is needed to prevent the pellets from clogging the tube.
2. Immediately shake the syringe for approximately 2 minutes to disperse the tablet.
3. Hold the syringe with the tip up and check that the tip has not clogged.
4. Attach the syringe to the tube whilst maintaining the above position.
5. Shake the syringe and position it with the tip pointing down. Immediately inject 5-10 mL into the tube. Invert the syringe after injection and shake (the syringe must be held with the tip pointing up to avoid clogging of the tip).
6. Turn the syringe with the tip down and immediately inject another 5-10 mL into the tube. Repeat this procedure until the syringe is empty.
7. Fill the syringe with 25 mL of water and 5 mL of air and repeat step 5 if necessary to wash down any sediment left in the syringe. For some tubes, 50 mL water is needed.

7 PARALLEL PRODUCT AUTHORISATION HOLDER

Imbat Limited
Unit L2
North Ring Business Park
Santry
Dublin 9

8 PARALLEL PRODUCT AUTHORISATION NUMBER

PPA1151/017/002

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 22nd September 2006

Date of last renewal: 22nd September 2011

10 DATE OF REVISION OF THE TEXT

July 2017