

## PACKAGE LEAFLET: INFORMATION FOR THE USER

**Lexapro® 5mg Film-coated tablets**  
**Lexapro® 10mg Film-coated tablets**  
**Lexapro® 15mg Film-coated tablets**  
**Lexapro® 20mg Film-coated tablets**  
escitalopram (as oxalate)

### Read all of this leaflet carefully before you start taking this medicine

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

#### In this leaflet:

- What Lexapro is and what it is used for
- Before you take Lexapro
- How to take Lexapro
- Possible side effects
- How to store Lexapro
- Further information

## 1. WHAT LEXAPRO IS AND WHAT IT IS USED FOR

Lexapro belongs to a group of antidepressants called selective serotonin reuptake inhibitors (SSRIs). These medicines act on the serotonin-system in the brain by increasing the serotonin level. Disturbances in the serotonin-system are considered an important factor in the development of depression and related diseases.

Lexapro contains escitalopram and is used to treat depression (major depressive episodes) and anxiety disorders (such as panic disorder with or without agoraphobia, social anxiety disorder, generalised anxiety disorder and obsessive-compulsive disorder).

## 2. BEFORE YOU TAKE LEXAPRO

### Do not take Lexapro

- if you are allergic (hypersensitive) to escitalopram or any of the other ingredients of Lexapro (see section 6 "Further information").
- if you take other medicines that belong to a group called MAO inhibitors, including selegiline (used in the treatment of Parkinson's disease), moclobemide (used in the treatment of depression) and linezolid (an antibiotic).
- if you are born with or have had an episode of abnormal heart rhythm (seen at ECG: an examination to evaluate how the heart is functioning).
- If you take medicines for heart rhythm problems or that may affect the heart's rhythm (see section 2 "Taking other medicines")

### Take special care with Lexapro

Please tell your doctor if you have any other condition or illness, as your doctor may need to take this into consideration.

In particular, tell your doctor:

- if you have epilepsy. Treatment with Lexapro should be stopped if seizures occur for the first time or if there is an increase in the seizure frequency (see also section 4 "Possible side effects").
- if you suffer from impaired liver or kidney function. Your doctor may need to adjust your dosage.
- if you have diabetes. Treatment with Lexapro may alter glycaemic control. Insulin and/or oral hypoglycaemic dosage may need to be adjusted.
- if you have a decreased level of sodium in the blood.
- if you have a tendency to easily develop bleedings or bruises.
- if you are receiving electroconvulsive treatment.
- if you have coronary heart disease.
- if you suffer or have suffered from heart problems or have recently had a heart attack.
- if you have a low resting heart-rate and/or you know that you may have salt depletion as a result of prolonged severe diarrhoea and vomiting (being sick) or usage of diuretics (water tablets).
- if you experience a fast or irregular heartbeat, fainting, collapse or dizziness on standing up, which may indicate abnormal functioning of the heart rate.

### Please note

Some patients with manic-depressive illness may enter into a manic phase. This is characterised by unusual and rapidly changing ideas, inappropriate happiness and excessive physical activity. If you experience this, contact your doctor.

Symptoms such as restlessness or difficulty to sit or stand still can also occur during the first weeks of the treatment. Tell your doctor immediately if you experience these symptoms.

### Thoughts of suicide and worsening of your depression or anxiety disorder

If you are depressed and/or have anxiety disorders you can sometimes have thoughts of harming or killing yourself. These may be increased when first starting antidepressants, since these medicines all take time to work, usually about two weeks but sometimes longer.

**You** may be more likely to think like this:

- If you have previously had thoughts about killing or harming yourself.
- If you are a **young adult**. Information from clinical trials has shown an increased risk of suicidal behaviour in adults aged less than 25 years with psychiatric conditions who were treated with an antidepressant.

If you have thoughts of harming or killing yourself at any time, **contact your doctor or go to a hospital straight away.**

**You may find it helpful to tell a relative or close friend** that you are depressed or have an anxiety disorder, and ask them to read this leaflet. You might ask them to tell you if they think your depression or anxiety is getting worse, or if they are worried about changes in your behaviour.

### Use in children and adolescents under 18 years of age

Lexapro should normally not be used for children and adolescents under 18 years. Also, you should know that patients under 18 have an increased risk of side effects such as suicide attempts, suicidal thoughts and hostility (predominantly aggression, oppositional behaviour and anger) when they take this class of medicines. Despite this, your doctor may prescribe Lexapro for patients under 18 because he/she decides that this is in their best interest. If your doctor has prescribed Lexapro for a patient under 18 and you want to discuss this, please go back to your doctor. You should inform your doctor if any symptoms listed above develop or worsen when patients under 18 are taking Lexapro. Also, the long term safety effects concerning growth, maturation and cognitive and behavioural development of Lexapro in this age group have not yet been demonstrated.

### Taking other medicines

Please tell your doctor if you are taking or have recently taken any other medicines, including medicines obtained without prescription.

Tell your doctor if you are taking any of the following medicines:

- "Non-selective monoamine oxidase inhibitors (MAOIs)", containing phenelzine, iproniazid, isocarboxazid, nialamide and tranylcypromine as active ingredients. If you have taken any of these medicines you will need to wait 14 days before you start taking Lexapro. After stopping Lexapro you must allow 7 days before taking any of these medicines.
- "Reversible, selective MAO-A inhibitors", containing moclobemide (used to treat depression).
- "Irreversible MAO-B inhibitors", containing selegiline (used to treat Parkinson's disease). These increase the risk of side effects.
- The antibiotic linezolid.
- Lithium (used in the treatment of manic-depressive disorder) and tryptophan.
- Imipramine and desipramine (both used to treat depression).
- Sumatriptan and similar medicines (used to treat migraine) and tramadol (used against severe pain). These increase the risk of side effects.
- Cimetidine and omeprazole (used to treat stomach ulcers), fluvoxamine (antidepressant) and ticlopidine (used to reduce the risk of stroke). These may cause increased blood levels of Lexapro.
- St. John's Wort (hypericum perforatum) - a herbal remedy used for depression.
- Acetylsalicylic acid and non-steroidal anti-inflammatory drugs (medicines used for pain relief or to thin the blood, so called anti-coagulant). These may increase bleeding tendency.
- Warfarin, dipyridamole, and phenprocoumon (medicines used to thin the blood, so called anti-coagulant). Your doctor will probably check the coagulation time of your blood when starting and discontinuing Lexapro in order to verify that your dose of anti-coagulant is still adequate.
- Mefloquin (used to treat Malaria), bupropion (used to treat depression) and tramadol (used to treat severe pain) due to a possible risk of a lowered threshold for seizures.
- Neuroleptics (medicines to treat schizophrenia, psychosis) due to a possible risk of a lowered threshold for seizures, and antidepressants.
- Flecainide, propafenone, and metoprolol (used in cardiovascular diseases), clomipramine, and nortriptyline (antidepressants) and risperidone, thioridazine, and haloperidol (antipsychotics). The dosage of Lexapro may need to be adjusted.

Do not take Lexapro if you take medicines for heart rhythm problems or medicines that may affect the heart's rhythm, such as Class IA and III antiarrhythmics, antipsychotics (e.g. phenothiazine derivatives, pimozide, haloperidol), tricyclic antidepressants, certain antimicrobial agents (e.g. sparflaxacin, moxifloxacin, erythromycin IV, pentamidine, anti-malarial treatment particularly halofantrine), certain antihistamines (astemizole, mizolastine). If you have any further questions about this you should speak to your doctor.

### Taking Lexapro with food and drink

Lexapro can be taken with or without food (see section 3 "How to take Lexapro").

As with many medicines, combining Lexapro with alcohol is not advisable, although Lexapro is not expected to interact with alcohol.

### Fertility, pregnancy and breast-feeding

Inform your doctor if you are pregnant or planning to become pregnant. Do not take Lexapro if you are pregnant or breast-feeding unless you and your doctor have discussed the risks and benefits involved.

If you take Lexapro during the last 3 months of your pregnancy you should be aware that the following effects may be seen in your newborn baby: trouble with breathing, blue-ish skin, fits, body temperature changes, feeding difficulties, vomiting, low blood sugar, stiff or floppy muscles, vivid reflexes, tremor, jitteriness, irritability, lethargy, constant crying, sleepiness and sleeping difficulties. If your newborn baby has any of these symptoms, please contact your doctor immediately.

If used during pregnancy Lexapro should never be stopped abruptly.

Ask your doctor or pharmacist for advice before taking any medicine.

Make sure your midwife and/or doctor know you are on Lexapro. When taken during pregnancy, particularly in the last 3 months of pregnancy, medicines like Lexapro may increase the risk of a serious condition in babies, called persistent pulmonary hypertension of the newborn (PPHN), making the baby breathe faster and appear bluish. These symptoms usually begin during the first 24 hours after the baby is born. If this happens to your baby you should contact your midwife and/or doctor immediately.

Citalopram, a medicine like escitalopram, has been shown to reduce the quality of sperm in animal studies. Theoretically, this could affect fertility, but impact on human fertility has not been observed as yet.

### Driving and using machines

You are advised not to drive a car or operate machinery until you know how Lexapro affects you.

## 3. HOW TO TAKE LEXAPRO

Always take Lexapro exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

#### Adults

##### Depression

The normally recommended dose of Lexapro is 10mg taken as one daily dose. The dose may be increased by your doctor to a maximum of 20mg per day.

##### Panic disorder

The starting dose of Lexapro is 5mg as one daily dose for the first week before increasing the dose to 10mg per day. The dose may be further increased by your doctor to a maximum of 20mg per day.

##### Social anxiety disorder

The normally recommended dose of Lexapro is 10mg taken as one daily dose. Your doctor can either decrease your dose to 5mg per day or increase the dose to a maximum of 20mg per day, depending on how you respond to the medicine.

##### Generalised anxiety disorder

The normally recommended dose of Lexapro is 10mg taken as one daily dose. The dose may be increased by your doctor to a maximum of 20mg per day.

##### Obsessive-compulsive disorder

The recommended dose of Lexapro is 10mg taken as one daily dose. The dose may be increased by your doctor to a maximum of 20mg per day.

##### Elderly patients (above 65 years of age)

The recommended starting dose of Lexapro is 5mg taken as one daily dose. The dose may be increased by your doctor to 10mg per day.

##### Children and adolescents (below 18 years of age)

Lexapro should not normally be given to children and adolescents. For further information please see section 2 "Before you take Lexapro".

You can take Lexapro with or without food. Swallow the tablet with some water. Do not chew them, as the taste is bitter.

If necessary, you can divide the tablets by firstly placing the tablet on a flat surface with the score facing upwards. The tablets may then be broken by pressing down on each end of the tablet, using both forefingers as shown in the drawing.



#### Duration of treatment

It may take a couple of weeks before you start to feel better. Continue to take Lexapro even if it takes some time before you feel any improvement in your condition. Do not change the dose of your medicine without talking to your doctor first. Continue to take Lexapro for as long as your doctor recommends. If you stop your treatment too soon, your symptoms may return. It is recommended that treatment should be continued for at least 6 months after you feel well again.

#### If you take more Lexapro than you should

If you take more than the prescribed dose of Lexapro, contact your doctor or nearest hospital emergency department immediately. Do this even if there are no signs of discomfort. Some of the signs of an overdose could be dizziness, tremor, agitation, convulsion, coma, nausea, vomiting, change in heart rhythm, decreased blood pressure and change in body fluid/salt balance. Take the Lexapro box/container with you when you go to the doctor or hospital.

#### If you forget to take Lexapro

Do not take a double dose to make up for forgotten doses. If you do forget to take a dose, and you remember before you go to bed, take it straight away. Carry on as usual the next day. If you only remember during the night, or the next day, leave out the missed dose and carry on as usual.

#### If you stop taking Lexapro

Do not stop taking Lexapro until your doctor tells you to do so. When you have completed your course of treatment, it is generally advised that the dose of Lexapro is gradually reduced over a number of weeks.

When you stop taking Lexapro, especially if it is abruptly, you may feel discontinuation symptoms. These are common when treatment with Lexapro is stopped. The risk is higher, when Lexapro has been used for a long time or in high doses or when the dose is reduced too quickly. Most people find that the symptoms are mild and go away on their own within two weeks. However, in some patients they may be severe in intensity or they may be prolonged (2-3 months or more). If you get severe discontinuation symptoms when you stop taking Lexapro, please contact your doctor. He or she may ask you to start taking your tablets again and come off them more slowly. Discontinuation symptoms include: Feeling dizzy (unsteady or off-balance), feelings like pins and needles, burning sensations and (less commonly) electric shock sensations, including in the head, sleep disturbances (vivid dreams, nightmares, inability to sleep), feeling anxious, headaches, feeling sick (nausea), sweating (including night sweats), feeling restless or agitated, tremor (shakiness), feeling confused or disorientated, feeling emotional or irritable, diarrhoea (loose stools), visual disturbances, fluttering or pounding heartbeat (palpitations). If you have any further questions on the use of this product, ask your doctor or pharmacist.

## 4. POSSIBLE SIDE EFFECTS

Like all medicines, Lexapro can cause side effects, although not everybody gets them. The side effects usually disappear after a few weeks of treatment. Please be aware that many of the effects may also be symptoms of your illness and therefore will improve when you start to get better.

#### See your doctor if you get any of the following side effects during treatment:

Uncommon (affects 1 to 10 users in 1000):

- Unusual bleeds, including gastrointestinal bleeds

Rare (affects 1 to 10 users in 10000):

- If you experience swelling of skin, tongue, lips, or face, or have difficulties breathing or swallowing (allergic reaction), contact your doctor or go to a hospital straight away.
- If you have a high fever, agitation, confusion, trembling and abrupt contractions of muscles these may be signs of a rare condition called serotonin syndrome. If you feel like this contact your doctor.

If you experience the following side effects you should contact your doctor or go to the hospital straight away:

- Difficulties urinating
- Seizures (fits), see also section "Take special care with Lexapro"

- Yellowing of the skin and the white in the eyes are signs of liver function impairment/hepatitis
- Fast, irregular heart beat, fainting which could be symptoms of a life-threatening condition known as Torsade de Pointes

#### In addition to above the following side effects have been reported:

Very common (affects more than 1 user in 10):

- Feeling sick (nausea)

Common (affects 1 to 10 users in 100):

- Blocked or runny nose (sinusitis)
- Decreased or increased appetite
- Anxiety, restlessness, abnormal dreams, difficulties falling asleep, feeling sleepy, dizziness, yawning, tremors, prickling of the skin
- Diarrhoea, constipation, vomiting, dry mouth
- Increased sweating
- Pain in muscles and joints (arthralgia and myalgia)
- Sexual disturbances (delayed ejaculation, problems with erection, decreased sexual drive and women may experience difficulties achieving orgasm)
- Fatigue, fever
- Increased weight

Uncommon (affects 1 to 10 users in 1000):

- Nettle rash (urticaria), rash, itching (pruritus)
- Grinding one's teeth, agitation, nervousness, panic attack, confusion state
- Disturbed sleep, taste disturbance, fainting (syncope)
- Enlarged pupils (mydriasis), visual disturbance, ringing in the ears (tinnitus)
- Loss of hair
- Vaginal bleeding
- Decreased weight
- Fast heart beat
- Swelling of the arms or legs
- Nosebleeds

Rare (affects 1 to 10 users in 10000):

- Aggression, depersonalisation, hallucination
- Slow heart beat

Some patients have reported (frequency can not be estimated from the available data):

- Thoughts of harming yourself or thoughts of killing yourself, see also section "Take special care with Lexapro"
- Decreased levels of sodium in the blood (the symptoms are feeling sick and unwell with weak muscles or confused)
- Dizziness when you stand up due to low blood pressure (orthostatic hypotension)
- Abnormal liver function test (increased amounts of liver enzymes in the blood)
- Movement disorders (involuntary movements of the muscles)
- Painful erections (priapism)
- Bleeding disorders including skin and mucous bleeding (ecchymosis) and low level of blood platelets (thrombocytopenia)
- Sudden swelling of skin or mucosa (angioedemas)
- Increase in the amount of urine excreted (inappropriate ADH secretion)
- Flow of milk in women that are not nursing
- Mania
- An increased risk of bone fractures has been observed in patients taking this type of medicine
- Alteration of the heart rhythm (called "prolongation of QT interval", seen on ECG, electrical activity of the heart).

In addition, a number of side effects are known to occur with drugs that work in a similar way to escitalopram (the active ingredient of Lexapro). These are:

- Motor restlessness (akathisia)
- Anorexia

**If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.**

## 5. HOW TO STORE LEXAPRO

Always keep medicines out of the reach and sight of children. This medicinal product does not require any special storage conditions. Do not use Lexapro after the expiry date which is stated on the blister and carton after Exp. The expiry date refers to the last day of that month. You should return any left over tablets to your pharmacist for safe disposal. Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment. If your tablets appear to be discoloured, damaged or show any other signs of deterioration, please return these to your pharmacist who will advise you further.

## 6. FURTHER INFORMATION

#### What Lexapro contains

The active substance is escitalopram. Each Lexapro tablet contains 5mg, 10mg, 15mg or 20mg escitalopram (as oxalate). The other ingredients are: Tablet core: microcrystalline cellulose, silica, colloidal anhydrous, talc, croscarmellose sodium and magnesium stearate. Coating: hypromellose, macrogol 400 and titanium dioxide (E171).

#### What Lexapro looks like and the contents of the pack

Lexapro 5mg tablets are round, white, film-coated tablets marked with "EK" on one side and plain on the other.

Lexapro 10mg tablets are oval, white, scored film-coated tablets marked with "E" and "L" on each side of the score on one side and plain on the other.

Lexapro 15mg tablets are oval, white film-coated tablets. The tablets are scored and marked with "E" and "M" on each side of the score on one side of the tablet.

Lexapro 20mg tablets are oval, white, scored, film-coated tablets marked with "E" and "N" on each side of the score on one side and plain on the other.

Your medicine is available in blister packs of 28 film-coated tablets.

#### Manufacturer

Your tablets are manufactured by: H. Lundbeck A/S, Ottiliavej 9, DK-2500 Copenhagen, Denmark. Procured from within the EU by the PPA holder: Imbat Ltd, Unit L2, North Ring Business Park, Santry, Dublin 9. The product is repackaged by: Doncaster Pharmaceuticals Group Ltd, Kirk Sandall, Doncaster, South Yorkshire, DN3 1QR, UK. Distributed by: Eurodrug Ltd, Santry, Dublin 9.

PPA No: 1151/25/2 Lexapro 5mg Tablets  
1151/25/3 Lexapro 10mg Tablets  
1151/25/4 Lexapro 15mg Tablets  
1151/25/5 Lexapro 20mg Tablets

Leaflet revision & issue date (ref): 14.06.12

Lexapro® is a registered trademark of H. Lundbeck A/S, Denmark.

#### This medicinal product is authorised in the Member States of the EEA under the following names:

Austria: CipraleX	Latvia: CipraleX
Belgium: Siprolexa	Lithuania: CipraleX
Bulgaria: CipraleX	Luxembourg: Siprolexa
Cyprus: CipraleX	Malta: CipraleX
Czech Republic: CipraleX	Norway: CipraleX
Denmark: CipraleX	Poland: CipraleX
Estonia: CipraleX	Portugal: CipraleX
Finland: CipraleX	Romania: CipraleX
France: SeropleX	Slovakia: CipraleX
Germany: CipraleX	Slovenia: CipraleX
Greece: CipraleX	Spain: CipraleX
Hungary: CipraleX	Sweden: CipraleX
Iceland: CipraleX	United Kingdom: CipraleX
Ireland: Lexapro	
Italy: CipraleX	