

PACKAGE LEAFLET: INFORMATION FOR THE USER

BETOPTIC® 0.5% w/v Eye Drops, Solution (Betaxolol (as hydrochloride))

Your medicine is available using the above name but will be referred to as Betoptic 0.5% throughout the leaflet.

Read all of this leaflet carefully before you start using this medicine

- **Keep this leaflet.** You may need to read it again.
- **If you have any further questions**, ask your doctor or your pharmacist.
- This medicine has been prescribed for you. **Do not pass it on to others.** It may harm them, even if their symptoms are the same as yours.
- **If any of the side effects get serious**, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

IN THIS LEAFLET

1. What Betoptic 0.5% is and what it is used for
2. Before you use Betoptic 0.5%
3. How to use Betoptic 0.5%
4. Possible side effects
5. How to store Betoptic 0.5%
6. Contents of the pack and other information

1. WHAT BETOPTIC 0.5% IS AND WHAT IT IS USED FOR

Betoptic 0.5% belongs to a group of medicines known as beta blockers.

It is used to treat glaucoma or ocular hypertension (high pressure in the eye) by reducing the fluid pressure in your eye(s).

2. BEFORE YOU USE BETOPTIC 0.5%

Do not use Betoptic 0.5%...

- If you are **allergic** to betaxolol, beta-blockers or any of the other ingredients listed in section 6. Ask your doctor for advice.
- If you have now or have had in the past, **respiratory problems** such as **severe asthma, severe chronic obstructive bronchitis** (severe lung condition which may cause wheeziness, difficulty in breathing and/or long-standing cough).
- If you have a **slow heart beat, heart failure**, or **disorders of heart rhythm (irregular heartbeats)**.

Take special care...

Before you use this medicine, tell your doctor if you have now or have had in the past

- **coronary heart disease** (symptoms can include chest pain or tightness, breathlessness or choking), **heart failure, low blood pressure** (hypotension)
- **disturbances of heart rate** such as slow heart beat (bradycardia)
- **breathing problems, asthma or chronic obstructive pulmonary disease** (lung disease which may cause wheeziness, difficulty in breathing and/or long-standing cough)
- **poor blood circulation disease** (such as Raynaud's disease or Raynaud's syndrome)
- **diabetes**, as betaxolol may mask the signs and symptoms of low blood sugar
- **overactivity of the thyroid gland** as betaxolol may mask the signs and symptoms
- **a muscle weakening disease** such as myasthenia gravis. Your symptoms may get worse
- **angle-closure glaucoma**
- **dry eyes** (Sicca Syndrome)

Tell your doctor before you have an operation that you are using Betoptic 0.5% as betaxolol may change the effects of some medicines used during anaesthesia. **If any of these apply** you may still be able to use Betoptic 0.5%, but **discuss it with your doctor first**.

Using other medicines

Betoptic 0.5% can affect or be affected by other medicines you are using, including other eye drops for the treatment of glaucoma. Tell your doctor if you are using or intend to use medicines to **lower blood pressure, heart medicine or medicines to treat diabetes**.

Betoptic 0.5% may reduce the effectiveness of adrenaline, which can be used to treat serious allergic reactions (anaphylaxis). **Tell your doctor if you have a history of anaphylaxis or allergic reactions**.

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

If you are using more than one type of eye drop, wait 5 minutes between using each one. Eye ointments should be administered last.

Pregnancy and breast-feeding

Do not use Betoptic 0.5% if you are **pregnant** unless your doctor considers it necessary.

Do not use Betoptic 0.5% if you are **breast-feeding**. Betaxolol may get into your breast milk.

Ask your doctor for advice before taking any medicine during breast-feeding.

Driving and using machines

If your sight is affected in any way following the use of Betoptic 0.5%, **you should not drive or use any machines**.

Important information if you wear Contact Lenses

Do not use the drops while wearing **contact lenses**. Wait at least 15 minutes after use before putting your lenses back in. There is a preservative in Betoptic 0.5% (benzalkonium chloride) that can discolour soft contact lenses.

3. HOW TO USE BETOPTIC 0.5%

Betoptic 0.5% should only be used in the eye(s).



The usual dose

The usual dose is 1 drop in the affected eye(s) twice daily.

Remove the loose collar from the cap when you open the bottle.

Always use Betoptic 0.5% exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

How to use



- Wash your hands before you start.
- Twist off the bottle cap.
- Hold the bottle pointing down, between your thumb and fingers.
- Tilt your head back.
- Pull down your lower eyelid with a finger, until there is a 'pocket' between the eyelid and your eye. The drop will go in here (picture 1).
- Bring the bottle tip close to the eye. Do this in front of a mirror if it helps.
- **Do not touch your eye or eyelid**, surrounding areas or other surfaces with the dropper. It could infect the drops.

- **Gently press on the base** of the bottle to release one drop at a time (picture 2).
- **Do not squeeze the bottle**, only a gentle press on the bottom is needed.
- If you use drops in both eyes, repeat the steps for your other eye. Put the bottle cap firmly back on immediately after use.
- After using Betoptic 0.5%, **press a finger into the corner of your eye by the nose** (picture 3) for 2 minutes. This helps to stop betaxolol getting into the rest of the body.
- **If a drop misses your eye**, try again.
- **If you miss a dose**, just take it as soon as possible. However, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **Do not use a double dose to make up for a missed dose.**
- **If you use more Betoptic 0.5% than you should** it can be washed out of your eye with warm water.

If you have any further questions on the use of Betoptic 0.5%, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Betoptic 0.5% eye drops solution can cause side effects, although not everybody gets them.

You can usually carry on taking the drops, unless the effects are serious. If you are worried, talk to a doctor or pharmacist. Do not stop using Betoptic 0.5% without speaking to your doctor.

In clinical trials with Betaxolol eye drops, 12 out of every 100 patients reported eye discomfort. Other side effects experienced by patients during clinical trials with BETAXOLOL eye drops are:

Common (affects 1 to 10 users in 100)

- Blurred vision, watery eyes
- Headache

Uncommon (affects 1 to 10 users in 1,000)

- Inflammation of the eye surface, conjunctivitis or symptoms of conjunctivitis, visual impairment, sensitivity to light, painful, dry or tired eyes, excessive blinking, irritated, red or swollen eyes, eye discharge, weeping eyelids, bloodshot eyes
- Slow heart beat or unusually rapid heartbeat
- Asthma, difficulty breathing, blocked nose
- Nausea

Rare (affects 1 to 10 users in 10,000)

- Cataract formation, drooping of the eyelid, loss of eyelashes, dry eye syndrome
- Anxiety, difficulty sleeping (insomnia), depression
- Fainting
- Low blood pressure
- Cough, runny nose
- Taste disturbances
- Inflamed, itchy skin or rash, hair loss
- Muscular weakness and pain not caused by exercise
- Libido decreased

The following side effects have also been reported by people using BETAXOLOL eye drops.

The frequency cannot be estimated from the available data:

- Hypersensitivity reaction
- Dizziness
- Inflammation of the eyelid
- Changes in the rhythm or speed of the heartbeat
- Lack or loss of strength

Additional adverse reactions have been seen with other ophthalmic beta-blockers and could potentially occur with Betoptic eye drops. The frequency is unknown:

- Generalised allergic reactions including swelling beneath the skin (that can occur in areas such as the face and limbs, and can obstruct the airway which may cause difficulty swallowing or breathing), hives (or itchy rash), localised and generalised rash, itchiness, severe sudden life-threatening allergic reaction.
- Low blood glucose levels.

- Nightmares, memory loss.
- Stroke, reduced blood supply to the brain, increases in signs and symptoms of myasthenia gravis (muscle disorder), unusual sensations (like pins and needles).
- Detachment of the layer below the retina that contains blood vessels following filtration surgery which may cause visual disturbances, corneal erosion (damage to the front layer of the eyeball), decreased sensitivity of the eye, double vision.
- Chest pain, palpitations, oedema (fluid build-up), congestive heart failure (heart disease with shortness of breath and swelling of the feet and legs due to fluid build-up), a type of heart rhythm disorder, heart attack, heart failure.
- Raynaud's phenomenon, cold hands and feet.
- Constriction of the airways in the lungs (predominantly in patients with pre-existing disease)
- Indigestion, diarrhoea, dry mouth, abdominal pain, vomiting.
- Skin rash with white silvery coloured appearance (psoriasisiform rash) or worsening of psoriasis.
- Sexual dysfunction.
- Tiredness.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via

HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2

Tel: +353 1 6764971 Fax: +353 1 6762517

Website: www.hpra.ie E-mail: medsafety@hpra.ie

By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE BETOPTIC 0.5%

- Keep out of the sight and reach of children.
- Do not store above 25°C.
- Keep the bottle tightly closed.
- Keep in the outer carton in order to protect from light.
- Do not use the drops after the expiry date (marked 'Exp') on the bottle and the carton. The expiry date refers to the last day of that month.
- **Stop using the bottle 4 weeks after first opening**, to prevent infections.
- Medicines should not be disposed of via waste water or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Betoptic 0.5% contains

Each ml contains 5mg of the active substance, betaxolol (as hydrochloride).

The other ingredients are: sodium chloride, disodium edetate, benzalkonium chloride, hydrochloric acid and/or sodium hydroxide (to adjust pH) and purified water.

What Betoptic 0.5% looks like and contents of the pack

Betoptic 0.5% is a clear, colourless to pale yellow, sterile eye drops solution supplied in a plastic bottle with a screw cap.

Each bottle contains 5ml of solution.

Manufacturer

Manufactured by: SA Alcon-Couvreur NV, Rijksweg 14, B-2870 Puurs, Belgium.

Procured from within the EU and repackaged by:

Doncaster Pharmaceuticals Group Ltd, Kirk Sandall, Doncaster, DN3 1QR, UK.

PPA holder: Imbat Ltd., Santry, Dublin 9.

Distributed by: Eurodrug Ltd., Santry, Dublin 9.

PPA1151/170/001

Leaflet issue and revision date (ref): 19.07.18

BETOPTIC® is a registered trademark of Novartis AG.