

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Accupro 5 mg Film - Coated tablets.

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains:

Quinapril hydrochloride 5.416 mg (equivalent to 5 mg quinapril base).

Excipients:

Each 5 mg tablet contains Lactose.

For a full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Film-coated tablet.

Product imported from Italy:

Reddish-brown oval film-coated tablets scored on both sides and "5" embossed on both sides.

The scoreline is to allow breaking for ease of swallowing and not to divide into equal halves.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

1. For the treatment of all grades of essential hypertension. Accupro is effective as monotherapy or concomitantly with diuretics in patients with hypertension.
2. For the treatment of congestive heart failure when given concomitantly with a diuretic and/or cardiac glycoside. Treatment of congestive heart failure with Accupro should always be initiated under close medical supervision.

4.2 Posology and method of administration

For oral use.

Adults

Hypertension.

Monotherapy: The recommended initial dosage is 10 mg once daily. Depending upon clinical response, patient's dosage may be titrated (by doubling the dose, allowing adequate time for dosage adjustment) to a maintenance dosage of 20 to 40 mg/day given as a single dose or divided into 2 doses. Long-term control is maintained in most patients with a single daily dosage regimen. Patients have been treated with dosages up to 80 mg/day.

Concomitant Diuretics: In order to determine if excess hypotension will occur, an initial dosage of 5 mg of Accupro is recommended in patients who are also being treated with a diuretic. After this the dosage of Accupro should be titrated (by doubling the dose allowing adequate time for dosage adjustment) to the optimal response (see section 4.5).

Congestive Heart Failure

In order to closely monitor patients for symptomatic hypotension, a single 5 mg initial dosage is recommended. After this, patients should be titrated to an effective dose: (up to 40 mg/day) given in 1 or 2 doses with concomitant diuretic and/or cardiac glycoside therapy. Patients are usually maintained effectively on doses of 10-20 mg/day given with

concomitant therapy.

In the treatment of severe or unstable congestive heart failure, Accupro should always be initiated in hospital under close medical supervision.

Elderly

Age alone does not appear to affect the efficacy or safety profile of quinapril. Therefore, the recommended initial dosage in hypertension of quinapril in elderly patients is 10mg given once daily followed by titration to the optimal response.

Paediatric population

Currently available data are described in sections 5.1 and 5.2 but no recommendation on a posology can be made.

Patients with renal insufficiency

In patients with a creatinine clearance of less than 60 ml/min, an initial dosage in essential hypertension of 5 mg once daily is recommended followed by titration to the optimal response. Kinetic data indicate that the apparent elimination half-life of quinaprilat increases as creatinine clearance decreases (see section 4.4).

4.3 Contraindications

Hypersensitivity to quinapril or any of the excipients listed in section 6.1

Accupro is contraindicated in the second and third trimesters of pregnancy (see sections 4.4 and 4.6).

Use in patients with a history of angioneurotic oedema relating to previous treatment with an ACE inhibitor.

Hereditary or idiopathic angioneurotic oedema.

Quinapril should not be used in patients with dynamic left ventricular outflow obstruction.

4.4 Special warnings and precautions for use

Quinapril should be used with caution in selected patients with aortic stenosis.

Sensitivity reactions:

Sensitivity reactions may occur in patients with or without a history of allergy or bronchial asthma, e.g., purpura, photosensitivity, urticaria, necrotising angitis, respiratory distress including pneumonitis and pulmonary oedema, anaphylactic reactions.

Impaired Renal Function: In patients with renal insufficiency monitoring of renal function during therapy should be performed as deemed appropriate, although in the majority renal function will not alter or may improve.

As a consequence of inhibiting the renin-angiotensin-aldosterone system, changes in renal function may be anticipated in susceptible individuals. In patients with severe heart failure whose renal function may depend on the activity of the renin-angiotensin-aldosterone system, treatment with ACE inhibitors including quinapril, may be associated with oliguria and/or progressive azotemia and rarely acute renal failure and/or death.

The half-life of quinaprilat is prolonged as creatinine clearance falls. Patients with a creatinine clearance of <60 ml/min require a lower initial dosage of quinapril (see section 4.2). These patients' dosage should be titrated upwards based upon therapeutic response, and renal function should be closely monitored although initial studies do not indicate that quinapril produces further deterioration in renal function.

In clinical studies in hypertensive patients with unilateral or bilateral renal artery stenosis, increases in blood urea

nitrogen and serum creatinine have been observed in some patients following ACE inhibitor therapy. These increases were almost always reversible upon discontinuation of the ACE inhibitor and/or diuretic therapy. In such patients, renal function should be monitored during the first few weeks of therapy.

Some patients with hypertension or heart failure with no apparent pre-existing renal vascular disease have developed increases (>1.25 times the upper limit of normal) in blood urea nitrogen and serum creatinine, usually minor and transient, especially when quinapril has been given concomitantly with a diuretic. Increases in blood urea nitrogen and serum creatinine have been observed in 2% and 2%, respectively of hypertensive patients on quinapril monotherapy and in 4% and 3%, respectively of hypertensive patients on quinapril/HCTZ. These increases are more likely to occur in patients with pre-existing renal impairment. Dosage reduction and/or discontinuation of a diuretic and/or quinapril may be required.

There is insufficient experience in patients with severe renal impairment (creatinine clearance <10 ml/min). Treatment is therefore not recommended in these patients.

Impaired Hepatic Function: Quinapril when combined with a diuretic should be used with caution in patients with impaired hepatic function or progressive liver disease, since minor alterations of fluid and electrolyte balance may precipitate hepatic coma. The metabolism of quinapril to quinaprilat is normally dependent upon hepatic esterase. Quinaprilat concentrations are reduced in patients with alcoholic cirrhosis due to impaired deesterification of quinapril.

Rarely, ACE inhibitors have been associated with a syndrome beginning as a cholestatic jaundice and progressing to a fulminant hepatic necrosis (in some cases fatal). Patients who during ACE inhibitor therapy experience jaundice or clearly elevated hepatic enzymes should discontinue quinapril and receive appropriate medical follow-up.

Anaphylactoid reactions

Desensitisation: Patients receiving ACE inhibitors during desensitising treatment with hymenoptera venom have sustained life-threatening anaphylactoid reactions. In the same patients these reactions have been avoided when ACE inhibitors were temporarily withheld, but they have reappeared upon inadvertent rechallenge.

Haemodialysis: Patients haemodialysed using high-flux polyacrylonitrile ('AN69') membranes are highly likely to experience anaphylactoid reactions if they are treated with ACE inhibitors. This combination should therefore be avoided, either by use of alternative antihypertensive drugs or alternative membranes for haemodialysis.

LDL apheresis: Patients undergoing low-density lipoprotein apheresis with dextran-sulphate absorption when treated concomitantly with an ACE inhibitor have reported anaphylactoid reactions. This method should therefore not be used in patients treated with ACE inhibitors.

Angioneurotic oedema: Angioneurotic oedema has been reported in patients treated with ACE inhibitors including Accupro. If laryngeal stridor or angioedema of the face, tongue, or glottis occurs, treatment should be discontinued immediately, the patient treated appropriately in accordance with accepted medical care, and carefully observed until the swelling disappears. Where swelling is confined to the face, lips and mouth, the condition will usually resolve without further treatment, although antihistamines may be useful in relieving symptoms. Angioedema associated with laryngeal involvement may be fatal. These patients should be followed carefully until the swelling has resolved. However, where there is involvement of the tongue, glottis or larynx, likely to cause airways obstruction, appropriate therapy such as subcutaneous adrenaline 1:1000 (0.3 to 0.5ml) should be administered promptly when indicated.

Patients with a history of angioedema unrelated to ACE inhibitor therapy may be at increased risk of angioedema while receiving an ACE inhibitor (see Section 4.3).

Ethnic Differences:

Black patients receiving ACE inhibitor therapy have been shown to have a higher incidence of angioedema compared to non-black patients.

It should also be noted that in controlled clinical trials, ACE inhibitors have an effect on blood pressure that is less in black patients than in non-blacks.

Intestinal angioedema: Intestinal angioedema has been reported in patients treated with ACE inhibitors. These patients

presented with abdominal pain (with or without nausea or vomiting); in some cases there was no prior history of facial angioedema and C-1 esterase levels were normal. The angioedema was diagnosed by procedures including abdominal CT scan or ultrasound, or at surgery, and symptoms resolved after stopping the ACE inhibitor. Intestinal angioedema should be included in the differential diagnosis of patients on ACE inhibitors presenting with abdominal pain.

Caution should be exercised in those known to be hypersensitive to other ACE inhibitors, and particularly those with obstructive airways disease. Patients with a history of angioedema unrelated to ACE inhibitor therapy may be at increased risk of angioedema while receiving an ACE inhibitor (see also section 4.3).

Other hypersensitivity reactions have been reported.

Hypotension: Symptomatic hypotension is rarely seen in uncomplicated hypertensive patients treated with Accupro but it is a possible consequence of ACE inhibition therapy particularly in salt/volume depleted patients such as those previously treated with diuretics, who have a dietary salt reduction, who are on dialysis, have diarrhoea or vomiting, or has severe renin-dependent hypertension. (see sections 4.5 and 4.8). Any electrolyte or fluid inadequacy should be corrected preferably before initial dose of the product. Careful medical supervision is necessary for a period after dosing. If symptomatic hypotension occurs, the patient should be placed in the supine position and, if necessary, receive an intravenous infusion of normal saline. A transient hypotensive response is not a contraindication to further doses; however, lower doses of quinapril or any concomitant diuretic therapy should be considered if this event occurs.

In patients with congestive heart failure, who are at risk of excessive hypotension, quinapril therapy should be started at the recommended dose under close medical supervision; these patients should be followed closely for the first two weeks of treatment and whenever the dosage of quinapril is increased.

Similar considerations apply to patients with ischaemic heart or cerebrovascular disease in whom an excessive fall in blood pressure could result in a myocardial infarction or cerebrovascular accident.

Neutropenia/agranulocytosis: ACE inhibitors have been rarely associated with agranulocytosis and bone marrow depression in patients with uncomplicated hypertension but more frequently in patients with renal impairment, especially if they also have collagen vascular disease. Agranulocytosis has been rarely reported during treatment with quinapril.

As with other ACE inhibitors, monitoring of white blood cell counts in patients with collagen vascular disease and/or renal diseases should be considered.

Hypoglycaemia: ACE inhibitors may enhance insulin sensitivity in diabetic patients and have been associated with hypoglycaemia in patients on insulin or oral hypoglycaemic agents. Glycaemic control should be closely monitored particularly during the first month of treatment with an ACE inhibitor (see Section 4.5).

Cough: Cough has been reported with the use of ACE inhibitors including quinapril. Characteristically, the cough is non-productive, persistent and resolves after discontinuation of therapy. ACE inhibitor-induced cough should be considered as part of the differential diagnosis of cough.

Surgery/Anaesthesia:

In patients undergoing major surgery or during anaesthesia with agents that produce hypotension, quinapril may block angiotensin II formation secondary to compensatory renin release. If hypotension occurs and is considered to be due to this mechanism, it can be corrected by volume expansion (see Section 4.5).

Hyperkalemia and Potassium-sparing Diuretics:

Patients on quinapril alone may have increased serum potassium levels. When administered concomitantly, quinapril may reduce the hypokalemia induced by thiazide diuretics. Because of the risk of further potentiating increases in serum potassium it is advised that combination therapy with potassium-sparing diuretics be initiated with caution and the patient's serum potassium levels be closely monitored (see Hypotension above and Section 4.5 Interaction with other medicinal products and other forms of interaction).

Patients with rare hereditary problems with galactose intolerance, the Lapp lactase deficiency or glucose-galactose

malabsorption should not take this medicine.

Pregnancy:

ACE inhibitors should not be initiated during pregnancy. Unless continued ACE inhibitor therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with ACE inhibitors should be stopped immediately, and, if appropriate, alternative therapy should be started (see sections 4.3 and 4.6).

4.5 Interaction with other medicinal products and other forms of interaction

Tetracycline and other drugs that interact with magnesium: Because of the presence of magnesium carbonate in the formulation Accupro has been shown in healthy volunteers to reduce the absorption of tetracycline in concomitant administration by 28-37%. This interaction should be considered if coprescribing quinapril and tetracycline. It is recommended that concomitant administration of tetracycline be avoided.

Concomitant diuretic therapy: Patients treated with diuretics may occasionally experience an excessive reduction of blood pressure after initiation of therapy with Accupro. This hypotensive effect may be effectively minimised by either discontinuing the diuretic or increasing the salt intake prior to the initial dose of Accupro. If discontinuation of the diuretic is not possible, the starting dose of quinapril should be reduced. In patients in whom a diuretic is continued, medical supervision should be provided for up to two hours following administration of the initial dose (see Section 4.4 and section 4.2).

Other anti-hypertensive agents: B-blockers, methyldopa and diuretics may enhance the hypotensive effects of quinapril, and should only be used under careful supervision. Concomitant propranolol did not affect the pharmacokinetics of quinapril in a single dose study.

Calcium antagonists: There is no experience of concomitant use with Accupro.

Atorvastatin: Co-administration of multiple 10mg doses of atorvastatin with 80mg quinapril resulted in no significant change in the steady state pharmacokinetic parameters of atorvastatin.

Lithium: Increased serum lithium levels and symptoms of lithium toxicity have been reported in patients receiving concomitant lithium and ACE inhibitor therapy due to the sodium-losing effect of these agents. These drugs should be co-administered with caution and frequent monitoring of serum lithium levels is recommended. If a diuretic is also used, it may increase the risk of lithium toxicity.

Non-steroidal anti-inflammatory agents including selective cyclooxygenase-2 inhibitors (COX-2 inhibitors): In patients who are elderly, volume-depleted (including those on diuretic therapy), or with compromised renal function, co-administration of NSAIDs, including selective COX-2 inhibitors, with ACE inhibitors, including quinapril, may result in deterioration of renal function, including possible acute renal failure. These effects are usually reversible. Monitor renal function periodically in patients receiving quinapril and NSAID therapy.

The antihypertensive effect of ACE inhibitors, including quinapril may be attenuated by NSAIDs. It has been described that NSAIDs and ACE inhibitors exert an additive effect on the increase in serum potassium.

Gold:

Nitritoid reactions (symptoms include facial flushing, nausea, vomiting, and hypotension) have been reported rarely in patients on therapy with injectable gold (sodium aurothiomalate) and concomitant ACE inhibitor therapy.

Allopurinol, cytostatic and immunosuppressive agents, systemic corticosteroids or procainamide:

Concomitant administration with ACE inhibitors may lead to an increased risk for leucopenia (See section 4.4).

Alcohol, barbiturates and narcotics:

Potential of orthostatic hypotension may occur.

Agents increasing serum potassium: Quinapril is an angiotensin-converting enzyme inhibitor capable of lowering aldosterone levels, which in turn can result in elevation in serum potassium. Concomitant treatments with potassium

sparing diuretics, potassium supplements or potassium salts should only be used with caution and with appropriate monitoring of serum potassium, especially in patients with impaired renal function.

Surgery/anaesthesia: Although no data are available to indicate there is an interaction between Accupro and anaesthetic agents that produces hypotension, caution should be exercised when patients undergo major surgery or anaesthesia since angiotensin converting enzyme inhibitors have been shown to block angiotensin II formation secondary to compensatory renin release. This may lead to hypotension which can be corrected by volume expansion.(See Section 4.4)

Antacids

Antacids may decrease the bioavailability of quinapril

Antidiabetic drugs: Concomitant administration of ACE inhibitors and anti-diabetic medicines (insulin, oral hypoglycaemic agents) may cause an increased blood glucose lowering effect with the risk of hypoglycaemia. This phenomenon may be more likely to occur during the first weeks of combined treatment and in patients with renal impairment. Glycaemic control should be closely monitored particularly during the first month of treatment with an ACE inhibitor (see section 4.4).

4.6 Fertility, pregnancy and lactation

Pregnancy:

The use of ACE inhibitors is not recommended during the first trimester of pregnancy (see section 4.4). The use of ACE inhibitors is contraindicated during the 2nd and 3rd trimester of pregnancy (see sections 4.3 and 4.4).

Epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded. Unless continued ACE inhibitor therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with ACE inhibitors should be stopped immediately, and, if appropriate, alternative therapy should be started.

Exposure to ACE inhibitor therapy during the second and third trimesters is known to induce human foetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia). Should exposure to ACE inhibitor have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended. Infants whose mothers have taken ACE inhibitors should be closely observed for hypotension (see sections 4.3 and 4.4). oliguria, and hyperkalaemia. If oliguria occurs, attention should be directed toward support of blood pressure and renal perfusion.

Lactation:

Limited pharmacokinetic data demonstrate very low concentrations in breast milk (see section 5.2). Although these concentrations seem to be clinically irrelevant, the use of Accupro in breastfeeding is not recommended for preterm infants and for the first few weeks after delivery, because of the hypothetical risk of cardiovascular and renal effects and because there is not enough clinical experience.

In the case of an older infant, the use of Accupro in a breast-feeding mother may be considered if this treatment is necessary for the mother and the child is observed for any adverse effect.

4.7 Effects on ability to drive and use machines

The ability to engage in activities such as operating machinery or operating a motor vehicle may be impaired especially when initiating quinapril therapy.

4.8 Undesirable effects

- Very common (≥1/10)
- Common (≥1/100 to <1/10)
- Uncommon (≥1/1,000 to <1/100)
- Rare (≥1/10,000 to <1/1,000)
- Very rare (≤1/10,000)
- Not known (cannot be estimated from the available data)

The most frequently reported adverse reactions found in controlled clinical trials were headache (7.2%), dizziness (5.5%), cough (3.9%), fatigue (3.5%), rhinitis (3.2%), nausea and/or vomiting (2.8%), and myalgia (2.2%).

System Organ Class	Frequency	Undesirable effects
Blood and lymphatic system disorders	Not Known	Agranulocytosis, haemolytic anaemia, neutropenia, thrombocytopenia
Immune system disorders	Not Known	Anaphylactoid reaction
Metabolism and nutrition disorders	Common	Hyperkalaemia
Psychiatric disorders	Common	Insomnia
	Uncommon	Confusional state, depression, nervousness
Nervous system disorders	Common	Dizziness, headache, paraesthesia
	Uncommon	Transient ischaemic attack, somnolence
	Rare	Balance disorder, syncope
	Not known	Cerebrovascular accident
Eye disorders	Uncommon	Amblyopia
	Very Rare	Vision blurred
Ear and labyrinth disorders	Uncommon	Vertigo, tinnitus
Cardiac disorders	Uncommon	Myocardial infarction, angina pectoris, tachycardia, palpitations
Vascular disorders	Common	Hypotension
	Uncommon	Vasodilatation
	Not known	Orthostatic hypotension
Respiratory, thoracic and mediastinal disorders	Common	Dyspnoea, cough
	Uncommon	Dry throat
	Rare	Eosinophilic pneumonia
	Not known	Bronchospasm. In individual cases, upper airways obstruction by angioedema (that may be fatal)
Gastrointestinal disorders	Common	Vomiting, diarrhoea, dyspepsia, abdominal pain, nausea
	Uncommon	Flatulence, dry mouth

	Rare	Glossitis, constipation, dysgeusia
	Very Rare	Ileus, small bowel angioedema
	Not Known	Pancreatitis*
Hepato-biliary disorders	Not Known	Hepatitis, jaundice cholestatic
Skin and subcutaneous tissue disorders	Uncommon	Angioedema, rash, pruritus, hyperhidrosis
	Rare	Erythema multiforme, pemphigus, urticaria
	Very Rare	Dermatitis psoriasiform
	Not Known	Stevens Johnson syndrome, toxic epidermal necrolysis, exfoliative dermatitis, alopecia, photosensitivity reaction. Skin disorders may be associated with pyrexia, muscle and joint pain (myalgia, arthralgia, arthritis), vascular inflammation (vasculitis), inflammation of serous tissues and certain changes in laboratory values (eosinophilia, leukocytosis and/or antinuclear antibody increased, red blood sedimentation rate increased).
Musculoskeletal, connective tissue and bone disorders	Common	Back pain, myalgia
Renal and urinary disorders	Uncommon	Renal impairment, proteinuria
Reproductive system and breast disorders	Uncommon	Erectile dysfunction
General disorders and administration site conditions	Common	Fatigue, asthenia, chest pain
	Uncommon	Generalised oedema, pyrexia, oedema peripheral
Investigations	Common	Blood creatinine increased, blood urea increased**
	Not Known	Haemoglobin decreased, haematocrit decreased, decreases in haematocrit and WCXC, hepatic enzyme increased, blood bilirubin increased. In patients with a congenital G-6-PDH deficiency, individual cases of haemolytic anaemia have been reported.
Infections and infestations	Common	Pharyngitis, rhinitis
	Uncommon	Bronchitis, upper respiratory tract infection, urinary tract infection, sinusitis

* Pancreatitis has been reported rarely in patients treated with ACE inhibitors; in some cases this has proved fatal.
 ** Such increases are more likely to occur in patients receiving concomitant diuretic therapy than those on monotherapy with quinapril. These observed increases will often reverse on continued therapy.

Vasculitis and gynecomastia have been reported with other ACE inhibitors and it cannot be excluded that these unwanted effects are class specific.

4.9 Overdose

The oral LD50 of quinapril in mice and rats ranges from 1440 to 4280 mg/kg.

No data are available with respect to overdosage in humans. The most likely clinical manifestation would be symptoms attributable to severe hypotension, which should normally be treated by intravenous volume expansion.

Treatment is symptomatic and supportive, consistent with established medical care.

Haemodialysis and peritoneal dialysis have little effect on the elimination of quinapril and quinaprilat.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

ATC Code – CO9AA06, ACE inhibitors, plain.

Accupro is rapidly deesterified to quinaprilat (quinapril diacid, the principal metabolite) which, in human and animal studies, is a potent angiotensin-converting enzyme (ACE) inhibitor. The primary mode of action of Accupro in humans and animals is to inhibit ACE, thereby decreasing vasopressor activity and aldosterone secretion. Removal of angiotensin II negative feedback on renin secretion leads to increased plasma renin activity. Accupro has antihypertensive activity in the presence of low to normal plasma renin concentrations.

Other possible mechanisms contributing to the activity of ACE inhibitors include bradykinin-induced vasodilation, release of prostaglandins, attenuation of sympathetic nervous system activity, and inhibition of tissue enzyme-converting activity.

Administration of 10-40 mg of quinapril to patients with mild to moderate hypertension results in a reduction of both sitting and standing blood pressure with minimal effect on heart rate. Antihypertensive activity commences within one hour with peak effects usually achieved by two to four hours after dosing. Achievement of maximum blood pressure lowering effects may require two weeks of therapy in some patients. At the recommended doses, antihypertensive effects are maintained in most patients throughout the 24 hour dosing interval and continue during long term therapy.

In a randomised clinical trial using target doses of 2.5, 5, 10 and 20 mg of quinapril, in 112 children and adolescents with hypertension or high normal blood pressure over 8 weeks (2 weeks double blind and 6 weeks extension) failed to reach its primary objective of reduction of diastolic blood pressure after 2 weeks. For systolic blood pressure (secondary objective of efficacy) at Week 2 only there was a statistically significant linear dose response across treatments with a significant difference between the quinapril 20 mg QD and placebo treatment groups.

Long term effects of quinapril on growth, puberty and general development have not been studied.

5.2 Pharmacokinetic properties

Peak plasma Accupro concentrations are observed within 1 hour of oral administration. The extent of absorption is approximately 60%, and is not influenced by food. Following absorption, Accupro is deesterified to its major active metabolite, quinaprilat, and to minor inactive metabolites. Accupro has an apparent half-life of approximately one hour. Peak plasma quinaprilat concentrations are observed approximately 2 hours following an oral dose of quinapril.

Quinaprilat is eliminated primarily by renal excretion and has an effective accumulation half-life of 3 hours. In

patients with renal insufficiency and creatinine clearance of ≤ 40 ml/min, peak and trough quinaprilat concentrations increase, time to peak concentration increases, apparent half-life increases, and time to steady state may be delayed. The elimination of quinaprilat is also reduced in elderly patients (>65 years) and correlates well with the impaired renal function which frequently occurs in the elderly. Studies in rats indicate that Accupro and its metabolites do not cross the blood-brain barrier.

Lactation:

After a single oral dose of 20 mg of quinapril in six breast-feeding women, the M/P (milk to plasma ratio) for quinapril was 0.12. Quinapril was not detected in milk after 4 hours after the dose. Quinaprilat milk levels were undetectable (<5 $\mu\text{g/L}$) at all time points. It is estimated that a breastfed infant would receive about 1.6% of the maternal weight-adjusted dosage of quinapril.

The pharmacokinetics of quinapril has been studied in a single dose study (0.2 mg/kg) in 24 children aged 2.5 months to 6.8 years and a multiple dose study (0.016-0.468 mg/kg) in 38 children aged 5-16 years old, weighing 66-98 kg on average.

As in adults, quinapril was rapidly converted to quinaprilat. Quinaprilat concentrations generally peaked 1 to 2 hours post dose and declined with a mean half-life of 2.3 hours. In infants and young children the exposure following a single 0.2-mg/kg dose is comparable to that observed in adults after a single 10-mg dose. In a multiple dose study in school age and adolescents, the AUC and C_{max} values of quinaprilat were observed to increase linearly with increasing dose of quinapril on a mg/kg basis."

5.3 Preclinical safety data

The results of the preclinical tests do not add anything of further significance to the prescriber.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Heavy magnesium carbonate
Lactose
Gelatin
Crospovidone
Magnesium stearate
Candelilla wax
Hypromellose
Hydroxypropylcellulose
Macrogol 400
Red iron oxide (E172)
Titanium dioxide (E171)

6.2 Incompatibilities

Not applicable

6.3 Shelf life

The shelf-life expiry date of this product shall be the date shown on the container and outer package of the product on the market in the country of origin.

6.4 Special precautions for storage

Do not store above 25°C

6.5 Nature and contents of container

Polyamide/aluminium/PVC blister containing 28 film-coated tablets in an over-labelled outer carton.

6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product

No special requirements

7 PARALLEL PRODUCT AUTHORISATION HOLDER

B&S Healthcare
Unit 4
Bradfield Road
Ruislip
Middlesex
HA4 0NU
United Kingdom

8 PARALLEL PRODUCT AUTHORISATION NUMBER

PPA 1328/045/002

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of First Authorisation: 8th December 2006

Date of Last Renewal: 8th December 2011

10 DATE OF REVISION OF THE TEXT

January 2014