

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Omnexel 400 microgram Prolonged Release Tablets, Film-coated.

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each prolonged release film-coated tablet contains 0.4 mg tamsulosin hydrochloride.

Excipients:

For a full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Film-coated, Prolonged Release Tablets
(Oral Controlled Absorption System)

Product imported from the Netherlands:

Approximately 9 mm in diameter, round, bi-convex, yellow, film-coated tablets, debossed with the code '04'.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Lower urinary tract symptoms (LUTS) associated with benign prostatic hyperplasia (BPH).

4.2 Posology and method of administration

Oral use.

One tablet daily.

Omnexel can be taken independently of food.

The tablet must be swallowed whole and not be crunched or chewed as this interferes with the prolonged release of the active substance.

No dose adjustment is warranted in renal impairment.

No dose adjustment is warranted in patients with mild to moderate hepatic insufficiency (see also 4.3, Contraindications).

There is no relevant indication for use of Omnexel in children.

4.3 Contraindications

Hypersensitivity to tamsulosin hydrochloride, including drug-induced angioedema or to any of the excipients.

A history of orthostatic hypotension.

Severe hepatic insufficiency.

4.4 Special warnings and precautions for use

As with other α_1 -adrenoceptor antagonists, a reduction in blood pressure can occur in individual cases during treatment with Omnexel, as a result of which, rarely, syncope can occur. At the first signs of orthostatic hypotension (dizziness, weakness), the patient should sit or lie down until the symptoms have disappeared.

Before therapy with Omnexel is initiated, the patient should be examined in order to exclude the presence of other conditions, which can cause the same symptoms as benign prostatic hyperplasia.

Digital rectal examination and, when necessary, determination of prostate specific antigen (PSA) should be performed before treatment and at regular intervals afterwards.

The treatment of patients with severe renal impairment (creatinine clearance of < 10 ml/min) should be approached with caution, as these patients have not been studied.

The 'Intraoperative Floppy Iris Syndrome' (IFIS, a variant of small pupil syndrome) has been observed during cataract surgery in some patients on or previously treated with tamsulosin. IFIS may lead to increased procedural complications during the operation. The initiation of therapy with tamsulosin in patients for whom cataract surgery is scheduled is not recommended. Discontinuing tamsulosin 1-2 weeks prior to cataract surgery is anecdotally considered helpful, but the benefit and duration of stopping of therapy prior to cataract surgery has not yet been established.

During pre-operative assessment, cataract surgeons and ophthalmic teams should consider whether patients scheduled for cataract surgery are being or have been treated with tamsulosin in order to ensure that appropriate measures will be in place to manage the IFIS during surgery.

It is possible that a remnant of the tablet is observed in the faeces.

4.5 Interaction with other medicinal products and other forms of interaction

Interaction studies have only been performed in adults.

No interactions have been seen when tamsulosin hydrochloride was given concomitantly with either atenolol, enalapril, nifedipine or theophylline.

Concomitant cimetidine brings about a rise in plasma levels of tamsulosin, whereas furosemide a fall, but as levels remain within the normal range posology need not be adjusted.

In vitro, neither diazepam nor propranolol, trichlormethiazide, chlormadinone, amitriptyline, diclofenac, glibenclamide, simvastatin and warfarin change the free fraction of tamsulosin in human plasma. Neither does tamsulosin change the free fractions of diazepam, propranolol, trichlormethiazide and chlormadinone.

No interactions at the level of hepatic metabolism have been seen during *in vitro* studies with liver microsomal fractions (representative of the cytochrome P₄₅₀-linked drug metabolising enzyme system), involving amitriptyline, salbutamol, glibenclamide and finasteride. Diclofenac and warfarin, however, may increase the elimination rate of tamsulosin.

Concurrent administration of other α_1 -adrenoceptor antagonists could lead to hypotensive effects.

4.6 Fertility, pregnancy and lactation

Not applicable, as Omnexel is intended for male patients only.

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. However, patients should be aware of the fact that dizziness can occur.

4.8 Undesirable effects

MedDRA System organ class	Common (>1/100, <1/10)	Uncommon (>1/1,000, <1/100)	Rare (>1/10,000, <1/1,000)	Very Rare (<1/10,000)
<i>Nervous Systems Disorders</i>	Dizziness (1.3%)	Headache	Syncope	
<i>Cardiac disorders</i>		Palpitations		
<i>Vascular disorders</i>		Postural hypotension		
<i>Respiratory disorders</i>		Rhinitis		
<i>Gastro-intestinal disorders</i>		Constipation, diarrhoea, nausea, vomiting		
<i>Skin and subcutaneous tissue disorders</i>		Rash, pruritus, urticaria	Angioedema	
<i>Reproductive system and breast disorders</i>		Abnormal ejaculation		Priapism
<i>General disorders and administration site conditions</i>		Asthenia		

During cataract surgery a small pupil situation, known as Intraoperative Floppy Iris Syndrome (IFIS), has been associated with therapy of tamsulosin during post-marketing surveillance (*see also section 4.4, Special warnings and precautions for use*).

4.9 Overdose

Acute overdose with 5 mg of tamsulosin hydrochloride has been reported. Acute hypotension (systolic blood pressure 70 mm Hg), vomiting and diarrhoea were observed, which were treated with fluid replacement and the patient could be discharged the same day.

In case of acute hypotension occurring after overdosage cardiovascular support should be given. Blood pressure can be restored and heart rate brought back to normal by lying the patient down. If this does not help then volume expanders and, when necessary, vasopressors could be employed. Renal function should be monitored and general supportive measures applied. Dialysis is unlikely to be of help as tamsulosin is very highly bound to plasma proteins.

Measures, such as emesis, can be taken to impede absorption. When large quantities are involved, gastric lavage can be applied and activated charcoal and an osmotic laxative, such as sodium sulphate, can be administered.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: α_1 -adrenoceptor antagonists.

ATC code: G04C A02. Preparations for the exclusive treatment of prostatic disease.

Mechanism of action

Tamsulosin binds selectively and competitively to the postsynaptic α_1 -adrenoceptors, in particular to subtypes α_{1A} and α_{1D} . It brings about relaxation of prostatic and urethral smooth muscle.

Pharmacodynamic effects

Omnexel increases the maximum urinary flow rate. It relieves obstruction by relaxing smooth muscle in prostate and urethra thereby improving voiding symptoms.

It also improves the storage symptoms in which bladder instability plays an important role.

These effects on storage and voiding symptoms are maintained during long-term therapy. The need for surgery or catheterisation is significantly delayed.

α_1 -adrenoceptor antagonists can reduce blood pressure by lowering peripheral resistance. No reduction in blood pressure of any clinical significance was observed during studies with Omnexel.

5.2 Pharmacokinetic properties

Absorption

Omnexel is a prolonged release tablet of the non-ionic gel matrix type. The Ocas formulation provides consistent slow release of tamsulosin, resulting in an adequate exposure, with little fluctuation, over 24 hours.

Tamsulosin administered as Omnexel mg is absorbed from the intestine. Of the administered dose, approximately 57% is estimated to be absorbed.

The rate and extent of absorption of tamsulosin administered as Omnexel are not affected by food.

Tamsulosin shows linear pharmacokinetics.

After a single dose of Omnexel in the fasted state, plasma concentrations of tamsulosin peak at a median time of 6 hours. In steady state, which is reached by day 4 of multiple dosing, plasma concentrations of tamsulosin peak at 4 to 6 hours, in the fasted and fed state. Peak plasma concentrations increase from approximately 6 ng/ml after the first dose to 11 ng/ml in steady state.

As a result of the prolonged release characteristics of Omnic Ocas the trough concentration of tamsulosin in plasma amounts to 40% of the peak plasma concentration under fasted and fed conditions.

There is a considerable inter-patient variation in plasma levels both after single and multiple dosing.

Distribution

In man, tamsulosin is about 99% bound to plasma proteins. The volume of distribution is small (about 0.2 l/kg).

Biotransformation

Tamsulosin has a low first pass effect, being metabolised slowly. Most tamsulosin is present in plasma in the form of unchanged active substance. It is metabolised in the liver.

In rats, hardly any induction of microsomal liver enzymes was seen to be caused by tamsulosin.

None of the metabolites are more active than the original compound.

Elimination

Tamsulosin and its metabolites are mainly excreted in the urine. The amount excreted as unchanged active substance is estimated to be about 4 - 6% of the dose, administered as Omnexel.

After a single dose of Omnexel mg and in steady state, elimination half-lives of about 19 and 15 hours, respectively, have been measured.

5.3 Preclinical safety data

Single and repeat dose toxicity studies were performed in mice, rats and dogs. In addition, reproduction toxicity in rats, carcinogenicity in mice and rats and *in vivo* and *in vitro* genotoxicity were examined.

The general toxicity profile, as seen with high doses of tamsulosin, is consistent with the known pharmacological actions of the α -adrenoceptor antagonists.

At very high dose levels the ECG was altered in dogs. This response is considered to be not clinically relevant.

Tamsulosin showed no relevant genotoxic properties.

Increased incidences of proliferative changes of mammary glands of female rats and mice have been reported. These findings, which are probably mediated by hyperprolactinemia and only occurred at high dose levels, are regarded as irrelevant.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Macrogol 7.000.000

Macrogol 8.000

Magnesium stearate (E470b)

Butylhydroxytoluene (E321)

Colloidal silica anhydrous (E551)

Hypromellose (E464)

Iron oxide yellow (E172)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

The shelf life expiry date of this product is the date shown on the blister strips and outer carton of the product as marketed in the country of origin.

6.4 Special precautions for storage

Store in the original pack.

6.5 Nature and contents of container

Over-labelled cardboard carton containing three aluminium blister strips (10 tablets per blister).

Pack size: 30 tablets

6.6 Special precautions for disposal and other handling

No special requirements.

7 PARALLEL PRODUCT AUTHORISATION HOLDER

Profind Wholesale Ltd
Unit 625, Kilshane Avenue
Northwest Business Park
Ballycoolin
Dublin 15
Ireland

8 PARALLEL PRODUCT AUTHORISATION NUMBER

PPA1500/80/1

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of First Authorisation: 3rd May 2011

10 DATE OF REVISION OF THE TEXT