

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

COSOPT Preservative-Free 20 mg/ml + 5 mg/ml, eye drops, solution, single-dose container

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml contains 22.26 mg of dorzolamide hydrochloride corresponding to 20 mg dorzolamide and 6.83 mg of timolol maleate corresponding to 5 mg timolol.

For a full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Eye drops, solution, single dose container

Clear, colourless to nearly colourless, slightly viscous solution.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Indicated in the treatment of elevated intraocular pressure (IOP) in patients with open-angle glaucoma or pseudoexfoliative glaucoma when topical beta-blocker monotherapy is not sufficient.

4.2 Posology and method of administration

The dose is one drop of COSOPT Preservative-Free in the (conjunctival sac of the) affected eye(s) two times daily.

If another topical ophthalmic agent is being used, COSOPT Preservative-Free and the other agent should be administered at least ten minutes apart.

This medicinal product is a sterile solution that does not contain a preservative. The solution from one individual single dose container is to be used immediately after opening for administration to the affected eye(s). Since sterility cannot be maintained after the individual single dose container is opened, any remaining contents must be discarded immediately after administration.

Patients should be instructed to avoid allowing the tip of the container to come into contact with the eye or surrounding structures.

Patients should also be instructed that ocular solutions, if handled improperly, can become contaminated by common bacteria known to cause ocular infections. Serious damage to the eye and subsequent loss of vision may result from using contaminated solutions.

Usage Instructions

1. Open the sachet which contains 15 individual single dose containers. There are three strips of 5 single dose containers each in the sachet.
2. First wash your hands then break off one single dose container from the strip and twist open the top.
3. Tilt your head back and pull your lower eyelid down slightly to form a pocket between your eyelid and eye.

4. Instill one drop in the affected eye(s) as directed by your physician. Each single dose container contains enough solution for both eyes.
5. When using nasolacrimal occlusion or closing the eyelids for 2 minutes, the systemic absorption is reduced. This may result in a decrease in systemic side effects and an increase in local activity.
6. Close your eye for a moment and press the inner corner of the eye with your finger for about one minute. This helps to prevent the eye drop from draining down the tear duct.
7. After instillation, discard the used single dose container even if there is solution remaining.
8. Store the remaining single dose containers in the sachet; the remaining single dose containers must be used within 15 days after opening of the sachet.

Pediatric population

Efficacy in paediatric patients has not been established.

Safety in paediatric patients below the age of 2 years has not been established. (For information regarding safety in paediatric patients ≥ 2 and <6 years of age, see section 5.1) .

4.3 Contraindications

COSOPT Preservative-Free is contraindicated in patients with:

- reactive airway disease, including bronchial asthma or a history of bronchial asthma, or severe chronic obstructive pulmonary disease
- sinus bradycardia, sick sinus syndrome, sino-atrial block, second or third degree atrioventricular block not controlled with pacemaker, overt cardiac failure, cardiogenic shock
- severe renal impairment ($\text{CrCl} < 30 \text{ ml/min}$) or hyperchloremic acidosis
- hypersensitivity to one or both active substances or to any of the excipients.

The above are based on the components and are not unique to the combination.

4.4 Special warnings and precautions for use

Cardiovascular/Respiratory Reactions

Like other topically applied ophthalmic agents timolol is absorbed systemically. Due to betaadrenergic component, timolol, the same types of cardiovascular, pulmonary and other adverse reactions seen with systemic beta-adrenergic blocking agents may occur. Incidence of systemic ADRs after topical ophthalmic administration is lower than for systemic administration. To reduce the systemic absorption, see 4.2.

Cardiac disorders:

In patients with cardiovascular diseases (e.g. coronary heart disease, Prinzmetal's angina and cardiac failure) and hypotension therapy with beta-blockers should be critically assessed and the therapy with other active substances should be considered. Patients with cardiovascular diseases should be watched for signs of deterioration of these diseases and of adverse reactions.

Due to its negative effect on conduction time, betablockers should only be given with caution to patients with first degree heart block.

Vascular disorders

Patients with severe peripheral circulatory disturbance/disorders (i.e. severe forms of Raynaud's

disease or Raynaud's syndrome) should be treated with caution.

Respiratory disorders:

Respiratory reactions, including death due to bronchospasm in patients with asthma have been reported following administration of some ophthalmic beta-blockers.

Cosopt preservative-free should be used with caution, in patients with mild/moderate chronic obstructive pulmonary disease (COPD) and only if the potential benefit outweighs the potential risk.

Hepatic Impairment

This medicinal product has not been studied in patients with hepatic impairment and should therefore be used with caution in such patients.

Immunology and Hypersensitivity

As with other topically-applied ophthalmic agents, this medicinal product may be absorbed systemically. Dorzolamide contains a sulfonamido group, which also occurs in sulfonamides. Therefore, the same types of adverse reactions found with systemic administration of sulfonamides may occur with topical administration, including severe reactions such as Stevens-Johnson syndrome and toxic epidermal necrolysis. If signs of serious reactions or hypersensitivity occur, discontinue use of this preparation.

Local ocular adverse effects, similar to those observed with dorzolamide hydrochloride eye drops, have been seen with this medicinal product. If such reactions occur, discontinuation of COSOPT Preservative-Free should be considered.

While taking beta-blockers, patients with a history of atopy or a history of severe anaphylactic reaction to a variety of allergens may be more reactive to repeated challenge with such allergens and may be unresponsive to the usual doses of adrenaline used to treat anaphylactic reactions.

Concomitant Therapy

The effect on intra-ocular pressure or the known effects of systemic beta-blockade may be potentiated when timolol is given to the patients already receiving a systemic beta-blocking agent. The response of these patients should be closely observed. The use of two topical beta-adrenergic blocking agents is not recommended (see section 4.5).

The use of dorzolamide and oral carbonic anhydrase inhibitors is not recommended.

Withdrawal of Therapy

As with systemic beta-blockers, if discontinuation of ophthalmic timolol is needed in patients with coronary heart disease, therapy should be withdrawn gradually.

Additional Effects of Beta-Blockade

Hypoglycaemia/diabetes

Beta-blockers should be administered with caution in patients subject to spontaneous hypoglycaemia or to patients with labile diabetes, as beta-blockers may mask the signs and symptoms of acute hypoglycaemia.

Beta-blockers may also mask the signs of hyperthyroidism. Abrupt withdrawal of beta-blocker therapy may precipitate a worsening of symptoms.

Corneal diseases

Ophthalmic β -blockers may induce dryness of eyes. Patients with corneal diseases should be treated with caution.

Surgical anaesthesia

Beta-blocking ophthalmological preparations may block systemic beta-agonist effects e.g. of adrenaline. The anaesthesiologist should be informed when the patient is receiving timolol.

Therapy with beta-blockers may aggravate symptoms of myasthenia gravis.

Additional Effects of Carbonic Anhydrase Inhibition

Therapy with oral carbonic anhydrase inhibitors has been associated with urolithiasis as a result of acid-base disturbances, especially in patients with a prior history of renal calculi. Although no acid-base disturbances have been observed with COSOPT (preserved formulation), urolithiasis has been reported infrequently. Because COSOPT Preservative-Free contains a topical carbonic anhydrase inhibitor that is absorbed systemically, patients with a prior history of renal calculi may be at increased risk of urolithiasis while using this medicinal product.

Other

The management of patients with acute angle-closure glaucoma requires therapeutic interventions in addition to ocular hypotensive agents. This medicinal product has not been studied in patients with acute angle-closure glaucoma.

Corneal oedema and irreversible corneal decompensation have been reported in patients with pre-existing chronic corneal defects and/or a history of intraocular surgery while using dorzolamide. There is an increased potential for developing corneal oedema in patients with low endothelial cell counts. Precautions should be used when prescribing COSOPT Preservative-Free to these groups of patients.

Choroidal detachment has been reported with administration of aqueous suppressant therapies (e.g. timolol, acetazolamide) after filtration procedures.

As with the use of other antiglaucoma medicines, diminished responsiveness to ophthalmic timolol maleate after prolonged therapy has been reported in some patients. However, in clinical studies in which 164 patients have been followed for at least three years, no significant difference in mean intraocular pressure has been observed after initial stabilization.

Contact Lens Use

This medicinal product has not been studied in patients wearing contact lenses.

Paediatric population

See section 5.1.

4.5 Interaction with other medicinal products and other forms of interaction

Specific medicine interaction studies have not been performed with Cosopt Preservative-Free.

In a clinical study, this medicinal product was used concomitantly with the following systemic medications without evidence of adverse interactions: ACE-inhibitors, calcium channel blockers, diuretics, non-steroidal anti-inflammatory medicines including aspirin, and hormones (e.g., estrogen, insulin, thyroxine).

There is a potential for additive effects resulting in hypotension and/or marked bradycardia when ophthalmic beta-blockers solution is administered concomitantly with oral calcium channel blockers, catecholamine-depleting medicines or beta-adrenergic blocking agents, antiarrhythmics (including amiodarone), digitalis glycosides, parasympathomimetics, guanethidine, narcotics, and monoamine oxidase (MAO) inhibitors.

Potentiated systemic beta-blockade (e.g., decreased heart rate, depression) has been reported during combined treatment with CYP2D6 inhibitors (e.g. quinidine, fluoxetine, paroxetine) and timolol.

Although COSOPT (preserved formulation) alone has little or no effect on pupil size, mydriasis resulting from

concomitant use of ophthalmic beta-blockers and adrenaline (epinephrine) has been reported occasionally

Beta-blockers may increase the hypoglycemic effect of antidiabetic agents.

Oral beta-adrenergic blocking agents may exacerbate the rebound hypertension which can follow the withdrawal of clonidine.

4.6 Fertility, pregnancy and lactation

Use During Pregnancy

Cosopt preservative-Free should not be used during pregnancy.

Dorzolamide

No adequate clinical data in exposed pregnancies are available. In rabbits, dorzolamide produced teratogenic effect at maternotoxic doses (see Section 5.3).

Timolol

There are no adequate data for the use of timolol in pregnant women. Timolol should not be used during pregnancy unless clearly necessary. To reduce the systemic absorption, see section 4.2.

Epidemiological studies have not revealed malformative effects but show a risk for intra uterine growth retardation when betablockers are administered by the oral route. In addition, signs and symptoms of beta-blockade (e.g. bradycardia, hypotension, respiratory distress and hypoglycaemia) have been observed in the neonate when beta-blockers have been administered until delivery. If this medical product is administered until delivery, the neonate should be carefully monitored during the first days of life.

Use During Lactation

It is not known whether dorzolamide is excreted in human milk. In lactating rats receiving dorzolamide, decreases in the body weight gain of offspring were observed.

Beta-blockers are excreted in breast milk. However, at therapeutic doses of timolol in eye drops it is not likely that sufficient amounts would be present in breast milk to produce clinical symptoms of beta-blockade in the infant. To reduce the systemic absorption, see 4.2. If treatment with COSOPT Preservative-Free is required, then lactation is not recommended.

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. Possible side effects such as blurred vision may affect some patients' ability to drive and/or operate machinery.

4.8 Undesirable effects

In a clinical study for COSOPT Preservative-Free the observed adverse reactions have been consistent with those that were reported previously with COSOPT (preserved formulation), dorzolamide hydrochloride and/or timolol maleate.

During clinical studies, 1035 patients were treated with COSOPT (preserved formulation). Approximately 2.4% of all patients discontinued therapy with COSOPT (preserved formulation) because of local ocular adverse reactions; approximately 1.2% of all patients discontinued because of local adverse reactions suggestive of allergy or hypersensitivity (such as lid inflammation and conjunctivitis).

COSOPT Preservative-Free has been shown to have a similar safety profile to COSOPT (preservative containing formulation) in a repeat dose double-masked, comparative study.

The following adverse reactions have been reported with COSOPT Preservative-Free or one of its components either during clinical trials or during post-marketing experience:

Like other topically applied ophthalmic medicines, timolol is absorbed into the systemic circulation. This may cause similar undesirable effects as seen with systemic betablocking agents. Incidence of systemic ADRs after topical ophthalmic administration is lower than for systemic administration.

Listed adverse reactions include reactions seen within the class of ophthalmic beta-blockers.

[Very Common: ($\geq 1/10$), Common: ($\geq 1/100$, $<1/10$), Uncommon: ($\geq 1/1000$, $<1/100$), and Rare: ($\geq 1/10,000$, $<1/1000$), Not known (cannot be estimated from the available data)]

System Organ Class (MedDRA)	Formulation	Very Common	Common	Uncommon	Rare	Not Known**
Immune system disorders	<u>COSOPT Preservative-Free</u>				signs and symptoms of systemic allergic reactions, including angioedema, urticaria, pruritus, rash, anaphylaxis	
	<u>Timolol maleate eye drops, solution</u>				signs and symptoms of allergic reactions including angioedema, urticaria, localized and generalized rash, anaphylaxis	pruritus
Metabolism and nutrition disorders	<u>Timolol maleate eye drops, solution</u>					hypoglycaemia
Psychiatric disorders	<u>Timolol maleate eye drops, solution</u>			depression*	insomnia*, nightmares*, memory loss,	
Nervous system disorders	<u>Dorzolamide hydrochloride eye drops, solution</u>		headache*		dizziness*, paraesthesia*	
	<u>Timolol maleate eye drops, solution</u>		headache*	dizziness*, syncope*	paraesthesia*, increase in signs and symptoms of myasthenia gravis, decreased	

					libido*, cerebrovascular accident*, cerebral ischaemia	
Eye disorders	<u>COSOPT Preservative- Free</u>	burning and stinging	conjunctival injection, blurred vision, corneal erosion, ocular itching, tearing			
	<u>Dorzolamide hydrochloride eye drops, solution</u>		eyelid inflammation*, eyelid irritation*	iritidocyclitis*	irritation including redness*, pain*, eyelid crusting*, transient myopia (which resolved upon discontinuation of therapy), corneal oedema*, ocular hypotony*, choroidal detachment (following filtration surgery)*	
	<u>Timolol maleate eye drops, solution</u>		signs and symptoms of ocular irritation including blepharitis*, keratitis*, decreased corneal sensitivity, and dry eyes*	visual disturbances including refractive changes (due to withdrawal of miotic therapy in some cases) *	ptosis, diplopia, choroidal detachment following filtration surgery* (see Special warning and precautions for use 4.4)	itching, tearing, redness, blurred vision, corneal erosion
Ear and labyrinth disorders	<u>Timolol maleate eye drops, solution</u>				tinnitus*	
Cardiac disorders	<u>Timolol maleate eye drops, solution</u>			bradycardia*	chest pain*, palpitation*, oedema*, arrhythmia*, congestive heart failure*, cardiac arrest*,	atrioventricular block, cardiac failure

					heart block	
Vascular disorders	<u>Timolol maleate eye drops, solution</u>				hypotension*, claudication, Raynaud's phenomenon*, cold hands and feet*	
Respiratory, thoracic, and mediastinal disorders	<u>COSOPT Preservative-Free</u>		sinusitis		shortness of breath, respiratory failure, rhinitis, rarely bronchospasm	
	<u>Dorzolamide hydrochloride eye drops, solution</u>				epistaxis*	
	<u>Timolol maleate eye drops, solution</u>			dyspnoea*	bronchospasm (predominantly in patients with pre-existing bronchospastic disease)*, respiratory failure, cough*	
Gastrointestinal disorders	<u>COSOPT Preservative-Free</u>	dysgeusia				
	<u>Dorzolamide hydrochloride eye drops, solution</u>		nausea*		throat irritation, dry mouth*	
	<u>Timolol maleate eye drops, solution</u>			nausea*, dyspepsia*	diarrhoea, dry mouth*	dysgeusia, abdominal pain, vomiting
Skin and subcutaneous tissue disorders	<u>COSOPT Preservative-Free</u>				contact dermatitis, Stevens-Johnson syndrome, toxic epidermal necrolysis	
	<u>Dorzolamide hydrochloride eye drops, solution</u>				rash*	
	<u>Timolol maleate eye drops, solution</u>				alopecia*, psoriasiform rash or exacerbation of psoriasis*	skin rash
Musculoskeletal	<u>Timolol</u>				systemic lupus	Myalgia

and connective tissue disorders	<u>maleate eye drops, solution</u>				erythematosis	
Renal and urinary disorders	<u>COSOPT Preservative-Free</u>			Urolithiasis		
Reproductive system and breast disorders	<u>Timolol maleate eye drops, solution</u>				Peyronie's disease*, decreased libido	sexual dysfunction
General disorders and administration site conditions	<u>Dorzolamide hydrochloride eye drops, solution</u>		asthenia/ fatigue*			
	<u>Timolol maleate eye drops, solution</u>			asthenia/ fatigue		

*These adverse reactions were also observed with COSOPT (preserved formulation) during post-marketing experience.

**Additional adverse reactions have been seen with ophthalmic beta-blockers and may potentially occur with COSOPT Preservative-Free.

4.9 Overdose

No data are available in humans in regard to overdose by accidental or deliberate ingestion of COSOPT (preserved formulation) or COSOPT Preservative-Free.

Symptoms

There have been reports of inadvertent overdoses with timolol maleate ophthalmic solution resulting in systemic effects similar to those seen with systemic beta-adrenergic blocking agents such as dizziness, headache, shortness of breath, bradycardia, bronchospasm, and cardiac arrest. The most common signs and symptoms to be expected with overdoses of dorzolamide are electrolyte imbalance, development of an acidotic state, and possibly central nervous system effects.

Only limited information is available with regard to human overdose by accidental or deliberate ingestion of dorzolamide hydrochloride. With oral ingestion, somnolence has been reported. With topical application the following have been reported: nausea, dizziness, headache, fatigue, abnormal dreams, and dysphagia.

Treatment

Treatment should be symptomatic and supportive. Serum electrolyte levels (particularly potassium) and blood pH levels should be monitored. Studies have shown that timolol does not dialyze readily.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antiglaucoma preparations and miotics, Beta blocking agents, Timolol, combinations, ATC code: S01ED51.

Mechanism of Action

COSOPT Preservative-Free is comprised of two components: dorzolamide hydrochloride and timolol maleate. Each of these two components decreases elevated intraocular pressure by reducing aqueous humor secretion, but does so by a different mechanism of action.

Dorzolamide hydrochloride is a potent inhibitor of human carbonic anhydrase II. Inhibition of carbonic anhydrase in the ciliary processes of the eye decreases aqueous humor secretion, presumably by slowing the formation of bicarbonate ions with subsequent reduction in sodium and fluid transport. Timolol maleate is a nonselective beta-adrenergic receptor blocking agent. The precise mechanism of action of timolol maleate in lowering intraocular pressure is not clearly established at this time, although a fluorescein study and tonography studies indicate that the predominant action may be related to reduced aqueous formation. However, in some studies a slight increase in outflow facility was also observed.

The combined effect of these two agents results in additional intraocular pressure reduction (IOP) compared to either component administered alone.

Following topical administration, COSOPT Preservative-Free reduces elevated intraocular pressure, whether or not associated with glaucoma. Elevated intraocular pressure is a major risk factor in the pathogenesis of optic nerve damage and glaucomatous visual field loss. This medicinal product reduces intraocular pressure without the common side effects of miotics such as night blindness, accommodative spasm and pupillary constriction.

Pharmacodynamic effects

Clinical Effects

Clinical studies of up to 15 months duration were conducted to compare the IOP-lowering effect of COSOPT (preserved formulation) b.i.d. (dosed morning and bedtime) to individually- and concomitantly-administered 0.5% timolol and 2.0% dorzolamide in patients with glaucoma or ocular hypertension for whom concomitant therapy was considered appropriate in the trials. This included both untreated patients and patients inadequately controlled with timolol monotherapy. The majority of patients were treated with topical beta-blocker monotherapy prior to study enrollment. In an analysis of the combined studies, the IOP-lowering effect of COSOPT (preserved formulation) b.i.d. was greater than that of monotherapy with either 2% dorzolamide t.i.d. or 0.5% timolol b.i.d. The IOP-lowering effect of COSOPT (preserved formulation) b.i.d. was equivalent to that of concomitant therapy with dorzolamide b.i.d. and timolol b.i.d. The IOP-lowering effect of COSOPT (preserved formulation) b.i.d. was demonstrated when measured at various time points throughout the day and this effect was maintained during long-term administration.

In an active-treatment-controlled, parallel, double-masked study in 261 patients with elevated intraocular pressure ≥ 22 mmHg in one or both eyes, COSOPT Preservative-Free had an IOP-lowering effect equivalent to that of COSOPT (preserved formulation). The safety profile of COSOPT Preservative-Free was similar to COSOPT (preserved formulation).

Paediatric population

A 3 month controlled study, with the primary objective of documenting the safety of 2% dorzolamide hydrochloride ophthalmic solution in children under the age of 6 years has been conducted. In this study, 30 patients under 6 and greater than or equal to 2 years of age whose IOP was not adequately controlled with monotherapy by dorzolamide or timolol received COSOPT (preserved formulation) in an open label phase.

Efficacy in those patients has not been established. In this small group of patients, twice daily administration of COSOPT (preserved formulation) was generally well tolerated with 19 patients completing the treatment period and 11 patients discontinuing for surgery, a change in medication, or other reasons.

5.2 Pharmacokinetic properties

Dorzolamide Hydrochloride

Unlike oral carbonic anhydrase inhibitors, topical administration of dorzolamide hydrochloride allows for the active substance to exert its effects directly in the eye at substantially lower doses and therefore with less systemic exposure. In clinical trials, this resulted in a reduction in IOP without the acid-base disturbances or alterations in electrolytes characteristic of oral carbonic anhydrase inhibitors.

When topically applied, dorzolamide reaches the systemic circulation.

To assess the potential for systemic carbonic anhydrase inhibition following topical administration, active substance and metabolite concentrations in red blood cells (RBCs) and plasma and carbonic anhydrase inhibition in RBCs were measured. Dorzolamide accumulates in RBCs during chronic dosing as a result of selective binding to CA-II while extremely low concentrations of free active substance in plasma are maintained. The parent active substance forms a single N-desethyl metabolite that inhibits CA-II less potently than the parent active substance but also inhibits a less active isoenzyme (CA-I). The metabolite also accumulates in RBCs where it binds primarily to CA-I. Dorzolamide binds moderately to plasma proteins (approximately 33%). Dorzolamide is primarily excreted unchanged in the urine; the metabolite is also excreted in urine. After dosing ends, dorzolamide washes out of RBCs nonlinearly, resulting in a rapid decline of active substance concentration initially, followed by a slower elimination phase with a half-life of about four months.

When dorzolamide was given orally to simulate the maximum systemic exposure after long term topical ocular administration, steady state was reached within 13 weeks. At steady state, there was virtually no free active substance or metabolite in plasma; CA inhibition in RBCs was less than that anticipated to be necessary for a pharmacological effect on renal function or respiration. Similar pharmacokinetic results were observed after chronic, topical administration of dorzolamide hydrochloride. However, some elderly patients with renal impairment (estimated CrCl 30-60 ml/min) had higher metabolite concentrations in RBCs, but no meaningful differences in carbonic anhydrase inhibition and no clinically significant systemic side effects were directly attributable to this finding.

Timolol Maleate

In a study of plasma active substance concentration in six subjects, the systemic exposure to timolol was determined following twice daily topical administration of timolol maleate ophthalmic solution 0.5%. The mean peak plasma concentration following morning dosing was 0.46 ng/ml and following afternoon dosing was 0.35 ng/ml.

5.3 Preclinical safety data

The ocular and systemic safety profile of the individual components is well established.

Dorzolamide

In rabbits given maternotoxic doses of dorzolamide associated with metabolic acidosis, malformations of the vertebral bodies were observed.

Timolol

Animal studies have not shown teratogenic effect.

Furthermore, no adverse ocular effects were seen in animals treated topically with dorzolamide hydrochloride and timolol maleate ophthalmic solution or with concomitantly-administered dorzolamide hydrochloride and timolol maleate. *In vitro* and *in vivo* studies with each of the components did not reveal a mutagenic potential. Therefore, no significant risk for human safety is expected with therapeutic doses of COSOPT Preservative-Free.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Hydroxyethylcellulose
Mannitol (E421)
Sodium citrate (E331)
Sodium hydroxide (E524) for pH adjustment
Water for injection

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

The shelf life expiry date for this product shall be the date shown on the sachet and outer package of the product on the market in the country of origin.

After first opening of the sachet: 15 days. Discard any unused single dose containers after that time.

Discard the opened single dose container immediately after first use.

6.4 Special precautions for storage

Do not store above 30°C. Do not freeze.
Store in original packaging in order to protect from light.

6.5 Nature and contents of container

COSOPT Preservative-Free is available in 0.2 ml low density polyethylene single dose containers in an aluminum sachet containing 15 single -dose containers.

Pack size: 60 x 0.2 ml (4 sachets with 15 single dose containers)

Not all pack sizes may be marketed.

6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product

No special requirements.

7 PARALLEL PRODUCT AUTHORISATION HOLDER

LTT Pharma Limited
Unit 18
Oxleasow Road
East Moon Moat
Redditch 2
Worcestershire
B98 0RE
UK

8 PARALLEL PRODUCT AUTHORISATION NUMBER

PPA1562/43/1

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 11th February 2011

10 DATE OF REVISION OF THE TEXT

August 2013