

# COVERSYL® ARGININE 10mg film-coated tablets

(perindopril arginine)

## Patient Information Leaflet

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- \* Keep this leaflet. You may need to read it again.
- \* If you have any further questions, ask your doctor or pharmacist.
- \* This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- \* If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

### In this leaflet:

- 1 What Coversyl Arginine is and what it is used for
- 2 What you need to know before you take Coversyl arginine
- 3 How to take Coversyl Arginine
- 4 Possible side effects
- 5 How to store Coversyl Arginine
- 6 Content of the pack and other information

## 1 What Coversyl Arginine is and what it is used for

Coversyl Arginine is an angiotensin converting enzyme (ACE) inhibitor. These work by widening the blood vessels, which makes it easier for your heart to pump blood through them.

Coversyl Arginine is used:

- \* to treat **high blood pressure** (hypertension),
- \* to reduce the risk of cardiac events, such as heart attack, in patients with **stable coronary artery disease** (a condition where the blood supply to the heart is reduced or blocked) and who have already had a heart attack and/or an operation to improve the blood supply to the heart by widening the vessels that supply it.

## 2 What you need to know before you take Coversyl arginine

**Do not take Coversyl Arginine**

- \* if you are allergic to perindopril, or any of the other ingredients of this medicine (listed in Section 6), or to any other ACE inhibitor
- \* if you have experienced symptoms such as wheezing, swelling of the face, tongue or throat, intense itching or severe skin rashes with previous ACE inhibitor treatment or if you or a member of your family have had these symptoms in any other circumstances (a condition called angioedema).
- \* if you are more than 3 months pregnant (It is also better to avoid Coversyl arginine in early pregnancy – see pregnancy section).
- \* if you have diabetes mellitus or kidney problems and you are treated with aliskiren.

**Warning and precaution**

If any of the following apply to you please talk to your doctor, or pharmacist or nurse before taking Coversyl Arginine if you:

- \* have aortic stenosis (narrowing of the main blood vessel leading from the heart) or hypertrophic cardiomyopathy (heart muscle disease) or renal artery stenosis (narrowing of the artery supplying the kidney with blood),
- \* have any other heart problems,
- \* have liver problems,
- \* have kidney problems or if you are receiving dialysis,
- \* suffer from a collagen vascular disease (disease of the connective tissue) such as systemic lupus erythematosus or scleroderma,
- \* have diabetes,
- \* are on a salt restricted diet or use salt substitutes which contain potassium,
- \* are to undergo anaesthesia and/or major surgery,
- \* are to undergo LDL apheresis (which is removal of cholesterol from your blood by a machine),
- \* are going to have desensitisation treatment to reduce the effects of an allergy to bee or wasp stings,
- \* have recently suffered from diarrhoea or vomiting or are dehydrated,
- \* have been told by your doctor that you have an intolerance to some sugars.
- \* take aliskiren (medicines used to treat hypertension).
- \* are of black origin since you may have a higher risk of angioedema and this medicine may be less effective in lowering your blood pressure than in non-black patients.

**Angioedema**

Angioedema (a severe allergic reaction with swelling of the face, lips, tongue or throat with difficulty in swallowing or breathing) has been reported in patients treated with ACE inhibitors, including Coversyl arginine. This may occur at any time during treatment. If you develop such symptoms, you should stop taking Coversyl arginine and see a doctor immediately. See also section 4.

You must tell your doctor if you think you are (or might become) pregnant. Coversyl Arginine is not recommended in early pregnancy, and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see pregnancy section).

**Children and adolescent**

Coversyl arginine is not recommended for use in children and adolescents.

**Other medicines and Coversyl arginine**

Please tell your doctor or pharmacist if you are taking or might take any other medicines.

Treatment with Coversyl Arginine can be affected by other medicines. These include:

- \* other medicines for high blood pressure, including aliskiren, diuretics (medicines which increase the amount of urine produced by the kidneys),
- \* potassium-sparing drugs (e.g. triamterene, amiloride), potassium supplements or potassium-containing salt substitutes,
- \* potassium-sparing drugs used in the treatment of heart failure : eplerenone and spironolactone at doses between 12,5 mg to 50 mg by day,
- \* lithium for mania or depression,
- \* non-steroidal anti-inflammatory drugs (e.g. ibuprofen) for pain relief or high dose aspirin,
- \* medicines to treat diabetes (such as insulin or metformin),
- \* baclofen (used to treat muscle stiffness in diseases such as multiple sclerosis),
- \* medicines to treat mental disorders such as depression, anxiety, schizophrenia etc (e.g. tricyclic antidepressants, antipsychotics),
- \* immunosuppressants (medicines which reduce the defence mechanism of the body) used for the treatment of auto-immune disorders or following transplant surgery (e.g. ciclosporin, tacrolimus),
- \* trimethoprim (for the treatment of infections),
- \* estramustine (used in cancer therapy),
- \* allopurinol (for the treatment of gout),
- \* procainamide (for the treatment of an irregular heart beat),
- \* vasodilators including nitrates (products that make the blood vessels become wider),
- \* heparin (medicines used to thin blood),
- \* medicines used for the treatment of low blood pressure, shock or asthma (e.g. ephedrine, noradrenaline or adrenaline).
- \* gold salts, especially with intravenous administration (used to treat symptoms of rheumatoid arthritis).

**Coversyl arginine with food and drink**

It is preferable to take Coversyl Arginine before a meal.

**Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

**Pregnancy**

You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking Coversyl Arginine before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Coversyl Arginine. Coversyl Arginine is not recommended in early pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

**Breast-feeding**

**Tell** your doctor if you are breast-feeding or about to start breast-feeding. Coversyl Arginine is not recommended for mothers who are breast-feeding, and your doctor may choose another treatment for you if you wish to breast-feed, especially if your baby is newborn, or was born prematurely.

**Driving and using machines**

Coversyl Arginine usually does not affect alertness but dizziness or weakness due to low blood pressure may occur in certain patients. If you are affected in this way, your ability to drive or to operate machinery may be impaired.

**Coversyl arginine contains lactose**

Coversyl Arginine contains lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

## 3 How to take Coversyl Arginine

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure. Swallow your tablet with a glass of water, preferably at the same time each day, in the morning, before a meal. Your doctor will decide on the correct dose for you.

The recommended dosages are as follows:

High blood pressure: the usual starting and maintenance dose is 5 mg once daily. After one month, this can be increased to 10 mg once a day if required. 10 mg a day is the maximum recommended dose for high blood pressure.

If you are 65 or older, the usual starting dose is 2.5 mg once a day. After a month this can be increased to 5 mg once a day and then if necessary to 10 mg once daily.

**Stable coronary artery disease:** the usual starting dose is 5 mg once daily. After two weeks, this can be increased to 10 mg once daily, which is the maximum recommended dose in this indication.

If you are 65 or older, the usual starting dose is 2.5 mg once a day. After a week this can be increased to 5 mg once a day and after a further week to 10 mg once daily.

**Use in children and adolescent**

Use in children and adolescent is not recommended.

**If you take more Coversyl arginine than you should**

If you take too many tablets, contact your nearest accident and emergency department or tell your doctor immediately. The most likely effect in case of overdose is low blood pressure which can make you feel dizzy or faint. If this happens, lying down with the legs raised can help.

**If you forget to take Coversyl Arginine**

It is important to take your medicine every day as regular treatment works better. However, if you forget to take a dose of Coversyl Arginine, take the next dose at the usual time. Do not take a double dose to make up for a forgotten dose.

# COVERSYL® ARGININE 10mg film-coated tablets

(perindopril arginine)

## Patient Information Leaflet (continued)

### If you stop taking Coversyl Arginine

As the treatment with Coversyl Arginine is usually life-long, you should discuss with your doctor before stopping this medicinal product.

If you have any further questions on the use of this product, ask your doctor, pharmacist or nurse.

## 4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

- Stop taking the medicinal product and see a doctor immediately, if you experience any of the following side effects that can be serious:**
- \* swelling of the face, lips, mouth, tongue or throat, difficulty in breathing (angioedema) (See section 2 "Warning and precaution") (Uncommon -may affect up to 1 in 100 people),
  - \* severe dizziness or fainting due to low blood pressure (Common - may affect up to 1 in 10 people),
  - \* unusual fast or irregular heart beat,chest pain(angina) or heart attack (Very rare - may affect up to 1 in 10,000 people),
  - \* weakness of arms or legs, or problems speaking which could be a sign of a possible stroke (Very rare – may affect up to 1 in 10,000 people),
  - \* sudden wheeziness, chest pain, shortness of breath, or difficulty in breathing (bronchospasm) (Uncommon - may affect up to 1 in 100 people),
  - \* inflamed pancreas which may cause severe abdominal and back pain accompanied with feeling very unwell (Very rare - may affect up to 1 in 10,000 people),
  - \* yellowing of the skin or eyes (jaundice) which could be a sign of hepatitis (Very rare - may affect up to 1 in 10,000 people),
  - \* skin rash which often starts with red itchy patches on your face, arms or legs (erythema multiforme) (Very rare - may affect up to 1 in 10,000 people).

### Tell your doctor if you notice any of the following side effects:

- Common (may affect up to 1 in 10 people):
- \* headache,
  - \* dizziness,
  - \* vertigo,
  - \* pins and needles,
  - \* vision disturbances,
  - \* tinnitus (sensation of noises in the ears),
  - \* cough,
  - \* shortness of breath(dyspnoea),
  - \* gastro-intestinal disorders (nausea, vomiting, abdominal pain, taste disturbances, dyspepsia or difficulty of digestion, diarrhoea, constipation),
  - \* allergic reactions (such as skin rashes, itching),
  - \* muscle cramps, feeling of tiredness,

- Uncommon (may affect to 1 in 100 people ):
- \* mood swings,
  - \* sleep disturbances,
  - \* dry mouth,
  - \* intense itching or severe skin rashes,
  - \* formation of blister clusters over the skin,
  - \* kidney problems,
  - \* impotence,
  - \* sweating,
  - \* excess of eosinophils (a type of white blood cells),
  - \* somnolence,
  - \* fainting,
  - \* palpitations,
  - \* tachycardia,
  - \* vasculitis (inflammation of blood vessels),
  - \* photosensitivity reaction (increased sensitivity of the skin to sun),
  - \* arthralgia (joint pain),
  - \* myalgia (muscle pain),
  - \* chest pain,
  - \* malaise,
  - \* oedema peripheral,
  - \* fever,
  - \* fall,
  - \* change in laboratory parameters: high blood level of potassium reversible on discontinuation, low level of sodium, hypoglycaemia (very low blood sugar level) in case of diabetic patients, increased blood urea, and increased blood creatinine.

- Rare (may affect up to 1 in 1000 people):
- \* changes in laboratory parameters: Increased level of liver enzymes, high level of serum bilirubin.

- Very rare (may affect up to 1 in 10,000 people ):
- \* confusion,
  - \* eosinophilic pneumonia (a rare type of pneumonia),
  - \* rhinitis (blocked up or runny nose),
  - \* acute renal failure,
  - \* changes in blood values such as a lower number of white and red blood cells, lower haemoglobin, lower number of blood platelets

### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL – Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; e-mail: medsafety@hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

## 5 How to store Coversyl Arginine

Keep out of the reach and sight of children.  
Do not use Coversyl Arginine after the expiry date which is stated on the container and outer packaging after (EXP).  
The expiry date refers to the last day of that month.


Keep the bottle tightly closed in order to protect from moisture.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## 6 Content of the pack and other information

### What Coversyl Arginine 10 mg contains

- \* The active substance is perindopril arginine. One film-coated tablet contains 6.790 mg perindopril (corresponding to 10 mg perindopril arginine)
- \* The other ingredients in the tablet core are: lactose monohydrate, magnesium stearate, maltodextrin, hydrophobic colloidal silica, sodium starch glycolate (type A), and in the tablet film-coating: glycerol, hypromellose, copper chlorophyllin, macrogol 6000, magnesium stearate, titanium dioxide.

**What Coversyl Arginine 10 mg looks like and contents of the pack**  
Coversyl Arginine 10 mg tablets are green, round, biconvex, film-coated tablets engraved with  on one face and a company logo on the other face.

The tablets are available in containers of 30 tablets.

**The Parallel Product Authorisation Holder is:**  
LTT Pharma Limited, Unit 18, Oxleasow Road, East Moons Moat, Redditch, Worcestershire, B98 0RE, UK.

**Repackaged by:**  
Lexon (UK) Limited, Unit 18, Oxleasow Road, East Moons Moat, Redditch, Worcestershire, B98 0RE, UK.

**The manufacturer is:**  
Manufacturers: Les Laboratoires Servier Industrie, 905 route de Saran, 45520 Gidy, France.  
and  
Servier (Ireland) Industries Ltd., Gorey Road, Arklow, Co. Wicklow, Ireland and  
EGIS Pharmaceuticals PLC. H-1106 Budapest, 30-38 Kereszturi Way, Hungary.  
or  
EGIS Pharmaceuticals Plc, H-9900 Körmend, King Matthias' 65th Street, Hungary.

COVERSYL is a registered trademark of BIOFARMA.

Revision date: 22/12/2014

**POM** PPA1562/079/002

Blind or partially sighted?  
Is this leaflet hard to see or read?  
Phone LTT Pharma Limited,  
Tel: +44 (0)1527 505414 for help.

<b>This medicinal product is authorised in the Member States of the EEA under the following names:</b>	
Austria	Coversum-Arginin 10 mg-Filmtabletten
Belgium	COVERSYL 10 mg
Cyprus	COVERSYL 10 mg
Czech Republic	Prestarium Neo Forte
Denmark	COVERSYL NOVUM 10 mg
Estonia	Prestarium Arginine 10 mg
Finland	COVERSYL NOVUM 10 mg
France	COVERSYL 10 mg
Germany	COVERSUM Arginin 10 mg
Greece	COVERSYL 10 mg
Hungary	ARMIX Arginin 10 mg
Iceland	COVERSYL NOVUM 10 mg
Ireland	COVERSYL Arginine 10 mg film-coated tablets
Italy	COVERSYL 10 mg
Latvia	PRESTARIUM 10 mg
Lithuania	PRESTARIUM 10 mg
Luxembourg	COVERSYL 10 mg
Malta	COVERSYL 10 mg
Netherlands	COVERSYL arg 10 mg
Norway	PERINDOPRILARGININ SERVIER 10 mg
Poland	PRESTARIUM 10 mg
Portugal	COVERSYL 10 mg
Slovakia	PRESTARIUM A 10 mg
Slovenia	BIOPREXANIL 10 mg
Sweden	COVERSYL NOVUM 10 mg
United Kingdom	COVERSYL Arginine 10 mg