

5. HOW TO STORE REQUIP-MODUTAB

- Keep this medicine out of the sight and reach of children.
- Do not use Requip Mobutab after the expiry date which is stated on the blister and on the carton. The expiry date refers to the last day of that month.
- Do not store above 25°C. Store in the original package in order to protect from light.

Do not throw away any medicines via waste water or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Requip-Modutab contains

- The active substance is 4mg of ropinirole (as hydrochloride)
 - The other ingredients are **tablet core** - hypromellose, hydrogenated castor oil, carmellose sodium, povidone (K 29-32), maltodextrin, magnesium stearate, lactose monohydrate, anhydrous colloidal silica mannitol (E421), iron oxide yellow (E172) and glycerol dibehenate
- Film-coat** - opadry light brown OY-27207, (hypromellose, titanium dioxide (E171), macrogol 400, sunset yellow FCF aluminium lake (E110), indigo carmine aluminium lake (E132)).

What Requip-Modutab looks like and contents of the pack

- Requip-Modutab is a light brown prolonged-release tablets marked “GS” on one side and ‘WXG’ on reverse side.
- Requip-Modutab comes in blister packs of 84 prolonged- release tablets.

Licence Holder:

Product Licence Holder: LTT Pharma Limited, Unit 18, Oxleasow Road, East Moons Moat, Redditch, Worcestershire, B98 0RE.

Repackaged by:

Lexon (UK) Limited, Unit 18, Oxleasow Road, East Moons Moat, Redditch, Worcestershire, B98 0RE.

Manufacturer

This medicine is manufactured by SmithKline Beecham Plc T/A SmithKline Beecham Pharmaceuticals, Manor Royal, Crawley, West Sussex, RH10 9QJ United Kingdom or Glaxo Wellcome GmbH & Co.KG, Industriestrasse 32-36, 23843, Bad Oldesole, Germany or GlaxoSmithKline Pharmaceuticals S.A ., Ul. Grunwaldzka, 61-322, Poznan, Poland or Glaxo Wellcome S.A., Avenida de Extremadura 3, Aranda de Duero, Burgos, Spain.

Requip is a registered trademark of the GlaxoSmithKline Group of Companies.

Leaflet revision date: 14/02/2018



PPA1562/115/002

Blind or partially sighted?

Is this leaflet hard to see or read?

Phone LTT Pharma Limited,

Tel: +44 (0)1527 505414 for help.

Package leaflet: Information for the user
Requip-Modutab® 4mg Prolonged-release Tablets
ropinirole (as hydrochloride)

Ref: 115/002/140218/1/F

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- Keep this leaflet. You may need to read it again.
- If you have any more questions, ask your doctor or pharmacist.
- **If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.**

What is in this leaflet

1. What Requip-Modutab is and what it is used for
2. What you need to know before you take Requip-Modutab
3. How to take Requip-Modutab
4. Possible side effects
5. How to store Requip-Modutab
6. Contents of the pack and other information

1. WHAT REQUIP-MODUTAB IS AND WHAT IT IS USED FOR

The active ingredient in Requip-Modutab is ropinirole, which belongs to a group of medicines called dopamine agonists. Dopamine agonists affect the brain in a similar way to a natural substance called dopamine.

Requip-Modutab prolonged-release tablets are used to treat Parkinson's disease.

People with Parkinson's disease have low levels of dopamine in some parts of their brains. Ropinirole has effects similar to those of natural dopamine, so it helps to reduce the symptoms of Parkinson's disease.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE REQUIP-MODUTAB

Do not take Requip-Modutab:

- if you are **allergic** to ropinirole or any of the other ingredients of this medicine (listed in section 6).
 - if you have **serious kidney disease**
 - if you have **liver disease**
- Tell your doctor if you think any of these may apply to you.

Warnings and precautions

Talk to your doctor or pharmacist before taking Requip-Modutab:

- if you are **pregnant** or think you may be pregnant
- if you are **breast feeding**
- if you are **under 18 years old**
- if you have a **serious heart complaint**
- if you have a **serious mental health problem and/or behaviours** (see section 4).
- if you have an **intolerance to some sugars** (such as lactose).

Tell your doctor if you experience symptoms such as depression, apathy, anxiety, lack of energy, sweating or pain after stopping or reducing your Requip-Modutab treatment. If the problems persist more than a few weeks, **your doctor may need to adjust your dose.**

Tell your doctor if you or your family/carer notices that you are developing urges or cravings to behave in ways that are unusual for you and you cannot resist the impulse, drive or temptation to carry out certain activities that could harm yourself or others. These are called impulse control disorders and can include behaviours such as addictive gambling, excessive eating or spending, an abnormally high sex drive or an increase in sexual thoughts or feelings. **Your doctor may need to adjust or stop your dose.**

→ **Tell your doctor** if you think any of these may apply to you. Your doctor may decide that Requip-Modutab is not suitable for you, or that you need extra check-ups while you are taking it.

While you are taking Requip-Modutab

Tell your doctor if you or your family notices that you are developing any **unusual behaviours** (such as an **unusual urge to gamble or increased sexual urges and/or behaviours**) while you are taking Requip-Modutab. Your doctor may need to adjust or stop your dose.

Smoking and Requip-Modutab

Tell your doctor if you start smoking, or give up smoking, while you are taking Requip-Modutab. Your doctor may need to adjust your dose.

Other medicines and Requip-Modutab

Tell your doctor or pharmacist if you are taking, or have recently taken or might take any other medicines, including any herbal medicines or other medicines you obtained without a prescription.

Remember to tell your doctor or pharmacist if you begin taking a new medicine while you are taking Requip-Modutab.

Some medicines can affect the way Requip-Modutab works, or make it more likely that you will have side effects. Requip-Modutab can also affect the way some other medicines work.

These include:

- the **anti-depressant fluvoxamine**
 - medication for other **mental health problems**, for example **sulpiride**
 - **HRT** (hormone replacement therapy)
 - **metoclopramide**, which is used to treat **nausea and heartburn**
 - the **antibiotics ciprofloxacin** or **enoxacin**
 - any other medicine for **Parkinson's disease.**
- **Tell your doctor** if you are taking, or have recently taken, any of these.

You will require additional blood tests if you are taking these medicines with Requip-Modutab:

- Vitamin K antagonists (used to reduce blood clotting) such as Warfarin (coumadin).

Requip-Modutab with food and drink
You can take Requip-Modutab with or without food, as you prefer.

Pregnancy and breast feeding
Requip-Modutab is not recommended if you are pregnant, unless your doctor advises that the benefit to you of taking Requip-Modutab is greater than the risk to your unborn baby.
Requip-Modutab is not recommended if you are breast feeding, as it can affect your milk production.

Tell your doctor immediately if you are pregnant, if you think you might be pregnant, or if you are planning to become pregnant. Your doctor will also advise you if you are breast feeding or planning to do so. Your doctor may advise you to stop taking Requip-Modutab.

Driving and using machines
Requip-Modutab can make you feel drowsy. **It can make people feel extremely sleepy**, and it sometimes makes people **fall asleep very suddenly without warning**.
Requip-Modutab can cause hallucinations (seeing, hearing or feeling things that are not there). If affected, do not drive or use machines. If you could be affected: **do not drive, do not operate machines and do not** put yourself in any situation where feeling sleepy or falling asleep could put you (or other people) at risk of serious injury or death. Do not take part in these activities until you are no longer affected.

→ **Talk to your doctor** if this causes problems for you. Requip-Modutab prolonged-release tablets contain a sugar called **lactose**. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking Requip-Modutab.

The 4 mg prolonged-release tablet contains a colouring called **sunset yellow FCF (E110)** which may cause allergic reactions.

3. HOW TO TAKE REQUIP MODUTAB

Always take Requip Modutab exactly as your doctor or pharmacist has told you to. Check with your doctor or pharmacist if you are not sure.

Do not give Requip Modutab to children. Requip Modutab is not normally prescribed for people under 18.

You may be given Requip Modutab on its own to treat the symptoms of your Parkinson's disease. Or you may be given Requip Modutab as well as another medicine called L-dopa (*also called levodopa*). If you are taking L-dopa you may experience some uncontrollable movements (dyskinesias) when you first start taking Requip Modutab. Tell your doctor if this happens, as your doctor may need to adjust the doses of the medicines you are taking.

Requip-Modutab tablet(s) are designed to release drug over a 24hr period. If you have a condition where your medicine passes through your body too quickly, e.g., diarrhoea, the tablet(s) may not dissolve completely and may not work properly. You may see tablet(s) in your stool. If this happens, let your doctor know as soon as possible.

How much Requip-Modutab will you need to take?
It may take a while to find out the best dose of Requip-Modutab for you.


The recommended starting dose of Requip-Modutab prolonged-release tablets is 2 mg once daily for the first week. Your doctor may increase your dose to 4 mg of Requip-Modutab prolonged-release tablets once daily, from the second week of treatment. If you are very elderly, your doctor may increase your dose more slowly.

After that, the doctor may adjust your dose until you are taking the dose that is best for you. Some people take up to 24 mg of Requip-Modutab prolonged-release tablets each day.

If at the start of your treatment, you experience side effects that you find difficult to tolerate, speak to your doctor. Your doctor may advise you to switch to a lower dose of ropinirole film-coated (immediate-release) tablets which you will take three times a day.

Do not take any more Requip-Modutab than your doctor has recommended.
It may take a few weeks for Requip-Modutab to work for you.

Taking your dose of Requip-Modutab
Take Requip-Modutab once a day, at the same time each day.
Swallow your Requip-Modutab prolonged-release tablet(s) whole, with a glass of water.



Do not break, chew or crush the prolonged-release tablet(s) — if you do, there is a danger you could overdose, because the medicine will be released into your body too quickly.

If you are switching from ropinirole film-coated (immediate-release) tablets
Your doctor will base your dose of Requip-Modutab prolonged-release tablets on the dose of ropinirole film-coated (immediate-release) tablets you were taking. Take your ropinirole film-coated immediate-release) tablets as normal the day before you switch. Then take your Requip-Modutab prolonged-release tablets next morning and do not take any more ropinirole film-coated (immediate-release) tablets.

If you take more Requip-Modutab than you should Contact a doctor or pharmacist immediately. If possible, show them the Requip-Modutab pack.
Someone who has taken an overdose of Requip-Modutab may have any of these symptoms: feeling sick (nausea), being sick (vomiting), dizziness (a spinning sensation), feeling drowsy, mental or physical tiredness, fainting, hallucinations.

If you forget to take Requip-Modutab
Do not take extra prolonged-release tablets or a double dose to make up for a missed dose.

If you have missed taking Requip Modutab for one day or more, ask your doctor for advice on how to start taking it again.

If you stop taking Requip-Modutab
Do not stop taking Requip-Modutab without advice.

Take Requip-Modutab for as long as your doctor recommends. Do not stop unless your doctor advises you to.

If you suddenly stop taking Requip-Modutab, your Parkinson's disease symptoms may quickly get much worse. A sudden stop could cause you to develop a medical condition called neuroleptic malignant syndrome which may represent a major health risk. The symptoms include: akinesia (loss of muscle movement), rigid muscles, fever, unstable blood pressure, tachycardia (increased heart rate), confusion, depressed level of consciousness (e.g. coma).

If you need to stop taking Requip-Modutab, your doctor will reduce your dose gradually.

If you have any further questions on the use of this medicines, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everyone gets them. The side effects of Requip-Modutab are more likely to happen when you first start taking it, or when your dose has just been increased. They are usually mild, and may become less troublesome after you have taken the dose for a while. If you are worried about side effects, talk to your doctor.

Very common side effects
These may **affect more than 1 in 10 people** taking Requip-Modutab:

- fainting
- feeling drowsy
- feeling sick (nausea).

Common side effects
These may **affect up to 1 in 10 people** taking Requip-Modutab

- falling asleep very suddenly without feeling sleepy first (*sudden sleep onset episodes*)
- hallucinations ('seeing' things that are not really there)
- being sick (vomiting)
- feeling dizzy (a spinning sensation)
- heartburn
- stomach pain
- constipation
- swelling of the legs, feet or hands.

Uncommon side effects
These may affect **up to 1 in 100 people** taking Requip-Modutab:

- feeling dizzy or faint, especially when you stand up suddenly (this is caused by a drop in blood pressure)
- feeling very sleepy during the day (*extreme somnolence*)
- mental problems such as delirium (severe confusion), delusions (unreasonable ideas) or paranoia (unreasonable suspicions).

Some patients may have the following side effects (frequency not known: cannot be estimated from the available data)

- **allergic reactions** such as red, itchy **swellings** on the skin (hives), swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing, **rash** or intense itching (see Section 2)
- changes in **liver function**, which have shown up in blood tests
- act in an aggressive manner
- excessive use of Requip-Modutab (craving for large doses of dopaminergic drugs in excess of that required to control motor symptoms, known as dopamine dysregulation syndrome)
- inability to resist that impulse, drive or temptation to perform an action that could be harmful to you or others, which may include:
 - Strong impulse to gamble excessively despite serious personal or family consequences.
 - Altered or increased sexual interest and behaviour of significant concern to you or to others, for example, an increased sexual drive
 - Uncontrollable excessive shopping or spending.
 - binge eating (eating large amounts of food in a short time period) or compulsive eating (eating more food than is needed to satisfy your hunger).

- Depression, apathy, anxiety, lack of energy, sweating or pain may occur (called dopamine agonist withdrawal syndrome or DAWS) after stopping or reducing your Requip-Modutab treatment.

Tell your doctor if you experience any of these behaviours; they will discuss ways of managing or reducing the symptoms.

If you are taking Requip-Modutab with L-dopa
People who are taking Requip-Modutab with L-dopa may develop other side effects over time:

- uncontrollable movements (dyskinesias) are a very common side effect. If you are taking L-dopa you may experience some uncontrollable movements (dyskinesias) when you first start taking Requip-Modutab. Tell your doctor if this happens, as your doctor may need to adjust the doses of the medicines you are taking.
- feeling confused is a common side effect.

Reporting of side effects
If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL – Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; e-mail: medsafety@hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine