

# Summary of Product Characteristics

## 1 NAME OF THE MEDICINAL PRODUCT

Elantan LA 25mg Prolonged-release Capsules

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each prolonged release capsule contains 25 mg isosorbide mononitrate.

Contains sucrose and lactose

For a full list excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Prolonged-release Capsules

*Product imported from the UK:*

Hard capsule with brown cap and white body, containing white to off-white pellets.

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic Indications

In the prophylaxis of angina pectoris.

### 4.2 Posology and method of administration

For oral administration.

#### Adults

One capsule to be taken in the morning.

For patients with higher nitrate requirement, the dose may be increased to two capsules taken simultaneously. The lowest effective dose should be used.

#### Elderly

There is no evidence to suggest that an adjustment of the dosage is necessary.

#### Children

The safety and efficacy of Elantan LA 25 has yet to be established in children.

Attenuation of effect has occurred in some patients being treated with prolonged release preparations. In such patients intermittent therapy may be more appropriate (see section 4.4).

Treatment with Elantan LA, as with any other nitrate, should not be stopped suddenly. Both dosage and frequency should be tapered gradually (see section 4.4).

### 4.3 Contraindications

Elantan LA25 should not be used in cases of acute myocardial infarction with low filling pressure, acute circulatory failure (shock, vascular collapse), or very low blood pressure, hypertrophic obstructive cardiomyopathy (HOCM), constrictive pericarditis, cardiac tamponade, low cardiac filling pressures, aortic/mitral valve stenosis and diseases associated with a raised intra-cranial pressure e.g. following a head trauma and including cerebral haemorrhage.

This product should not be given to patients with a known sensitivity to Isosorbide mononitrate, the listed ingredients or other nitrates.

Elantan LA25 should not be used in patients with marked anaemia, severe hypotension, closed angle glaucoma or hypovolaemia.

Phosphodiesterase type-5 inhibitors (e.g. sildenafil, tadalafil and vardenafil) have been shown to potentiate the hypotensive effects of nitrates, and their co-administration with nitrates or nitric oxide donors is therefore contraindicated (see section 4.5).

#### **4.4 Special warnings and precautions for use**

The capsules should be used with caution in patients who have a recent history of myocardial infarction or who are suffering from hypothyroidism, hypothermia, malnutrition, severe liver disease or severe renal disease.

Symptoms of circulatory collapse may arise after first dose, particularly in patients with labile circulation.

This product may give rise to postural hypotension and syncope in some patients. Severe postural hypotension with light-headedness and dizziness is frequently observed after the consumption of alcohol.

Hypotension induced by nitrates may be accompanied by paradoxical bradycardia and increased angina.

Elantan LA capsules contain lactose and therefore should not be used in patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption.

Elantan LA capsules also contain sucrose and therefore patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine. In the event of an acute angina attack, a sublingual treatment such as a GTN spray or tablet should be used instead of Elantan LA capsules.

If these capsules are not taken as indicated (see section 4.2.) tolerance to the medication could develop. In some patients being treated with prolonged release preparations, attenuation of effect is observed. In such patients, intermittent therapy may be more appropriate. The lowest effective dose should be used.

Treatment of Elantan LA, as with any other nitrate, should not be stopped suddenly. Both the dosage and frequency should be tapered gradually (see section 4.2).

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Concurrent administration of drugs with blood pressure lowering properties, e.g. beta-blockers, calcium channel blockers, vasodilators, alprostadil, aldesleukin, angiotensin II receptor antagonists etc and/or alcohol may potentiate the hypotensive effect of Elantan LA. This may also occur with neuroleptics and tricyclic antidepressants. Any blood pressure lowering effect of Elantan LA25 will be increased if used together with Phosphodiesterase type-5 inhibitors, which are used for erectile dysfunction (see special warnings and contraindications sections 4.4 & 4.3 respectively).

This might lead to life threatening cardiovascular complications. Patients who are on Elantan LA25 therapy therefore must not use Phosphodiesterase type-5 inhibitors.

Reports suggest that concomitant administration of Elantan LA may increase the blood level of dihydroergotamine and its hypertensive effect.

#### 4.6 Fertility, pregnancy and lactation

Safety in pregnancy has not been established for isosorbide mononitrate and it is not known whether nitrates are excreted in human milk. Therefore Elantan should only be used in pregnancy and during lactation if, in the opinion of the physician, the possible benefits of treatment outweigh the hazards.

#### 4.7 Effects on ability to drive and use machines

Dizziness, tiredness or blurred vision may occur at the start of treatment. If affected do not drive or operate machinery. This effect may be increased by alcohol.

#### 4.8 Undesirable effects

A very common (>10%) adverse reaction to Elantan LA25 is throbbing headache. The evidence of headache diminishes gradually with time and continued use.

At the start of therapy or when the dosage is increased, hypotension and/or light headedness in upright position are observed commonly (i.e. in 1-10% of patients). These symptoms may be associated with dizziness, drowsiness, reflex tachycardia and a feeling of weakness.

Infrequently (i.e. in <1% of patients) nausea, vomiting, flushing and allergic skin reaction (e.g. rash) may occur sometimes severely. In single cases, exfoliative dermatitis may occur.

Severe hypotensive response have been reported for organic nitrates and include nausea, vomiting, restlessness, pallor and excessive perspiration. Uncommonly collapse may occur (sometimes accompanied by bradyarrhythmia and syncope). Uncommonly severe hypotension may lead to enhanced angina symptoms.

A few reports of heartburn most likely due to nitrate-induced sphincter relaxation have been reported. Tachycardia and paroxysmal bradycardia have been reported.

#### 4.9 Overdose

Symptoms and signs:

Headache, hypotension, nausea, vomiting, sweating, tachycardia, vertigo, restlessness, warm flushed skin, blurred vision and syncope. A rise in intracranial pressure with confusion and neurological deficits can sometimes occur. Methaemoglobinaemia (cyanosis, hypoxaemia, restlessness, respiratory depression, convulsions, cardiac arrhythmias, circulatory failure, raised intracranial pressure) occurs rarely.

Management:

Consider oral activated charcoal if ingestion of a potentially toxic amount has occurred within 1 hour. Observe for at least 12 hours after the overdose. Monitor blood pressure and pulse. Correct hypotension by raising the foot of the bed and/or by expanding the intravascular volume. Other measures as indicated by the patient's clinical condition. If severe hypotension persists despite the above measures consider use of inotropes.

If methaemoglobinaemia (symptoms or >30 % methaemoglobin), IV administration of methylene blue 1-2 mg/kg body weight. If therapy fails with second dose after 1 hour or contraindicated, consider red blood cell concentrates or exchange transfusion.

In case of cerebral convulsions, diazepam or clonazepam IV, or if therapy fails, phenobarbital, phenytoin or propofol anaesthesia.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

**ATC Code:** CO1D A 14 – Vasodilator used in cardiac diseases.

Isosorbide mononitrate is an organic nitrate, which, in common with other cardioactive nitrates is a vasodilator. It produces decreased left and right ventricular end-diastolic pressures to a greater extent than the decrease in systemic arterial pressure, thereby reducing afterload and especially the preload to the heart.

Isosorbide mononitrate influences the oxygen supply to ischaemic myocardium by causing the redistribution of blood flow along collateral channels and from epicardial to endocardial regions by selective dilation of large epicardial vessels.

It reduces the requirement of the myocardium for oxygen by increasing venous capacitance, causing a pooling of blood in peripheral veins, thereby reducing ventricular volume and heart wall distension.

### 5.2 Pharmacokinetic properties

Isosorbide mononitrate is a vasodilator, which is rapidly absorbed following oral administration. These capsules have a bioavailability of 80 – 90 % when compared to the immediate release isosorbide mononitrate tablets.

The capsules contain pellets which are formulated to release 30 % of the dose immediately whilst 70 % of the dose is released slowly.

Isosorbide mononitrate is extensively metabolised to nitric oxide (NO – which is the active agent) and isosorbide (inactive). In patients with cirrhotic disease or cardiac failure or renal failure, pharmacokinetic parameters were similar to those obtained in healthy

### 5.3 Preclinical safety data

Acute toxicity:

Studies on acute toxicity in mice and rats with different routes of administration indicate a low acute toxicity (LD<sub>50</sub> oral approximately 2000-2500 mg/kg body weight (b.w.)).

Chronic toxicity:

Long term toxicity has been tested in rats for 78 weeks and in dogs for 52 weeks. First toxic reactions occurred in dosages of 90 mg/kg (dog) and 405 mg/kg (rat), respectively. Thus taking into account the recommended dosage of 20 to 30 mg/d in humans, the therapeutic index can be stated as high.

Reproduction studies:

These studies included a fertility and breeding study over two generations in rats; teratology studies in rats and rabbits; and a rat peri-postnatal study. The dosage levels used were generally high and produced maternal toxic effects at the highest dose. No teratogenic effects of isosorbide mononitrate were observed.

Mutagenicity:

Isosorbide mononitrate was tested in different studies in both *in vitro* and *in vivo* (Ames test, Human peripheral lymphocytes, bone marrow of rats and hamsters, V 79 test, SCE test) on possible mutagenic effects. As all tests were negative the mutagenic risks in humans is considered low.

Carcinogenicity:

Neither the long term toxicity studies in rats and dogs nor a special carcinogenicity study, performed in rats over 125 weeks (males) and 138 weeks (females) indicated neoplastigenic properties of isosorbide mononitrate. Therefore, it can be concluded that carcinogenic risk in humans is low.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose monohydrate  
Purified Talc  
Ethyl cellulose  
Macrogol 20000  
Hydroxypropyl cellulose  
Sucrose  
Corn starch  
Gelatin  
Titanium dioxide  
Iron oxide red (E172)  
Iron oxide black (E172)

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

The shelf-life expiry date of this product shall be the date shown on the blister and outer package of the product on the market in the country of origin.

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

Blisters of PVC/aluminium or of polypropylene and aluminium, in an over-labelled outer carton containing calendar packs containing 28 capsules.

### **6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product**

No special requirements.

## **7 PARALLEL PRODUCT AUTHORISATION HOLDER**

IPS Healthcare Limited  
Sterling House  
501 Middleton Road  
Chadderton  
Oldham  
Lancashire OL9 9LY  
United Kingdom

## **8 PARALLEL PRODUCT AUTHORISATION NUMBER**

PPA 1659/050/002

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 11<sup>th</sup> November 2011

**10 DATE OF REVISION OF THE TEXT**