

PACKAGE LEAFLET: INFORMATION FOR THE USER

Buttercup Bronchostop Berry Flavour Cough Pastilles

Thyme herb dry extract

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse after 7 days.

What is in this leaflet:

1. What Buttercup Bronchostop® Berry Flavour Cough Pastilles are and what they are used for
2. What you need to know before you take Buttercup Bronchostop® Berry Flavour Cough Pastilles
3. How to take Buttercup Bronchostop® Berry Flavour Cough Pastilles
4. Possible side effects
5. How to store Buttercup Bronchostop® Berry Flavour Cough Pastilles
6. Contents of the pack and other information

1. WHAT BUTTERCUP BRONCHOSTOP® BERRY FLAVOUR COUGH PASTILLES ARE AND WHAT THEY ARE USED FOR

Buttercup Bronchostop® Berry Flavour Cough Pastilles are a traditional herbal medicinal product used for the relief of coughs, such as chesty, dry, tickly, irritating coughs and catarrh exclusively based on long standing use.

Buttercup Bronchostop® Berry Flavour Pastilles are for use in adults and adolescents over 12 years of age.

You must talk to a doctor if you do not feel better or if you feel worse after 7 days.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE BUTTERCUP BRONCHOSTOP® BERRY FLAVOUR COUGH PASTILLES

Do not take Buttercup Bronchostop® Berry Flavour Cough Pastilles

- If you are allergic to Thyme or other members of the Lamiaceae family (e.g. Basil, Rosemary, Sage), or to any of the other ingredients of this medicine (listed in section 6).

- If you are pregnant or breast-feeding.

Warnings and precautions

Talk to your doctor or pharmacist before taking Buttercup Bronchostop® Berry Flavour Cough Pastilles, if you have asthma or are prone to allergic reactions. There is a possible risk that this medicine may trigger severe allergic reactions in patients with a history of allergic reactions or asthma. Treatment should be discontinued at the first sign of an allergic reaction (See Section 4 for possible side effects).

If you are short of breath, have a high temperature (fever), your phlegm is yellow-green or brown in colour (purulent sputum), you must consult your doctor or qualified healthcare professional.

You must talk to a doctor or qualified healthcare professional if you do not feel better or if you feel worse after 7 days.

Children

Do not give this medicine to children under 12 years.

Other medicines and Buttercup Bronchostop® Berry Flavour Cough Pastilles

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including herbal medicines or medicines obtained without a prescription.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

In the absence of sufficient data, the use of this medicinal product during pregnancy and breast-feeding is not recommended..

Driving and using machines

No studies on the effects on the ability to drive and to use machines have been performed. Buttercup Bronchostop® Berry Flavour Cough Pastilles contain no ingredients that are known to cause drowsiness.

Buttercup Bronchostop® Berry Flavour Cough Pastilles contain fructose, sorbitol, propylene glycol (E1520) and benzyl-alcohol (E1519).

This medicine contains 523 mg of sorbitol and 300 mg of fructose per pastille.

Sorbitol is a source of fructose. If your doctor has told you that you have an intolerance to some sugars or if you have been diagnosed with hereditary fructose intolerance (HFI), a rare genetic disorder in which a person cannot break down fructose, contact your doctor before you take this medicine.

This medicine contains 5.53 mg of propylene-glycol in each pastille.

This medicine contains 0.0018 mg benzyl-alcohol (E1519) in each pastille. Benzyl-alcohol (E1519) may cause allergic reactions. Ask you doctor or pharmacist for advice if you have a liver or kidney disease. This is because large amounts of benzyl alcohol can build-up in your body and may cause side effects (called “metabolic acidosis”).

This medicine contains less than 1 mmol sodium (23 mg) per pastille, that is to say essentially 'sodium-free'.

3. HOW TO TAKE BUTTERCUP BRONCHOSTOP® BERRY FLAVOUR COUGH PASTILLES

Always take Buttercup Bronchostop® Berry Flavour Cough Pastilles exactly as described in this leaflet or as your doctor, pharmacist or qualified Healthcare Professional has told you. Check with your doctor, pharmacist or qualified Healthcare Professional if you are not sure.

Dosage

For oral short-term use only.

Adults, the elderly and children over 12 years:

1 – 2 pastilles to be taken every 4 hours, 4 times a day. If required, up to a maximum of 12 pastilles can be taken per day.

Do not give to children under 12 years.

DO NOT EXCEED THE STATED DOSE.

If you take more Buttercup Bronchostop® Berry Flavour Cough Pastilles than you should

Talk to a doctor straight away, or contact your nearest hospital casualty department. Take the carton and this leaflet with you.

If you forget to take Buttercup Bronchostop® Berry Flavour Cough Pastilles

Take the next dose when it is due. Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Buttercup Bronchostop® Berry Flavour Cough Pastilles can cause side effects, although not everybody gets them.

Allergic reactions including Quincke's oedema and anaphylactic shock have been reported.

This medicine can cause the following serious allergic reactions. If you experience any of the following side effects, stop taking this medicine immediately and seek medical advice:

- Sudden and severe allergic reactions, with breathing difficulty, swelling, lightheadedness, fast heartbeat, sweating and loss of consciousness
- Rapid swelling under the skin in areas such as face, throat, arms and legs
- Difficulty breathing or shortness of breath

Other possible side effects

If any of the following side effects get serious or concern you, or you notice any other side effect, stop taking this medicine and seek medical advice as soon as possible:

- Allergic skin reactions including rash, itching and itchy rash.
- Stomach and gut disorders such as belly pain or discomfort, diarrhoea, vomiting, and nausea.

The frequency of the possible side effects occurring is not known (frequency cannot be estimated from the available data).

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via: HPRA Pharmacovigilance, Website: www.hpra.ie

5. HOW TO STORE BUTTERCUP BRONCHOSTOP® COUGH BERRY FLAVOUR PASTILLES

Do not store above 30°C.

Store the blister in the original package in order to protect from light.

Keep out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and carton. The expiry date refers to the last day of that month.

Do not throw away medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures help protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Buttercup Bronchostop® Berry Flavour Cough Pastilles contain

Each pastille contains the active ingredient 59.5 mg of dry extract from *Thymus vulgaris* L. and *Thymus zygis* L. herba (thyme herb) (7-13:1). Extraction solvent: water

Each pastille also contains 300 mg of fructose and 523 mg of sorbitol

The other ingredients are:

Maltodextrin, Acacia (E414), fructose, sorbitol liquid 70% non-crystallising (E420), Citric acid anhydrous (E330), saccharin sodium (E954), aronia flavour (which contains propylene-glycol (E1520)), fruit of the forest flavour (which contains propylene-glycol (E1520) and benzyl-alcohol (E1519)), paraffin light liquid and purified water.

What Buttercup Bronchostop® Berry Flavour Cough Pastilles look like and contents of the package

Buttercup Bronchostop® Berry Flavour Cough Pastilles are hexagonal, brown pastilles.

Buttercup Bronchostop® Berry Flavour Cough Pastilles are available in blister packs containing 10, 20 or 40 pastilles. Not all pack sizes may be marketed.

TR Holder and manufacturer

Kwizda Pharma GmbH, Effingergasse 21, A-1160 Vienna, Austria

Distributed by
Chefaro Ireland DAC, The Sharp Building, Hogan Place, Dublin 2, Ireland

TR2006/001/002

Is this leaflet hard to see or read?
Phone +44 (0) 1748 827299 for help.

This leaflet was revised in December 2024