

# Summary of Product Characteristics

## 1 NAME OF THE MEDICINAL PRODUCT

Motusol Rx 1 % w/w gel

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

1 g of gel contains diclofenac as 11.6 mg of diclofenac diethylamine corresponding to 10 mg of diclofenac sodium

### Excipient(s) with known effect

1 g of gel contains 50 mg propylene glycol (E1520) and 1 mg fragrance (contains 0.15 mg/g benzyl alcohol, citral, citronellol, coumarin, eugenol, farnesol, geranoil, d-limonene and linalool).

For the full list of excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Gel

White to almost white, homogeneous gel

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic indications

#### **For adults**

For the local symptomatic treatment of pain:

- of acute strains, sprains or contusions following blunt traumas
- of the soft tissues close to the joint (e.g. bursae, tendons, tendon sheaths, ligaments, muscle insertions and joint capsules) in osteoarthritis of knee and finger joints
- in epicondylitis;
- in acute muscular pain, e.g in the back area.

#### **For adolescents aged 14 years and over**

For short-term treatment.

For local, symptomatic treatment of pain in acute strains, sprains or contusions following blunt trauma.

### 4.2 Posology and method of administration

#### Posology

*Adults and adolescents aged 14 years or older:*

Depending on the size of the affected site to be treated, a cherry to walnut-sized quantity, corresponding to 1-4 g of gel (11.6 - 46.4 mg diclofenac diethylamine, corresponding to 10-40 mg diclofenac sodium) should be applied 3-4 times daily. This is sufficient to treat of an area of 400-800 cm<sup>2</sup>.

The maximum daily dose is 16 g of gel corresponding to 185.6 mg of diclofenac, diethylamine salt (corresponding to 160 mg diclofenac sodium).

The duration of use depends on the symptoms and the underlying disease. Motusol Rx 1 % w/w Gel should not be used longer than 1 week without medical advice.

If symptoms worsen or do not improve after 3 - 5 days, a doctor should be consulted.

#### Special patient groups

*Elderly patients:*

No special dose adjustment is required. Because of the potential undesirable-effect profile, elderly people should be carefully monitored.

*Patient with renal impairment:*

No dose reduction is required in patients with renal impairment

*Patient with hepatic impairment:*

No dose reduction is required in patients with hepatic impairment.

*Children and adolescents (under 14 years):*

There are insufficient data on efficacy and safety in children and adolescents under 14 years (see section 4.3).

Method of administration

For cutaneous use.

The gel is applied to the affected parts of the body thinly and rubbed gently into the skin. The hands should be washed unless they are the site being treated.

Before applying a bandage (see section 4.4) the gel should be left to dry for a few minutes on the skin.

### **4.3 Contraindications**

- hypersensitivity to the active substance or to any of the excipients listed in section 6.1
- patients with a history of hypersensitivity reactions, such as asthma, bronchospasmus, urticaria, acute rhinitis or angioedema in response to acetylsalicylic acid or non-steroidal anti-inflammatory drugs (NSAIDs).
- on open injuries, inflammations or infections of the skin as well as on eczema or mucous membranes;
- in the last trimester of pregnancy (see section 4.6);
- in children and adolescents under 14 years of age.

### **4.4 Special warnings and precautions for use**

The possibility of systemic undesirable effects from application of topical diclofenac cannot be excluded if the preparation is used on large areas of skin and over a prolonged period (see product information on systemic forms of diclofenac).

Motusol Rx 1 % w/w Gel must only be applied to intact, not diseased or injured skin. Eyes and oral mucous membranes must not come into contact with the medicinal product and it must not be taken orally.

Topical diclofenac may be used with a non-occlusive bandage but not with an airtight occlusive dressing (see section 5.2)

In acute conditions that are associated with severe redness, swelling, or overheating of joints, in prolonged joint pain or severe back pain that radiates into the legs and / or is associated with neurological deficiencies (e.g. numbness, tingling) a physician should be consulted.

If symptoms worsen or do not improve after 3 – 5 days, a doctor should be consulted.

Patients suffering from asthma, hay fever, swelling of nasal mucous membranes (so-called nasal polyps) or chronic obstructive pulmonary disease, chronic respiratory infections (particularly associated with hay fever-like symptoms), and patients with hypersensitivity to painkillers and anti-rheumatic medicinal products of all kinds are rather at risk to asthma attacks (so called analgesic intolerance / analgesic asthma), to local skin or mucous membrane swelling (so-called quincke edema) or to urticaria - than other patients when treated with Motusol Rx 1 % w/w Gel.

In these patients, Motusol Rx 1 % w/w Gel may only be used under certain precautions (emergency preparedness) and direct medical supervision. The same applies for patients who are also allergic to other substances e.g. with skin reactions, itching or urticaria.

If a skin rash occurs during the treatment with Motusol Rx 1 % w/w Gel, the treatment should be stopped.

During treatment photosensitivity can occur with the appearance of skin reactions after exposition to sunlight.

Preventative measures should be taken so that children do not contact the skin areas to which the gel has been applied.

This medicinal product contains fragrance with benzyl alcohol (0.15 mg/g), citral, citronellol, coumarin, eugenol, farnesol, geraniol, d-limonene and linalool which may cause allergic reactions.

In addition, benzyl alcohol may cause mild local irritation.

#### 4.5 Interaction with other medicinal products and other forms of interaction

Since the systemic absorption of diclofenac is very low with topical application, interactions are very unlikely in use as intended.

#### 4.6 Fertility, pregnancy and lactation

The systemic concentration of diclofenac is lower after topical administration, compared to oral formulations. With reference to experience from treatment with NSAIDs with systemic uptake, the following is recommended:

##### Pregnancy

Inhibition of prostaglandin synthesis may adversely affect the pregnancy and/or the embryo/fetal development. Data from epidemiological studies suggest an increased risk of miscarriage and of cardiac malformation and gastroschisis after use of a prostaglandin synthesis inhibitor in early pregnancy. The absolute risk for cardiovascular malformation was increased from less than 1%, up to approximately 1.5 %. The risk is believed to increase with dose and duration of therapy. In animals, administration of a prostaglandin synthesis inhibitor has been shown to result in increased pre- and post-implantation loss and embryo-fetal lethality. In addition, increased incidences of various malformations, including cardiovascular, have been reported in animals given a prostaglandin synthesis inhibitor during the organogenetic period.

During the first and second trimester of pregnancy, diclofenac should not be used unless clearly necessary. If diclofenac is used by a woman attempting to conceive, or during the first and second trimester of pregnancy, the dose should be kept as low and duration of treatment as short as possible.

During the third trimester of pregnancy, all prostaglandin synthesis inhibitors may expose

- the fetus to:
  - cardiopulmonary toxicity (with premature closure of the ductus arteriosus and pulmonary hypertension);
  - renal dysfunction, which may progress to renal failure with oligo hydroamniosis;
- the mother and the neonate, at the end of pregnancy, to:
  - possible prolongation of bleeding time, an anti-aggregating effect which may occur even at very low doses.
  - inhibition of uterine contractions resulting in delayed or prolonged labour.

Consequently, diclofenac is contraindicated during the third trimester of pregnancy.

##### Breast-feeding

Diclofenac passes into breast milk in small amounts. However, at therapeutic doses of Motusol Rx 1 % w/w Gel no effects on the breast-fed child are anticipated. Because of a lack of controlled studies in breast-feeding women, the medicinal product should only be used during breast-feeding under advice from a healthcare professional. Under this circumstance, Motusol Rx 1% w/w Gel should not be applied on the breasts of breast-feeding mothers, nor elsewhere on large areas of skin or for a prolonged period of time (see section 4.4).

#### 4.7 Effects on ability to drive and use machines

The topical use of diclofenac has no or negligible influence on the ability to drive and use machines.

#### 4.8 Undesirable effects

Adverse reactions are listed below, by system organ class and frequency. Frequencies are defined as: *Very common* ( $\geq 1/10$ ); *Common* ( $\geq 1/100$  to  $< 1/10$ ); *Uncommon* ( $\geq 1/1,000$  to  $< 1/100$ ); *Rare* ( $\geq 1/10,000$  to  $< 1/1,000$ ); *Very rare* ( $< 1/10,000$ ), *Not known* (cannot be estimated from the available data).

System organ class database	Adverse reactions and frequency
<b>Infections and infestations</b>	<i>Very rare</i> : Rash pustular

<b>Immune system disorders</b>	<i>Very rare:</i> Hypersensitivity (including urticaria), angioedema
<b>Respiratory, thoracic and mediastinal disorders</b>	<i>Very rare:</i> Asthma
<b>Gastrointestinal disorders</b>	<i>Very rare:</i> Gastrointestinal complaints
<b>Skin and subcutaneous tissue disorders</b>	<i>Common:</i> Dermatitis (including contact dermatitis), skin rash, erythema, eczema, pruritus <i>Uncommon:</i> Scaling, dehydration of the skin, oedema <i>Rare:</i> Dermatitis bullous <i>Very rare:</i> Photosensitivity reaction <i>Not known:</i> Burning sensation at the application site, dry skin

When the gel is applied on large areas of skin and over a prolonged period, the possibility of systemic undesirable effects (e.g. renal, hepatic or gastrointestinal undesirable effects, systemic hypersensitivity reactions) - as they occur possibly after systemic administration of diclofenac-containing medicinal products cannot be excluded.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the HPRA Pharmacovigilance

Website: [www.hpra.ie](http://www.hpra.ie).

#### **4.9 Overdose**

Due to the low systemic absorption of diclofenac in limited topical use an overdose is unlikely.

If the recommended dose is significantly exceeded, the gel should be removed from the skin and washed off with water.

Undesirable effects similar to those observed following an overdose of systemic diclofenac can occur if topical diclofenac is inadvertently ingested (1 tube of 100 g contains the equivalent of 1,160 diclofenac diethylamine corresponding to 1,000 mg diclofenac sodium).

In the event of accidental ingestion, resulting in significant systemic adverse effects, general therapeutic measures normally adopted to treat poisoning with non-steroidal anti-inflammatory medicinal products should be used. Gastric lavage and the use of activated charcoal should be considered, especially within a short time of ingestion.

A specific antidote does not exist.

### **5 PHARMACOLOGICAL PROPERTIES**

#### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Topical products for joint and muscular pain; Anti-inflammatory preparations, non-steroids for topical use

ATC code: M02AA15

#### Mechanism of action

Diclofenac is a potent non-steroidal anti-inflammatory drug. It develops its therapeutic efficacy mainly via inhibition of prostaglandin synthesis by cyclooxygenase 2 (COX-2). Diclofenac has proven to be effective via the prostaglandin synthesis inhibition in the conventional animal-experiment inflammation models. In humans, diclofenac reduces inflammatory-related pain, swellings and fever. Furthermore, diclofenac inhibits reversibly the ADP and the collagen-induced thrombocyte aggregation.

#### **5.2 Pharmacokinetic properties**

##### Absorption

The quantity of diclofenac absorbed through the skin is proportional to the duration of the skin contact and the size of the treated area, and depends on both the total dose applied and the degree of skin hydration. After local application of Motusol Rx 1 % w/w Gelto hand and knee joints, the active substance is absorbed through the skin and detectable in the plasma as well as the tissue in varying quantities – depending on the diffusion range – beneath the application site.

Absorption amounts to about 6 % of the applied dose of diclofenac after topical application of 2.5 g diclofenac gel on 500 cm<sup>2</sup> skin, determined by measuring total renal elimination of diclofenac and its hydroxylated metabolites, compared with the oral administration of diclofenac sodium. Due to a depot-effect in the skin, there is a delayed and prolonged release of active substance into the underlying tissue and the plasma. Under occlusive conditions (10 hours), percutaneous absorption of diclofenac in adults can be increased three-fold (serum concentration).

#### Distribution

99.7 % of diclofenac is bound to serum proteins, mainly albumin (99.4 %). Plasma levels after application of diclofenac gel are not sufficient to explain the observed therapeutic efficacy; this is more likely due to the presence of significantly higher active substance concentrations beneath the application site. Due to its properties (such as short plasma half-life, low pKa value, small distribution volume and high protein binding), diclofenac has an affinity to inflamed tissue.

Diclofenac preferentially distributes and persists in inflamed tissue it is found in concentrations up to 20 times higher than in plasma.

#### Biotransformation

Biotransformation of diclofenac involves partly glucuronidation of the intact molecule, but mainly single and multiple hydroxylation resulting in several phenolic metabolites, most of which are converted to glucuronide conjugates. Two of the phenolic metabolites are biologically active, however, to a much smaller extent than diclofenac.

#### Elimination

The total systemic clearance of diclofenac from plasma is 263 ± 56 ml/min. The terminal plasma half-life is 1-2 hours. Four of the metabolites, including the two active ones, also have short plasma half-lives of 1-3 hours. One metabolite, 3'-hydroxy-4'-methoxy-diclofenac, has a longer half-life but is virtually inactive. Diclofenac and its metabolites are excreted mainly in the urine.

Characteristics in patients:

No accumulation of diclofenac and its metabolites is to be expected in patients suffering from renal impairment. In patients with chronic hepatitis or nondecompensated cirrhosis, the kinetics and metabolism of diclofenac are the same as in patients without liver disease.

### **5.3 Preclinical safety data**

Based on conventional studies on safety pharmacology, genotoxicity and carcinogenic potential, the preclinical data do not reveal any specific hazards for humans, apart from those already described in other sections of the SPC. In animal studies, the chronic toxicity of diclofenac following systemic application mainly manifested as gastrointestinal lesions and ulcers. In a 2-year toxicity study, a dose-dependent increase in the incidence of thrombosis of the heart was observed in diclofenac-treated rats.

In animal studies on reproductive toxicity, systemically administered diclofenac caused inhibition of ovulation in rabbits and impairment of implantation and early embryonic development in rats. Gestation and duration of parturition were prolonged by diclofenac. The embryotoxic potential of diclofenac was investigated in three animal species (rat, mouse, rabbit). Fetal death and growth retardation occurred at materno-toxic dose levels. Based on the available non-clinical data, diclofenac is regarded as being non-teratogenic. Doses below the maternotoxic threshold had no impact on the postnatal development of the offspring.

Diclofenac poses a risk to the aquatic environment (see section 6.6).

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Carbomer

Cocoyl Caprylocaprate

Macrogol cetostearyl ether

Paraffin, Liquid

Diethylamine

Isopropyl alcohol

Propylene glycol (E1520)

Fragrance (contains citronellol, geraniol, benzyl alcohol, linalool, limonene, citral, farnesol, coumarin, eugenol)

Purified water

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

2 years

## **6.4 Special precautions for storage**

*Store in the original tube in order to protect from light.*

This medicinal product does not require any special temperature storage conditions.

## **6.5 Nature and contents of container**

The gel is packed in aluminium laminated tubes, closed with PE seal and PP screw caps, in pack sizes: 30 g, 50 g, 60 g, 100 g, 120 g, 150 g per tube.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal and other handling**

This medicinal product poses a risk to the environment (see section 5.3).

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7 MARKETING AUTHORISATION HOLDER**

Teva B.V.  
Swensweg 5  
2031GA Haarlem  
Netherlands

## **8 MARKETING AUTHORISATION NUMBER**

PA1986/093/001

## **9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 25<sup>th</sup> August 2020

Date of last renewal: 10<sup>th</sup> June 2025

## **10 DATE OF REVISION OF THE TEXT**

January 2025