

# Summary of Product Characteristics

## 1 NAME OF THE MEDICINAL PRODUCT

Sitagliptin/Metformin hydrochloride Krka 50 mg/1000 mg film-coated tablets

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 50 mg sitagliptin and 1000 mg of metformin hydrochloride.

For the full list of excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Film-coated tablet (tablet)

Dark pink, oval, biconvex, film coated tablets marked with C3 on one side of the tablet (dimensions approx.: 21 x 11 mm).

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic indications

For adult patients with type 2 diabetes mellitus:

Sitagliptin/Metformin hydrochloride Krka is indicated as an adjunct to diet and exercise to improve glycaemic control in patients inadequately controlled on their maximal tolerated dose of metformin alone or those already being treated with the combination of sitagliptin and metformin.

Sitagliptin/Metformin hydrochloride Krka is indicated in combination with a sulphonylurea (i.e., triple combination therapy) as an adjunct to diet and exercise in patients inadequately controlled on their maximal tolerated dose of metformin and a sulphonylurea.

Sitagliptin/Metformin hydrochloride Krka is indicated as triple combination therapy with a peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) agonist (i.e., a thiazolidinedione) as an adjunct to diet and exercise in patients inadequately controlled on their maximal tolerated dose of metformin and a PPAR $\gamma$  agonist.

Sitagliptin/Metformin hydrochloride Krka is also indicated as add-on to insulin (i.e., triple combination therapy) as an adjunct to diet and exercise to improve glycaemic control in patients when stable dose of insulin and metformin alone do not provide adequate glycaemic control.

### 4.2 Posology and method of administration

#### Posology

The dose of antihyperglycaemic therapy with Sitagliptin/Metformin hydrochloride Krka should be individualised on the basis of the patient's current regimen, effectiveness, and tolerability while not exceeding the maximum recommended daily dose of 100 mg sitagliptin.

#### ***Adults with normal renal function (GFR $\geq$ 90 mL/min)***

##### For patients inadequately controlled on maximal tolerated dose of metformin monotherapy

For patients not adequately controlled on metformin alone, the usual starting dose should provide sitagliptin dosed as 50 mg twice daily (100 mg total daily dose) plus the dose of metformin already being taken.

##### For patients switching from co-administration of sitagliptin and metformin

For patients switching from co-administration of sitagliptin and metformin, Sitagliptin/Metformin hydrochloride Krka should be initiated at the dose of sitagliptin and metformin already being taken.

For patients inadequately controlled on dual combination therapy with the maximal tolerated dose of metformin and a sulphonylurea

The dose should provide sitagliptin dosed as 50 mg twice daily (100 mg total daily dose) and a dose of metformin similar to the dose already being taken. When Sitagliptin/Metformin hydrochloride Krka is used in combination with a sulphonylurea, a lower dose of the sulphonylurea may be required to reduce the risk of hypoglycaemia (see section 4.4).

For patients inadequately controlled on dual combination therapy with the maximal tolerated dose of metformin and a PPAR $\gamma$  agonist

The dose should provide sitagliptin dosed as 50 mg twice daily (100 mg total daily dose) and a dose of metformin similar to the dose already being taken.

For patients inadequately controlled on dual combination therapy with insulin and the maximal tolerated dose of metformin

The dose should provide sitagliptin dosed as 50 mg twice daily (100 mg total daily dose) and a dose of metformin similar to the dose already being taken. When Sitagliptin/Metformin hydrochloride Krka is used in combination with insulin, a lower dose of insulin may be required to reduce the risk of hypoglycaemia (see section 4.4).

For the different doses on metformin, Sitagliptin/Metformin hydrochloride Krka is available in strengths of 50 mg sitagliptin and 850 mg metformin hydrochloride or 1,000 mg metformin hydrochloride.

All patients should continue their recommended diet with an adequate distribution of carbohydrate intake during the day.

*Special populations**Renal impairment*

No dose adjustment is needed for patients with mild renal impairment (glomerular filtration rate [GFR]  $\geq$  60 mL/min). A GFR should be assessed before initiation of treatment with metformin containing products and at least annually thereafter. In patients at increased risk of further progression of renal impairment and in the elderly, renal function should be assessed more frequently, e.g. every 3-6 months.

The maximum daily dose of metformin should preferably be divided into 2-3 daily doses. Factors that may increase the risk of lactic acidosis (see section 4.4) should be reviewed before considering initiation of metformin in patients with GFR < 60 mL/min.

If no adequate strength of Sitagliptin/Metformin hydrochloride Krka is available, individual monocomponents should be used instead of the fixed-dose combination.

<b>GFR mL/min</b>	<b><u>Metformin</u></b>	<b><u>Sitagliptin</u></b>
60-89	<i>Maximum daily dose is 3000 mg. Dose reduction may be considered in relation to declining renal function.</i>	<i>Maximum daily dose is 100 mg.</i>
45-59	<i>Maximum daily dose is 2000 mg. The starting dose is at most half of the maximum dose.</i>	<i>Maximum daily dose is 100 mg.</i>
30-44	<i>Maximum daily dose is 1000 mg. The starting dose is at most half of the maximum dose.</i>	<i>Maximum daily dose is 50 mg.</i>
< 30	<i>Metformin is contraindicated.</i>	<i>Maximum daily dose is 25 mg.</i>

*Hepatic impairment*

Sitagliptin/Metformin hydrochloride Krka must not be used in patients with hepatic impairment (see section 5.2).

*Elderly*

As metformin and sitagliptin are excreted by the kidney, Sitagliptin/Metformin hydrochloride Krka should be used with caution as age increases. Monitoring of renal function is necessary to aid in prevention of metformin-associated lactic acidosis, particularly in the elderly (see sections 4.3 and 4.4).

*Paediatric population*

Sitagliptin/Metformin hydrochloride Krka should not be used in children and adolescents 10 to 17 years of age because of insufficient efficacy. Currently available data are described in sections 4.8, 5.1, and 5.2. Sitagliptin/Metformin hydrochloride Krka has not been studied in paediatric patients under 10 years of age.

#### Method of administration

Sitagliptin/Metformin hydrochloride Krka should be given twice daily with meals to reduce the gastrointestinal adverse reactions associated with metformin.

### **4.3 Contraindications**

Sitagliptin/Metformin hydrochloride Krka is contraindicated in patients with:

- hypersensitivity to the active substances or to any of the excipients listed in section 6.1 (see sections 4.4. and 4.8).
- any type of acute metabolic acidosis (such as lactic acidosis, diabetic ketoacidosis);
- diabetic pre-coma;
- severe renal failure (GFR < 30 mL/min) (see section 4.4);
- acute conditions with the potential to alter renal function such as:
  - dehydration,
  - severe infection,
  - shock,
  - intravascular administration of iodinated contrast agents (see section 4.4);
- acute or chronic disease which may cause tissue hypoxia such as:
  - cardiac or respiratory failure,
  - recent myocardial infarction,
  - shock.
- hepatic impairment;
- acute alcohol intoxication, alcoholism;
- breast-feeding.

### **4.4 Special warnings and precautions for use**

#### General

Sitagliptin/Metformin hydrochloride Krka should not be used in patients with type 1 diabetes and must not be used for the treatment of diabetic ketoacidosis.

#### Acute pancreatitis

Use of DPP-4 inhibitors has been associated with a risk of developing acute pancreatitis. Patients should be informed of the characteristic symptom of acute pancreatitis: persistent, severe abdominal pain. Resolution of pancreatitis has been observed after discontinuation of sitagliptin (with or without supportive treatment), but very rare cases of necrotising or haemorrhagic pancreatitis and/or death have been reported. If pancreatitis is suspected, Sitagliptin/Metformin hydrochloride Krka and other potentially suspect medicinal products should be discontinued; if acute pancreatitis is confirmed, Sitagliptin/Metformin hydrochloride Krka should not be restarted.

Caution should be exercised in patients with a history of pancreatitis.

#### Lactic acidosis

Lactic acidosis, a rare but serious metabolic complication, most often occurs at acute worsening of renal function or cardiorespiratory illness or sepsis. Metformin accumulation occurs at acute worsening of renal function and increases the risk of lactic acidosis.

In case of dehydration (severe vomiting, diarrhoea, fever or reduced fluid intake), metformin should be temporarily discontinued and contact with a health care professional is recommended.

Medicinal products that can acutely impair renal function (such as antihypertensives, diuretics and NSAIDs) should be initiated with caution in metformin-treated patients. Other risk factors for lactic acidosis are excessive alcohol intake, hepatic insufficiency, inadequately controlled diabetes, ketosis, prolonged fasting and any conditions associated with hypoxia, as well as concomitant use of medicinal products that may cause lactic acidosis (see sections 4.3 and 4.5).

Patients and/or care-givers should be informed of the risk of lactic acidosis. Lactic acidosis is characterised by acidotic dyspnoea, abdominal pain, muscle cramps, asthenia and hypothermia followed by coma. In case of suspected symptoms, the patient should stop taking metformin and seek immediate medical attention. Diagnostic laboratory findings are decreased blood pH (< 7.35), increased plasma lactate levels (> 5 mmol/L) and an increased anion gap and lactate/pyruvate ratio.

### Patients with known or suspected mitochondrial diseases

In patients with known mitochondrial diseases such as Mitochondrial Encephalopathy with Lactic Acidosis, and Stroke-like episodes (MELAS) syndrome and Maternal inherited diabetes and deafness (MIDD), metformin is not recommended due to the risk of lactic acidosis exacerbation and neurologic complications which may lead to worsening of the disease.

In case of signs and symptoms suggestive of MELAS syndrome or MIDD after the intake of metformin, treatment with metformin should be withdrawn immediately and prompt diagnostic evaluation should be performed.

### Renal function

GFR should be assessed before treatment initiation and regularly thereafter (see section 4.2). Sitagliptin/Metformin hydrochloride Krka is contraindicated in patients with GFR < 30 mL/min and should be temporarily discontinued during conditions with the potential to alter renal function (see section 4.3).

### Hypoglycaemia

Patients receiving Sitagliptin/Metformin hydrochloride Krka in combination with a sulphonylurea or with insulin may be at risk for hypoglycaemia. Therefore, a reduction in the dose of the sulphonylurea or insulin may be necessary.

### Hypersensitivity reactions

Post-marketing reports of serious hypersensitivity reactions in patients treated with sitagliptin have been reported. These reactions include anaphylaxis, angioedema, and exfoliative skin conditions including Stevens-Johnson syndrome. Onset of these reactions occurred within the first 3 months after initiation of treatment with sitagliptin, with some reports occurring after the first dose. If a hypersensitivity reaction is suspected, Sitagliptin/Metformin hydrochloride Krka should be discontinued, other potential causes of the event should be assessed, and alternative treatment for diabetes should be instituted (see section 4.8).

### Bullous pemphigoid

There have been post-marketing reports of bullous pemphigoid in patients taking DPP-4 inhibitors including sitagliptin. If bullous pemphigoid is suspected, Sitagliptin/Metformin hydrochloride Krka should be discontinued.

### Surgery

Sitagliptin/Metformin hydrochloride Krka must be discontinued at the time of surgery under general, spinal or epidural anaesthesia. Therapy may be restarted no earlier than 48 hours following surgery or resumption of oral nutrition and provided that renal function has been re-evaluated and found to be stable.

### Administration of iodinated contrast agent

Intravascular administration of iodinated contrast agents may lead to contrast-induced nephropathy, resulting in metformin accumulation and an increased risk of lactic acidosis. Sitagliptin/Metformin hydrochloride Krka should be discontinued prior to or at the time of the imaging procedure and not restarted until at least 48 hours after, provided that renal function has been re-evaluated and found to be stable (see sections 4.3 and 4.5).

### Change in clinical status of patients with previously controlled type 2 diabetes

A patient with type 2 diabetes previously well controlled on Sitagliptin/Metformin hydrochloride Krka who develops laboratory abnormalities or clinical illness (especially vague and poorly defined illness) should be evaluated promptly for evidence of ketoacidosis or lactic acidosis. Evaluation should include serum electrolytes and ketones, blood glucose and, if indicated, blood pH, lactate, pyruvate, and metformin levels. If acidosis of either form occurs, treatment must be stopped immediately and other appropriate corrective measures initiated.

### Vitamin B<sub>12</sub> Deficiency

Metformin may reduce vitamin B<sub>12</sub> serum levels. The risk of low vitamin B<sub>12</sub> levels increases with increasing metformin dose, treatment duration, and/or in patients with risk factors known to cause vitamin B<sub>12</sub> deficiency. In case of suspicion of vitamin B<sub>12</sub> deficiency (such as anaemia or neuropathy), vitamin B<sub>12</sub> serum levels should be monitored. Periodic vitamin B<sub>12</sub> monitoring could be necessary in patients with risk factors for vitamin B<sub>12</sub> deficiency. Metformin therapy should be continued for as long as it is tolerated and not contra-indicated and appropriate corrective treatment for vitamin B<sub>12</sub> deficiency provided in line with current clinical guidelines.

Sitagliptin/Metformin hydrochloride Krka contains sodium.

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

## **4.5 Interaction with other medicinal products and other forms of interaction**

Co-administration of multiple doses of sitagliptin (50 mg twice daily) and metformin (1,000 mg twice daily) did not meaningfully alter the pharmacokinetics of either sitagliptin or metformin in patients with type 2 diabetes.

Pharmacokinetic drug interaction studies with Sitagliptin/Metformin hydrochloride Krka have not been performed; however, such studies have been conducted with the individual active substances, sitagliptin and metformin.

#### Concomitant use not recommended

##### Alcohol

Alcohol intoxication is associated with an increased risk of lactic acidosis, particularly in cases of fasting, malnutrition or hepatic impairment.

##### Iodinated contrast agents

Sitagliptin/Metformin hydrochloride Krka must be discontinued prior to or at the time of the imaging procedure and not restarted until at least 48 hours after, provided that renal function has been re-evaluated and found to be stable (see sections 4.3 and 4.4).

#### Combinations requiring precautions for use

Some medicinal products can adversely affect renal function, which may increase the risk of lactic acidosis, e.g. NSAIDs, including selective cyclo-oxygenase (COX) II inhibitors, ACE inhibitors, angiotensin II receptor antagonists and diuretics, especially loop diuretics. When starting or using such products in combination with metformin, close monitoring of renal function is necessary.

Concomitant use of medicinal products that interfere with common renal tubular transport systems involved in the renal elimination of metformin (e.g., organic cationic transporter-2 [OCT2] / multidrug and toxin extrusion [MATE] inhibitors such as ranolazine, vandetanib, dolutegravir, and cimetidine) could increase systemic exposure to metformin and may increase the risk for lactic acidosis. Consider the benefits and risks of concomitant use. Close monitoring of glycaemic control, dose adjustment within the recommended posology and changes in diabetic treatment should be considered when such products are co-administered.

Glucocorticoids (given by systemic and local routes) beta-2-agonists, and diuretics have intrinsic hyperglycaemic activity. The patient should be informed and more frequent blood glucose monitoring performed, especially at the beginning of treatment with such medicinal products. If necessary, the dose of the anti-hyperglycaemic medicinal product should be adjusted during therapy with the other medicinal product and on its discontinuation.

ACE-inhibitors may decrease the blood glucose levels. If necessary, the dose of the anti-hyperglycaemic medicinal product should be adjusted during therapy with the other medicinal product and on its discontinuation.

#### Effects of other medicinal products on sitagliptin

*In vitro* and clinical data described below suggest that the risk for clinically meaningful interactions following co-administration of other medicinal products is low.

*In vitro* studies indicated that the primary enzyme responsible for the limited metabolism of sitagliptin is CYP3A4, with contribution from CYP2C8. In patients with normal renal function, metabolism, including via CYP3A4, plays only a small role in the clearance of sitagliptin. Metabolism may play a more significant role in the elimination of sitagliptin in the setting of severe renal impairment or end-stage renal disease (ESRD). For this reason, it is possible that potent CYP3A4 inhibitors (i.e., ketoconazole, itraconazole, ritonavir, clarithromycin) could alter the pharmacokinetics of sitagliptin in patients with severe renal impairment or ESRD. The effects of potent CYP3A4 inhibitors in the setting of renal impairment have not been assessed in a clinical study.

*In vitro* transport studies showed that sitagliptin is a substrate for p-glycoprotein and organic anion transporter-3 (OAT3). OAT3 mediated transport of sitagliptin was inhibited *in vitro* by probenecid, although the risk of clinically meaningful interactions is considered to be low. Concomitant administration of OAT3 inhibitors has not been evaluated *in vivo*.

*Ciclosporin*: A study was conducted to assess the effect of ciclosporin, a potent inhibitor of p-glycoprotein, on the pharmacokinetics of sitagliptin. Co-administration of a single 100 mg oral dose of sitagliptin and a single 600 mg oral dose of ciclosporin increased the AUC and  $C_{max}$  of sitagliptin by approximately 29 % and 68 %, respectively. These changes in sitagliptin pharmacokinetics were not considered to be clinically meaningful. The renal clearance of sitagliptin was not meaningfully altered. Therefore, meaningful interactions would not be expected with other p-glycoprotein inhibitors.

Effects of sitagliptin on other medicinal products

*Digoxin:* Sitagliptin had a small effect on plasma digoxin concentrations. Following administration of 0.25 mg digoxin concomitantly with 100 mg of sitagliptin daily for 10 days, the plasma AUC of digoxin was increased on average by 11 %, and the plasma  $C_{max}$  on average by 18 %. No dose adjustment of digoxin is recommended. However, patients at risk of digoxin toxicity should be monitored for this when sitagliptin and digoxin are administered concomitantly.

*In vitro* data suggest that sitagliptin does not inhibit nor induce CYP450 isoenzymes. In clinical studies, sitagliptin did not meaningfully alter the pharmacokinetics of metformin, glyburide, simvastatin, rosiglitazone, warfarin, or oral contraceptives, providing *in vivo* evidence of a low propensity for causing interactions with substrates of CYP3A4, CYP2C8, CYP2C9, and organic cationic transporter (OCT). Sitagliptin may be a mild inhibitor of p-glycoprotein *in vivo*.

**4.6 Fertility, pregnancy and lactation**Pregnancy

There are no adequate data from the use of sitagliptin in pregnant women. Studies in animals have shown reproductive toxicity at high doses of sitagliptin (see section 5.3).

A limited amount of data suggests the use of metformin in pregnant women is not associated with an increased risk of congenital malformations. Animal studies with metformin do not indicate harmful effects with respect to pregnancy, embryonic or foetal development, parturition or postnatal development (see also section 5.3).

Sitagliptin/Metformin hydrochloride Krka should not be used during pregnancy. If a patient wishes to become pregnant or if a pregnancy occurs, treatment should be discontinued and the patient switched to insulin treatment as soon as possible.

Breast-feeding

No studies in lactating animals have been conducted with the combined active substances of this medicinal product. In studies performed with the individual active substances, both sitagliptin and metformin are excreted in the milk of lactating rats. Metformin is excreted in human milk in small amounts. It is not known whether sitagliptin is excreted in human milk. Sitagliptin/Metformin hydrochloride Krka must therefore not be used in women who are breast-feeding (see section 4.3).

Fertility

Animal data do not suggest an effect of treatment with sitagliptin on male and female fertility. Human data are lacking.

**4.7 Effects on ability to drive and use machines**

Sitagliptin/Metformin hydrochloride Krka has no or negligible influence on the ability to drive and use machines. However, when driving or using machines, it should be taken into account that dizziness and somnolence have been reported with sitagliptin.

In addition, patients should be alerted to the risk of hypoglycaemia when Sitagliptin/Metformin hydrochloride Krka is used in combination with a sulphonylurea or with insulin.

**4.8 Undesirable effects**Summary of the safety profile

There have been no therapeutic clinical studies conducted with sitagliptin/metformin combination however bioequivalence of sitagliptin/metformin combination with co-administered sitagliptin and metformin has been demonstrated (see section 5.2). Serious adverse reactions including pancreatitis and hypersensitivity reactions have been reported. Hypoglycaemia has been reported in combination with sulphonylurea (13.8 %) and insulin (10.9 %).

*Sitagliptin and metformin*Tabulated list of adverse reactions

Adverse reactions are listed below as MedDRA preferred term by system organ class and absolute frequency (Table 1). Frequencies are defined as: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1\ 000$  to  $< 1/100$ ); rare ( $\geq 1/10\ 000$  to  $< 1/1\ 000$ ); very rare ( $< 1/10\ 000$ ) and not known (cannot be estimated from the available data).

**Table 1: The frequency of adverse reactions identified from placebo-controlled clinical studies of sitagliptin and metformin alone, and post-marketing experience**

Adverse reaction	Frequency of adverse reaction
<b>Blood and lymphatic system disorders</b>	
thrombocytopenia	Rare
<b>Immune system disorders</b>	
hypersensitivity reactions including anaphylactic responses <sup>*,†</sup>	Frequency not known
<b>Metabolism and nutrition disorders</b>	
hypoglycaemia <sup>†</sup>	Common
Vitamin B <sub>12</sub> decrease/defficiency <sup>†</sup>	Common
<b>Nervous system disorders</b>	
somnolence	Uncommon
<b>Respiratory, thoracic and mediastinal disorders</b>	
interstitial lung disease <sup>*</sup>	Frequency not known
<b>Gastrointestinal disorders</b>	
diarrhoea	Uncommon
nausea	Common
flatulence	Common
constipation	Uncommon
upper abdominal pain	Uncommon
vomiting	Common
acute pancreatitis <sup>*,†,‡</sup>	Frequency not known
fatal and non-fatal haemorrhagic and necrotizing pancreatitis <sup>*,†</sup>	Frequency not known
<b>Skin and subcutaneous tissue disorders</b>	
pruritus <sup>*</sup>	Uncommon
angioedema <sup>*,†</sup>	Frequency not known
rash <sup>*,†</sup>	Frequency not known
urticaria <sup>*,†</sup>	Frequency not known
cutaneous vasculitis <sup>*,†</sup>	Frequency not known
exfoliative skin conditions including Stevens-Johnson syndrome <sup>*,†</sup>	Frequency not known
bullous pemphigoid <sup>*</sup>	Frequency not known
<b>Musculoskeletal and connective tissue disorders</b>	
arthralgia <sup>*</sup>	Frequency not known
myalgia <sup>*</sup>	Frequency not known
pain in extremity <sup>*</sup>	Frequency not known
back pain <sup>*</sup>	Frequency not known
arthropathy <sup>*</sup>	Frequency not known
<b>Renal and urinary disorders</b>	
impaired renal function <sup>*</sup>	Frequency not known
acute renal failure <sup>*</sup>	Frequency not known

<sup>\*</sup>Adverse reactions were identified through post-marketing surveillance.

<sup>†</sup>See section 4.4.

<sup>‡</sup> See *TECOS Cardiovascular Safety Study* below.

Description of selected adverse reactions

Some adverse reactions were observed more frequently in studies of combination use of sitagliptin and metformin with other anti-diabetic medicinal products than in studies of sitagliptin and metformin alone. These included hypoglycaemia (frequency very common with sulphonylurea or insulin), constipation (common with sulphonylurea), peripheral oedema (common with pioglitazone), and headache and dry mouth (uncommon with insulin).

#### *Sitagliptin*

In monotherapy studies of sitagliptin 100 mg once daily alone compared to placebo, adverse reactions reported were headache, hypoglycaemia, constipation, and dizziness.

Among these patients, adverse events reported regardless of causal relationship to medicinal product occurring in at least 5 % included upper respiratory tract infection and nasopharyngitis. In addition, osteoarthritis and pain in extremity were reported with frequency uncommon (> 0.5 % higher among sitagliptin users than that in the control group).

#### *Metformin*

Gastrointestinal symptoms were reported very commonly in clinical studies and post-marketing use of metformin.

Gastrointestinal symptoms such as nausea, vomiting, diarrhoea, abdominal pain and loss of appetite occur most frequently during initiation of therapy and resolve spontaneously in most cases. Additional adverse reactions associated with metformin include metallic taste (common); lactic acidosis, liver function disorders, hepatitis, urticaria, erythema, and pruritus (very rare). Frequency categories are based on information available from metformin Summary of Product Characteristics available in the EU.

#### *Paediatric population*

In clinical studies with sitagliptin/metformin in paediatric patients with type 2 diabetes mellitus aged 10 to 17 years, the profile of adverse reactions was generally comparable to that observed in adults. In paediatric patients on or not on background insulin, sitagliptin was associated with an increased risk of hypoglycaemia.

#### *TECOS Cardiovascular Safety Study*

The Trial Evaluating Cardiovascular Outcomes with Sitagliptin (TECOS) included 7,332 patients treated with sitagliptin, 100 mg daily (or 50 mg daily if the baseline eGFR was  $\geq 30$  and  $< 50$  mL/min/1.73 m<sup>2</sup>), and 7,339 patients treated with placebo in the intention-to-treat population. Both treatments were added to usual care targeting regional standards for HbA<sub>1c</sub> and CV risk factors. The overall incidence of serious adverse events in patients receiving sitagliptin was similar to that in patients receiving placebo.

In the intention-to-treat population, among patients who were using insulin and/or a sulphonylurea at baseline, the incidence of severe hypoglycaemia was 2.7 % in sitagliptin-treated patients and 2.5 % in placebo-treated patients; among patients who were not using insulin and/or a sulphonylurea at baseline, the incidence of severe hypoglycaemia was 1.0 % in sitagliptin-treated patients and 0.7 % in placebo-treated patients. The incidence of adjudication-confirmed pancreatitis events was 0.3 % in sitagliptin-treated patients and 0.2 % in placebo-treated patients.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance. Website: [www.hpra.ie](http://www.hpra.ie).

## **4.9 Overdose**

During controlled clinical studies in healthy subjects, single doses of up to 800 mg sitagliptin were administered. Minimal increases in QTc, not considered to be clinically relevant, were observed in one study at a dose of 800 mg sitagliptin. There is no experience with doses above 800 mg in clinical studies. In Phase I multiple-dose studies, there were no dose-related clinical adverse reactions observed with sitagliptin with doses of up to 600 mg per day for periods of up to 10 days and 400 mg per day for periods of up to 28 days.

A large overdose of metformin (or co-existing risks of lactic acidosis) may lead to lactic acidosis which is a medical emergency and must be treated in hospital. The most effective method to remove lactate and metformin is haemodialysis.

In clinical studies, approximately 13.5 % of the dose was removed over a 3- to 4-hour haemodialysis session. Prolonged haemodialysis may be considered if clinically appropriate. It is not known if sitagliptin is dialysable by peritoneal dialysis.

In the event of an overdose, it is reasonable to employ the usual supportive measures, e.g., remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring (including obtaining an electrocardiogram), and institute supportive therapy if required.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Drugs used in diabetes, Combinations of oral blood glucose lowering drugs, ATC code: A10BD07.

Sitagliptin/Metformin hydrochloride Krka combines two antihyperglycaemic medicinal products with complementary mechanisms of action to improve glycaemic control in patients with type 2 diabetes: sitagliptin, a dipeptidyl peptidase 4 (DPP-4) inhibitor, and metformin hydrochloride, a member of the biguanide class.

#### *Sitagliptin*

##### Mechanism of action

Sitagliptin phosphate is an orally-active, potent, and highly selective inhibitor of the dipeptidyl peptidase 4 (DPP-4) enzyme for the treatment of type 2 diabetes. The DPP-4 inhibitors are a class of agents that act as incretin enhancers. By inhibiting the DPP-4 enzyme, sitagliptin increases the levels of two known active incretin hormones, glucagon-like peptide-1 (GLP-1) and glucose-dependent insulinotropic polypeptide (GIP). The incretins are part of an endogenous system involved in the physiologic regulation of glucose homeostasis. When blood glucose concentrations are normal or elevated, GLP-1 and GIP increase insulin synthesis and release from pancreatic beta cells. GLP-1 also lowers glucagon secretion from pancreatic alpha cells, leading to reduced hepatic glucose production.

When blood glucose levels are low, insulin release is not enhanced and glucagon secretion is not suppressed. Sitagliptin is a potent and highly selective inhibitor of the enzyme DPP-4 and does not inhibit the closely-related enzymes DPP-8 or DPP-9 at therapeutic concentrations. Sitagliptin differs in chemical structure and pharmacological action from GLP-1 analogues, insulin, sulphonylureas or meglitinides, biguanides, peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) agonists, alpha-glucosidase inhibitors, and amylin analogues.

In a two-day study in healthy subjects, sitagliptin alone increased active GLP-1 concentrations, whereas metformin alone increased active and total GLP-1 concentrations to similar extents.

Co-administration of sitagliptin and metformin had an additive effect on active GLP-1 concentrations.

Sitagliptin, but not metformin, increased active GIP concentrations.

##### Clinical efficacy and safety

Overall, sitagliptin improved glycaemic control when used as monotherapy or in combination treatment in adult patients with type 2 diabetes.

In clinical studies, sitagliptin as monotherapy improved glycaemic control with significant reductions in haemoglobin A<sub>1c</sub> (HbA<sub>1c</sub>) and fasting and postprandial glucose. Reduction in fasting plasma glucose (FPG) was observed at 3 weeks, the first time point at which FPG was measured. The observed incidence of hypoglycaemia in patients treated with sitagliptin was similar to placebo. Body weight did not increase from baseline with sitagliptin therapy. Improvements in surrogate markers of beta cell function, including HOMA- $\beta$  (Homeostasis Model Assessment- $\beta$ ), proinsulin to insulin ratio, and measures of beta cell responsiveness from the frequently-sampled meal tolerance test were observed.

##### Studies of sitagliptin in combination with metformin

In a 24-week, placebo-controlled clinical study to evaluate the efficacy and safety of the addition of sitagliptin 100 mg once daily to ongoing metformin, sitagliptin provided significant improvements in glycaemic parameters compared with placebo. Change from baseline in body weight was similar for patients treated with sitagliptin relative to placebo. In this study there was a similar incidence of hypoglycaemia reported for patients treated with sitagliptin or placebo.

In a 24-week placebo-controlled factorial study of initial therapy, sitagliptin 50 mg twice daily in combination with metformin (500 mg or 1,000 mg twice daily) provided significant improvements in glycaemic parameters compared with either monotherapy. The decrease in body weight with the combination of sitagliptin and metformin was similar to that observed with metformin alone or placebo; there was no change from baseline for patients on sitagliptin alone. The incidence of hypoglycaemia was similar across treatment groups.

##### Study of sitagliptin in combination with metformin and a sulphonylurea

A 24-week placebo-controlled study was designed to evaluate the efficacy and safety of sitagliptin (100 mg once daily) added to glimepiride (alone or in combination with metformin). The addition of sitagliptin to glimepiride and metformin provided significant improvements in glycaemic parameters. Patients treated with sitagliptin had a modest increase in body weight (+1.1 kg) compared to those given placebo.

#### Study of sitagliptin in combination with metformin and a PPAR $\gamma$ agonist

A 26-week placebo-controlled study was designed to evaluate the efficacy and safety of sitagliptin (100 mg once daily) added to the combination of pioglitazone and metformin. The addition of sitagliptin to pioglitazone and metformin provided significant improvements in glycaemic parameters. Change from baseline in body weight was similar for patients treated with sitagliptin relative to placebo. The incidence of hypoglycaemia was also similar in patients treated with sitagliptin or placebo.

#### Study of sitagliptin in combination with metformin and insulin

A 24-week placebo-controlled study was designed to evaluate the efficacy and safety of sitagliptin (100 mg once daily) added to insulin (at a stable dose for at least 10 weeks) with or without metformin (at least 1,500 mg). In patients taking pre-mixed insulin, the mean daily dose was 70.9 U/day. In patients taking non-pre-mixed (intermediate/long-acting) insulin, the mean daily dose was 44.3 U/day. Data from the 73 % of patients who were taking metformin are presented in Table 2. The addition of sitagliptin to insulin provided significant improvements in glycaemic parameters. There was no meaningful change from baseline in body weight in either group.

**Table 2: HbA<sub>1c</sub> results in placebo-controlled combination therapy studies of sitagliptin and metformin\***

Study	Mean baseline HbA <sub>1c</sub> (%)	Mean change from baseline HbA <sub>1c</sub> (%)	Placebo-corrected mean change in HbA <sub>1c</sub> (%) (95 % CI)
Sitagliptin 100 mg once daily added to ongoing metformin therapy <sup>  </sup> (N=453)	8.0	-0.7 <sup>†</sup>	-0.7 <sup>†,‡</sup> (-0.8, -0.5)
Sitagliptin 100 mg once daily added to ongoing glimepiride + metformin therapy <sup>  </sup> (N=115)	8.3	-0.6 <sup>†</sup>	-0.9 <sup>†,‡</sup> (-1.1, -0.7)
Sitagliptin 100 mg once daily added to ongoing pioglitazone + metformin therapy <sup>¶</sup> (N=152)	8.8	-1.2 <sup>†</sup>	-0.7 <sup>†,‡</sup> (-1.0, -0.5)
Sitagliptin 100 mg once daily added to ongoing insulin + metformin therapy <sup>  </sup> (N=223)	8.7	-0.7 <sup>§</sup>	-0.5 <sup>§,‡</sup> (-0.7, -0.4)
Initial Therapy (twice daily) <sup>  </sup> : Sitagliptin 50 mg + metformin 500 mg (N=183)	8.8	-1.4 <sup>†</sup>	-1.6 <sup>†,‡</sup> (-1.8, -1.3)
Initial Therapy (twice daily) <sup>  </sup> : Sitagliptin 50 mg + metformin 1,000 mg (N=178)	8.8	-1.9 <sup>†</sup>	-2.1 <sup>†,‡</sup> (-2.3, -1.8)

\*All Patients Treated Population (an intention-to-treat analysis).

<sup>†</sup> Least squares means adjusted for prior antihyperglycaemic therapy status and baseline value.

<sup>‡</sup> p < 0.001 compared to placebo or placebo + combination treatment.

<sup>||</sup> HbA<sub>1c</sub> (%) at week 24.

<sup>¶</sup> HbA<sub>1c</sub> (%) at week 26.

<sup>§</sup> Least squares mean adjusted for insulin use at Visit 1 (pre-mixed vs. non-pre-mixed [intermediate- or long-acting]), and baseline value.

In a 52-week study, comparing the efficacy and safety of the addition of sitagliptin 100 mg once daily or glipizide (a sulphonylurea) in patients with inadequate glycaemic control on metformin monotherapy, sitagliptin was similar to glipizide in reducing HbA<sub>1c</sub> (-0.7 % mean change from baselines at week 52, with baseline HbA<sub>1c</sub> of approximately 7.5 % in both groups). The mean glipizide dose used in the comparator group was 10 mg per day with approximately 40 % of patients requiring a glipizide dose of ≤ 5 mg/day throughout the study. However, more patients in the sitagliptin group discontinued due to lack of efficacy than in the glipizide group. Patients treated with sitagliptin exhibited a significant mean decrease from baseline in body weight (-1.5 kg) compared to a significant weight gain in patients administered glipizide (+1.1 kg). In this study, the proinsulin to insulin ratio, a marker of efficiency of insulin synthesis and release, improved with sitagliptin and deteriorated with glipizide treatment. The incidence of hypoglycaemia in the sitagliptin group (4.9 %) was significantly lower than that in the glipizide group (32.0 %).

A 24-week placebo-controlled study involving 660 patients was designed to evaluate the insulin-sparing efficacy and safety of sitagliptin (100 mg once daily) added to insulin glargine with or without metformin (at least 1,500 mg) during intensification of insulin therapy. Among patients taking metformin, baseline HbA<sub>1c</sub> was 8.70 % and baseline insulin dose was 37 IU/day. Patients were instructed to titrate their insulin glargine dose based on fingerstick fasting glucose values. Among patients taking metformin, at Week 24, the increase in daily insulin dose was 19 IU/day in patients treated with sitagliptin and 24 IU/day in patients treated with placebo. The reduction in HbA<sub>1c</sub> for patients treated with sitagliptin, metformin, and insulin was -1.35 % compared to -0.90 % for patients treated with placebo, metformin, and insulin, a difference of -0.45 % [95 % CI: -0.62, -0.29]. The incidence of hypoglycaemia was 24.9 % for patients treated with sitagliptin, metformin, and insulin and 37.8 % for patients treated with placebo, metformin, and insulin. The difference was mainly due to a higher percentage of patients in the placebo group experiencing 3 or more episodes of hypoglycaemia (9.1 vs. 19.8 %). There was no difference in the incidence of severe hypoglycaemia.

### *Metformin*

#### Mechanism of action

Metformin is a biguanide with antihyperglycaemic effects, lowering both basal and postprandial plasma glucose. It does not stimulate insulin secretion and therefore does not produce hypoglycaemia.

Metformin may act via three mechanisms:

- by reduction of hepatic glucose production by inhibiting gluconeogenesis and glycogenolysis
- in muscle, by modestly increasing insulin sensitivity, improving peripheral glucose uptake and utilisation
- by delaying intestinal glucose absorption.

Metformin stimulates intracellular glycogen synthesis by acting on glycogen synthase. Metformin increases the transport capacity of specific types of membrane glucose transporters (GLUT-1 and GLUT-4).

#### Clinical efficacy and safety

In humans, independently of its action on glycaemia, metformin has favourable effects on lipid metabolism. This has been shown at therapeutic doses in controlled, medium-term or long-term clinical studies: metformin reduces total cholesterol, LDLc and triglyceride levels.

The prospective randomised (UKPDS) study has established the long-term benefit of intensive blood glucose control in type 2 diabetes. Analysis of the results for overweight patients treated with metformin after failure of diet alone showed:

- a significant reduction of the absolute risk of any diabetes-related complication in the metformin group (29.8 events/1,000 patient-years) versus diet alone (43.3 events/1,000 patient-years),  $p=0.0023$ , and versus the combined sulphonylurea and insulin monotherapy groups (40.1 events/1,000 patient-years),  $p=0.0034$
- a significant reduction of the absolute risk of any diabetes-related mortality: metformin 7.5 events/1,000 patient-years, diet alone 12.7 events/1,000 patient-years,  $p=0.017$
- a significant reduction of the absolute risk of overall mortality: metformin 13.5 events/1,000 patient-years versus diet alone 20.6 events/1,000 patient-years, ( $p=0.011$ ), and versus the combined sulphonylurea and insulin monotherapy groups 18.9 events/1,000 patient-years ( $p=0.021$ )
- a significant reduction in the absolute risk of myocardial infarction: metformin 11 events/1,000 patient-years, diet alone 18 events/1,000 patient-years, ( $p=0.01$ ).

The TECOS was a randomised study in 14,671 patients in the intention-to-treat population with an HbA<sub>1c</sub> of  $\geq 6.5$  to  $8.0\%$  with established CV disease who received sitagliptin (7,332) 100 mg daily (or 50 mg daily if the baseline eGFR was  $\geq 30$  and  $< 50$  mL/min/1.73 m<sup>2</sup>) or placebo (7,339) added to usual care targeting regional standards for HbA<sub>1c</sub> and CV risk factors. Patients with an eGFR  $< 30$  mL/min/1.73 m<sup>2</sup> were not to be enrolled in the study. The study population included 2,004 patients  $\geq 75$  years of age and 3,324 patients with renal impairment (eGFR  $< 60$  mL/min/1.73 m<sup>2</sup>).

Over the course of the study, the overall estimated mean (SD) difference in HbA<sub>1c</sub> between the sitagliptin and placebo groups was  $0.29\%$  (0.01), 95 % CI (-0.32, -0.27);  $p < 0.001$ .

The primary cardiovascular endpoint was a composite of the first occurrence of cardiovascular death, nonfatal myocardial infarction, nonfatal stroke, or hospitalization for unstable angina. Secondary cardiovascular endpoints included the first occurrence of cardiovascular death, nonfatal myocardial infarction, or nonfatal stroke; first occurrence of the individual components of the primary composite; all-cause mortality; and hospital admissions for congestive heart failure.

After a median follow up of 3 years, sitagliptin, when added to usual care, did not increase the risk of major adverse cardiovascular events or the risk of hospitalization for heart failure compared to usual care without sitagliptin in patients with type 2 diabetes (Table 3).

**Table 3: Rates of Composite Cardiovascular Outcomes and Key Secondary Outcomes**

	Sitagliptin 100 mg		Placebo		Hazard Ratio (95 % CI)	p-value <sup>†</sup>
	N (%)	Incidence rate per 100 patient years <sup>*</sup>	N (%)	Incidence rate per 100 patient years <sup>*</sup>		
<b>Analysis in the Intention-to-Treat Population</b>						
<b>Number of patients</b>	<b>7,332</b>		<b>7,339</b>			
<b>Primary Composite Endpoint</b> (Cardiovascular death, non-fatal myocardial infarction, non-fatal stroke, or hospitalization for unstable angina)	839 (11.4)	4.1	851 (11.6)	4.2	0.98 (0.89–1.08)	<0.001
<b>Secondary Composite Endpoint</b> (Cardiovascular death, non-fatal myocardial infarction, or nonfatal stroke)	745 (10.2)	3.6	746 (10.2)	3.6	0.99 (0.89–1.10)	<0.001
<b>Secondary Outcome</b>						
Cardiovascular death	380 (5.2)	1.7	366 (5.0)	1.7	1.03 (0.89–1.19)	0.711
All myocardial infarction (fatal and non-fatal)	300 (4.1)	1.4	316 (4.3)	1.5	0.95 (0.81–1.11)	0.487
All stroke (fatal and non-fatal)	178 (2.4)	0.8	183 (2.5)	0.9	0.97 (0.79–1.19)	0.760
Hospitalization for unstable angina	116 (1.6)	0.5	129 (1.8)	0.6	0.90 (0.70–1.16)	0.419
Death from any cause	547 (7.5)	2.5	537 (7.3)	2.5	1.01 (0.90–1.14)	0.875
Hospitalization for heart failure <sup>‡</sup>	228 (3.1)	1.1	229 (3.1)	1.1	1.00 (0.83–1.20)	0.983

\* Incidence rate per 100 patient-years is calculated as  $100 \times$  (total number of patients with  $\geq 1$  event during eligible exposure period per total patient-years of follow-up).

<sup>†</sup> Based on a Cox model stratified by region. For composite endpoints, the p-values correspond to a test of non-inferiority

seeking to show that the hazard ratio is less than 1.3. For all other endpoints, the p-values correspond to a test of differences in hazard rates.

‡ The analysis of hospitalization for heart failure was adjusted for a history of heart failure at baseline.

### Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with the reference medicinal product containing sitagliptin and metformin hydrochloride in all subsets of the paediatric population in type 2 diabetes mellitus (see section 4.2 for information on paediatric use).

The safety and efficacy of the addition of sitagliptin in paediatric patients aged 10 to 17 years with type 2 diabetes and inadequate glycaemic control on metformin with or without insulin was assessed in two studies over 54 weeks. The addition of sitagliptin (administered as sitagliptin + metformin or sitagliptin + metformin extended release (XR)) was compared to the addition of placebo to metformin or metformin XR.

While superiority of HbA1c reduction was demonstrated for sitagliptin + metformin / sitagliptin + metformin XR over metformin at Week 20 in the pooled analysis of these two studies, results from the individual studies were inconsistent. Furthermore, greater efficacy for sitagliptin + metformin / sitagliptin + metformin XR compared to metformin was not observed at Week 54. Therefore, sitagliptin/metformin should not be used in paediatric patients aged 10 to 17 years old because of insufficient efficacy (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

### Sitagliptin/metformin

A bioequivalence study in healthy subjects demonstrated that the sitagliptin/metformin hydrochloride combination tablets are bioequivalent to co-administration of sitagliptin phosphate and metformin hydrochloride as individual tablets.

The following statements reflect the pharmacokinetic properties of the individual active substances of Sitagliptin/Metformin hydrochloride Krka.

### *Sitagliptin*

#### Absorption

Following oral administration of a 100-mg dose to healthy subjects, sitagliptin was rapidly absorbed, with peak plasma concentrations (median  $T_{max}$ ) occurring 1 to 4 hours post-dose, mean plasma AUC of sitagliptin was 8.52  $\mu\text{M}\cdot\text{hr}$ ,  $C_{max}$  was 950 nM. The absolute bioavailability of sitagliptin is approximately 87 %. Since co-administration of a high-fat meal with sitagliptin had no effect on the pharmacokinetics, sitagliptin may be administered with or without food.

Plasma AUC of sitagliptin increased in a dose-proportional manner. Dose-proportionality was not established for  $C_{max}$  and  $C_{24hr}$  ( $C_{max}$  increased in a greater than dose-proportional manner and  $C_{24hr}$  increased in a less than dose-proportional manner).

#### Distribution

The mean volume of distribution at steady state following a single 100-mg intravenous dose of sitagliptin to healthy subjects is approximately 198 litres. The fraction of sitagliptin reversibly bound to plasma proteins is low (38 %).

#### Biotransformation

Sitagliptin is primarily eliminated unchanged in urine, and metabolism is a minor pathway. Approximately 79 % of sitagliptin is excreted unchanged in the urine.

Following a [ $^{14}\text{C}$ ]sitagliptin oral dose, approximately 16 % of the radioactivity was excreted as metabolites of sitagliptin. Six metabolites were detected at trace levels and are not expected to contribute to the plasma DPP-4 inhibitory activity of sitagliptin. *In vitro* studies indicated that the primary enzyme responsible for the limited metabolism of sitagliptin was CYP3A4, with contribution from CYP2C8.

*In vitro* data showed that sitagliptin is not an inhibitor of CYP isoenzymes CYP3A4, 2C8, 2C9, 2D6, 1A2, 2C19 or 2B6, and is not an inducer of CYP3A4 and CYP1A2.

#### Elimination

Following administration of an oral [ $^{14}\text{C}$ ]sitagliptin dose to healthy subjects, approximately 100 % of the administered radioactivity was eliminated in faeces (13 %) or urine (87 %) within one week of dosing. The apparent terminal  $t_{1/2}$  following a

100-mg oral dose of sitagliptin was approximately 12.4 hours. Sitagliptin accumulates only minimally with multiple doses. The renal clearance was approximately 350 mL/min.

Elimination of sitagliptin occurs primarily via renal excretion and involves active tubular secretion. Sitagliptin is a substrate for human organic anion transporter-3 (hOAT-3), which may be involved in the renal elimination of sitagliptin. The clinical relevance of hOAT-3 in sitagliptin transport has not been established. Sitagliptin is also a substrate of p-glycoprotein, which may also be involved in mediating the renal elimination of sitagliptin. However, ciclosporin, a p-glycoprotein inhibitor, did not reduce the renal clearance of sitagliptin. Sitagliptin is not a substrate for OCT2 or OAT1 or PEPT1/2 transporters. *In vitro*, sitagliptin did not inhibit OAT3 (IC<sub>50</sub>=160 µM) or p-glycoprotein (up to 250 µM) mediated transport at therapeutically relevant plasma concentrations. In a clinical study sitagliptin had a small effect on plasma digoxin concentrations indicating that sitagliptin may be a mild inhibitor of p-glycoprotein.

#### Characteristics in patients

The pharmacokinetics of sitagliptin were generally similar in healthy subjects and in patients with type 2 diabetes.

#### Renal impairment

A single-dose, open-label study was conducted to evaluate the pharmacokinetics of a reduced dose of sitagliptin (50 mg) in patients with varying degrees of chronic renal impairment compared to normal healthy control subjects. The study included patients with mild, moderate, and severe renal impairment, as well as patients with ESRD on haemodialysis. In addition, the effects of renal impairment on sitagliptin pharmacokinetics in patients with type 2 diabetes and mild, moderate, or severe renal impairment (including ESRD) were assessed using population pharmacokinetic analyses.

Compared to normal healthy control subjects, plasma AUC of sitagliptin was increased by approximately 1.2-fold and 1.6-fold in patients with mild renal impairment (GFR ≥ 60 to < 90 mL/min) and patients with moderate renal impairment (GFR ≥ 45 to < 60 mL/min), respectively.

Because increases of this magnitude are not clinically relevant, dose adjustment in these patients is not necessary.

Plasma AUC of sitagliptin was increased approximately 2-fold in patients with moderate renal impairment (GFR ≥ 30 to < 45 mL/min), and approximately 4-fold in patients with severe renal impairment (GFR < 30 mL/min), including patients with ESRD on haemodialysis. Sitagliptin was modestly removed by haemodialysis (13.5 % over a 3- to 4-hour haemodialysis session starting 4 hours post-dose).

#### Hepatic impairment

No dose adjustment for sitagliptin is necessary for patients with mild or moderate hepatic impairment (Child-Pugh score ≤ 9). There is no clinical experience in patients with severe hepatic impairment (Child-Pugh score > 9). However, because sitagliptin is primarily renally eliminated, severe hepatic impairment is not expected to affect the pharmacokinetics of sitagliptin.

#### Elderly

No dose adjustment is required based on age. Age did not have a clinically meaningful impact on the pharmacokinetics of sitagliptin based on a population pharmacokinetic analysis of Phase I and Phase II data. Elderly subjects (65 to 80 years) had approximately 19 % higher plasma concentrations of sitagliptin compared to younger subjects.

#### Paediatric population

The pharmacokinetics of sitagliptin (single dose of 50 mg, 100 mg or 200 mg) were investigated in paediatric patients (10 to 17 years of age) with type 2 diabetes. In this population, the dose adjusted AUC of sitagliptin in plasma was approximately 18 % lower compared to adult patients with type 2 diabetes for a 100 mg dose. No studies with sitagliptin have been performed in paediatric patients < 10 years of age.

#### Other patient characteristics

No dose adjustment is necessary based on gender, race, or body mass index (BMI). These characteristics had no clinically meaningful effect on the pharmacokinetics of sitagliptin based on a composite analysis of Phase I pharmacokinetic data and on a population pharmacokinetic analysis of Phase I and Phase II data.

#### Metformin

#### Absorption

After an oral dose of metformin, T<sub>max</sub> is reached in 2.5 h. Absolute bioavailability of a 500 mg metformin tablet is approximately 50-60 % in healthy subjects. After an oral dose, the non-absorbed fraction recovered in faeces was 20-30 %.

After oral administration, metformin absorption is saturable and incomplete. It is assumed that the pharmacokinetics of metformin absorption is non-linear. At the usual metformin doses and dosing schedules, steady state plasma concentrations are reached within 24-48 h and are generally less than 1 µg/mL. In controlled clinical studies, maximum metformin plasma levels ( $C_{max}$ ) did not exceed 5 µg/mL, even at maximum doses.

Food decreases the extent and slightly delays the absorption of metformin. Following administration of a dose of 850 mg, a 40% lower plasma peak concentration, a 25 % decrease in AUC and a 35 min prolongation of time to peak plasma concentration was observed. The clinical relevance of this decrease is unknown.

#### Distribution

Plasma protein binding is negligible. Metformin partitions into erythrocytes. The blood peak is lower than the plasma peak and appears at approximately the same time. The red blood cells most likely represent a secondary compartment of distribution. The mean  $V_d$  ranged between 63 – 276 L.

#### Biotransformation

Metformin is excreted unchanged in the urine. No metabolites have been identified in humans.

#### Elimination

Renal clearance of metformin is > 400 mL/min, indicating that metformin is eliminated by glomerular filtration and tubular secretion. Following an oral dose, the apparent terminal elimination half-life is approximately 6.5 h. When renal function is impaired, renal clearance is decreased in proportion to that of creatinine and thus the elimination half-life is prolonged, leading to increased levels of metformin in plasma.

### **5.3 Preclinical safety data**

No animal studies have been conducted with Sitagliptin/Metformin hydrochloride Krka.

In 16-week studies in which dogs were treated with either metformin alone or a combination of metformin and sitagliptin, no additional toxicity was observed from the combination. The NOEL in these studies was observed at exposures to sitagliptin of approximately 6 times the human exposure and to metformin of approximately 2.5 times the human exposure.

The following data are findings in studies performed with sitagliptin or metformin individually.

#### Sitagliptin

Renal and liver toxicity were observed in rodents at systemic exposure values 58 times the human exposure level, while the no-effect level was found at 19 times the human exposure level. Incisor teeth abnormalities were observed in rats at exposure levels 67 times the clinical exposure level; the no-effect level for this finding was 58-fold based on the 14-week rat study. The relevance of these findings for humans is unknown. Transient treatment-related physical signs, some of which suggest neural toxicity, such as open-mouth breathing, salivation, white foamy emesis, ataxia, trembling, decreased activity, and/or hunched posture were observed in dogs at exposure levels approximately 23 times the clinical exposure level. In addition, very slight to slight skeletal muscle degeneration was also observed histologically at doses resulting in systemic exposure levels of approximately 23 times the human exposure level. A no-effect level for these findings was found at an exposure 6-fold the clinical exposure level.

Sitagliptin has not been demonstrated to be genotoxic in preclinical studies. Sitagliptin was not carcinogenic in mice. In rats, there was an increased incidence of hepatic adenomas and carcinomas at systemic exposure levels 58 times the human exposure level. Since hepatotoxicity has been shown to correlate with induction of hepatic neoplasia in rats, this increased incidence of hepatic tumours in rats was likely secondary to chronic hepatic toxicity at this high dose. Because of the high safety margin (19-fold at this no-effect level), these neoplastic changes are not considered relevant for the situation in humans.

No treatment related effects on fertility were observed in male and female rats given sitagliptin prior to and throughout mating.

In a pre-/post-natal development study performed in rats sitagliptin showed no adverse reactions.

Reproductive toxicity studies showed a slight treatment-related increased incidence of foetal rib malformations (absent, hypoplastic and wavy ribs) in the offspring of rats at systemic exposure levels more than 29 times the human exposure levels. Maternal toxicity was seen in rabbits at more than 29 times the human exposure levels. Because of the high safety margins, these findings do not suggest a relevant risk for human reproduction. Sitagliptin is secreted in considerable amounts into the milk of lactating rats (milk/plasma ratio: 4:1).

### *Metformin*

Preclinical data for metformin reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

povidone  
microcrystalline cellulose  
mannitol  
sodium laurilsulfate  
magnesium stearate

#### Film coating

hypromellose  
titanium dioxide (E171)  
talc  
propylene glycol  
red ferric oxide (E172)

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

OPA/Alu/PVC//Alu blisters: 2 years  
PVC/PE/PVDC/PE/PVC//Alu blisters: 3 years

### **6.4 Special precautions for storage**

#### *OPA/Alu/PVC//Alu blisters:*

This medicinal product does not require any special temperature storage conditions.  
Store in the original package in order to protect from moisture.

#### *PVC/PE/PVDC/PE/PVC//Alu blisters:*

Do not store above 30°C. Store in the original package in order to protect from moisture.

### **6.5 Nature and contents of container**

Blister (OPA/Alu/PVC//Alu): 10, 14, 28, 30, 56, 60, 196, 200 film-coated tablets, in a box.  
Blister (OPA/Alu/PVC//Alu), calendar pack: 14, 28, 56, 196 film-coated tablets, in a box.  
Blister (PVC/PE/PVDC/PE/PVC//Alu): 10, 14, 28, 30, 56, 60, 196, 200 film-coated tablets, in a box.  
Blister (PVC/PE/PVDC/PE/PVC//Alu), calendar pack: 14, 28, 56, 196 film-coated tablets, in a box.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

No special requirements for disposal.

## **7 MARKETING AUTHORISATION HOLDER**

KRKA, d.d., Novo mesto  
Šmarješka cesta 6  
8501 Novo mesto

Slovenia

**8 MARKETING AUTHORISATION NUMBER**

PA1347/100/002

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 19<sup>th</sup> February 2021

Date of last renewal: 19<sup>th</sup> November 2025

**10 DATE OF REVISION OF THE TEXT**

September 2025