

Package leaflet: Information for the user

Enalapril maleate Accord 5 mg tablets
Enalapril maleate Accord 10 mg tablets
Enalapril maleate Accord 20 mg tablets
enalapril maleate

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.
- The full name of this medicine is Enalapril maleate Accord 5 mg, 10 mg, 20 mg tablets but within this leaflet it will be referred to as Enalapril maleate Accord tablets.

What is in this leaflet

1. What Enalapril maleate Accord tablets are and what they are used for
2. What you need to know before you take Enalapril maleate Accord tablets
3. How to take Enalapril maleate Accord tablets
4. Possible side effects
5. How to store Enalapril maleate Accord tablets
6. Contents of the pack and other information

1. What Enalapril maleate Accord tablets are and what they are used for

Enalapril maleate Accord tablets contain an active substance called Enalapril maleate. These belong to the group of medicines called ACE inhibitors (angiotensin converting enzyme inhibitors).

Enalapril maleate Accord tablets are used to:

- treat high blood pressure (hypertension)
- treat heart failure (weakening of heart function). It can lower the need to go to hospital and can help some patients live longer
- prevent the signs of heart failure. The signs include shortness of breath, tiredness after light physical activity such as walking, or swelling of the ankles and feet.

This medicine works by widening your blood vessels. This lowers your blood pressure. The medicine usually starts to work within an hour and the effect lasts for at least 24 hours. Some people will require several weeks of treatment until the best effect on your blood pressure is seen.

2. What you need to know before you take Enalapril maleate Accord tablets

Do not take Enalapril maleate Accord tablets if you:

- are allergic to Enalapril maleate Accord or any of the other ingredients of this medicine (listed in section 6)
- have ever had an allergic reaction to a type of medicine similar to this medicine called an ACE inhibitor
- have ever had swelling of your face, lips, mouth, tongue or throat which caused difficulty in swallowing or breathing (angioedema) when the reason why was not known or it was inherited
- have diabetes or impaired kidney function and you are treated with a blood pressure lowering medicine containing aliskiren

- are more than 3 months pregnant (It is also better to avoid Enalapril maleate Accord tablets in early pregnancy see Pregnancy section.)
- have taken or are currently taking sacubitril/valsartan, a medicine used to treat a type of long-term (chronic) heart failure in adults, as the risk of angioedema (rapid swelling under the skin in an area such as the throat) is increased.

Do not take this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking this medicine.

Warnings and precautions

Talk to your doctor or pharmacist before taking Enalapril maleate Accord tablets if you:

- have a heart problem
- have a condition involving the blood vessels in the brain
- have a blood problem such as low or lack of white blood cells (neutropenia/agranulocytosis), low blood platelet count (thrombocytopenia) or a decreased number of red blood cells (anaemia)
- have a liver problem
- have a kidney problem (including kidney transplantation), are on a salt-restricted diet, are taking potassium supplements, potassium-sparing agents, potassium-containing salt substitutes, or other drugs that may increase potassium in your blood (e.g., heparin [a medicine used to prevent blood clots], trimethoprim-containing products such as cotrimoxazole [medicines used to treat infections]). These may lead to higher levels of potassium in your blood which can be serious. Your doctor may need to adjust your dose of Enalapril maleate Accord tablets or monitor your blood level of potassium. See also information under the heading “Other medicines and Enalapril maleate Accord tablets.
- are having dialysis
- have been very sick (excessive vomiting) or had bad diarrhoea recently
- have diabetes. You should monitor your blood for low blood glucose levels, especially during the first month of treatment. The level of potassium in your blood can also be higher
- have ever had an allergic reaction with swelling of the face, lips, tongue or throat with difficulty in swallowing or breathing. You should be aware that black patients are at increased risk of these types of reactions to ACE inhibitors
- have low blood pressure (you may notice this as faintness or dizziness, especially when standing)
- have collagen vascular disease (e.g., lupus erythematosus, rheumatoid arthritis or scleroderma), are on therapy that suppresses your immune system, are taking the drugs allopurinol or procainamide, or any combinations of these
- are taking any of the following medicines, the risk of angioedema may be increased:
 - Racecadotril, a medicine used to treat diarrhoea.
 - Medicines used to prevent organ transplant rejection and for cancer (e.g., temsirolimus, sirolimus, everolimus).
 - Vildagliptin, a medicine used to treat diabetes
- are taking any of the following medicines used to treat high blood pressure:
 - an angiotensin II receptor blocker (ARB) (also known as sartans - for example valsartan, telmisartan, irbesartan, etc.), in particular if you have diabetes-related kidney problems
 - aliskiren.

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (e.g., potassium) in your blood at regular intervals.

See also information under the heading “Do not take Enalapril maleate Accord if you:”

You must tell your doctor if you think you are (or might become) pregnant. This medicine is not recommended in early pregnancy and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see Pregnancy section).

You should be aware that this medicine lowers the blood pressure in black patients less effectively than in non-black patients.

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking this medicine.

If you are about to have a procedure

If you are about to have any of the following, tell your doctor that you are taking Enalapril maleate Accord tablets:

- any surgery or receive anaesthetics (even at the dentist)
- a treatment to remove cholesterol from your blood called ‘LDL apheresis’
- a desensitisation treatment, to lower the effect of an allergy to bee or wasp stings.

If any of the above applies to you, talk to your doctor or dentist before the procedure.

Other medicines and Enalapril maleate Accord tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes herbal medicines. This is because Enalapril maleate Accord tablets can affect the way some medicines work. Also, some other medicines can affect the way Enalapril maleate Accord tablets work. Your doctor may need to change your dose and/or to take other precautions.

In particular tell your doctor or pharmacist if you are taking any of the following medicines:

- an angiotensin II receptor blocker (ARB) or aliskiren (see also information under the headings “Do not take Enalapril maleate Accord tablets if you:” and “Warnings and precautions”)
- other medicines to lower blood pressure, such as beta-blockers or water tablets (diuretics)
- potassium supplements (including salt substitutes), potassium-sparing diuretics and other medicines that can increase the amount of potassium in your blood (trimethoprim and co-trimoxazole for infections caused by bacteria; ciclosporin, an immunosuppressant medicine used to prevent organ transplant rejection; and heparin, a medicine used to thin blood to prevent clots). See also information under the heading “Warnings and precautions”.
- medicines for diabetes (including oral antidiabetic medicines and insulin)
- lithium (a medicine used to treat a certain kind of depression)
- medicines for depression called ‘tricyclic antidepressants’
- medicines for mental problems called ‘antipsychotic’
- certain cough and cold medicines and weight reducing medicines which contain something called a ‘sympathomimetic agent’
- certain pain or arthritis medicines including gold therapy
- an mTOR inhibitor (e.g., temsirolimus, sirolimus, everolimus; medicines used to treat certain types of cancer or to prevent the body’s immune system from rejecting a transplanted organ). See also information under the heading “Warnings and precautions”
- a medicine containing a neprilysin inhibitor such as sacubitril (available as fixed-dose combination with valsartan), racecadotril or vildagliptin. The risk of angioedema (swelling of the face, lips, tongue or throat with difficulty in swallowing or breathing) may be increased. See also information under the headings “Do not take Enalapril maleate Accord tablets if you:” and “Warnings and precautions”
- non-steroidal anti-inflammatory drugs, including COX-2-inhibitors (medicines that reduce inflammation, and can be used to help relieve pain)
- aspirin (acetylsalicylic acid)
- medicines used to dissolve blood clots (thrombolytics)
- alcohol

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking Enalapril maleate Accord tablets.

Enalapril maleate Accord tablets with food and drink

Enalapril maleate Accord tablets can be taken with or without food. Most people take Enalapril maleate Accord tablets with a drink of water.

Pregnancy and breast-feeding

Pregnancy

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will normally advise you to stop taking Enalapril maleate Accord tablets before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Enalapril maleate Accord tablets. This medicine is not recommended in early pregnancy and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

Breast-feeding

Tell your doctor if you are breast-feeding or about to start breast-feeding. Breast-feeding newborn babies (first few weeks after birth), and especially premature babies, is not recommended whilst taking this medicine. In the case of an older baby your doctor should advise you on the benefits and risks of taking this medicine whilst breast-feeding, compared to other treatments.

Driving and using machines

You may feel dizzy or sleepy while taking this medicine. If this happens, do not drive or use any tools or machines.

Enalapril maleate Accord tablets contain lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

Enalapril maleate Accord tablets contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially “sodium-free”.

3. How to take Enalapril maleate Accord tablets

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

- It is very important to continue taking this medicine for as long as your doctor prescribes it.
- Do not take more tablets than prescribed.
- The score line is only there to help you break the tablet if you have difficulty swallowing it whole.

High Blood Pressure

- The usual starting dose ranges from 5 to 20 mg taken once a day.
- Some patients may need a lower starting dose.
- The usual long term dose is 20 mg taken once a day.
- The maximal long term dose is 40 mg taken once a day.

Heart Failure

- The usual starting dose is 2.5 mg taken once a day.
- Your doctor will raise this amount step by step until the dose that is right for you has been achieved.

- The usual long term dose is 20 mg each day, taken in one or two doses.
- The maximal long term dose is 40 mg each day, divided in two doses.

Patients with kidney problems

Your dose of medicine will be changed depending on how well your kidneys are working:

- moderate kidney problems – 5 mg to 10 mg each day
- severe kidney problems – 2.5 mg each day
- if you are having dialysis – 2.5 mg each day. On days you are not having dialysis, your dose may be changed depending on how low your blood pressure is.

Elderly patients

Your dose will be decided by your doctor and will be based on how well your kidneys are working.

Use in children

Experience in the use of Enalapril maleate Accord tablets in children with high blood pressure is limited. If the child can swallow tablets, the dose will be worked out using the child's weight and blood pressure. The usual starting doses are:

- between 20 kg and 50 kg – 2.5 mg each day
- more than 50 kg – 5 mg each day.

The dose can be changed according to the needs of the child:

- a maximum of 20 mg daily can be used in children who are between 20 kg and 50 kg
- a maximum of 40 mg daily can be used in children who are more than 50 kg.

This medicine is not recommended in newborn babies (first few weeks after birth) and in children with kidney problems.

If you take more Enalapril maleate Accord tablets than you should

If you take more Enalapril maleate Accord tablets than you should, talk to your doctor or go to a hospital straight away. Take the medicine pack with you. The following effects may happen: feeling of light-headedness or dizziness. This is due to a sudden or excessive drop in blood pressure.

If you forget to take Enalapril maleate Accord tablets

- If you forget to take a tablet, skip the missed dose.
- Take the next dose as usual.
- Do not take a double dose to make up for a forgotten dose.

If you stop taking Enalapril maleate Accord tablets

Do not stop taking your medicine unless your doctor tells you to.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. The following side effects may happen with this medicine:

Stop taking Enalapril maleate Accord tablets and talk to a doctor straight away, if you notice any of the following:

- swelling of your face, lips, tongue or throat which may cause difficulty in breathing or swallowing
- swelling of your hands, feet or ankles
- if you develop a raised red skin rash (hives).

You should be aware that black patients are at increased risk of these types of reactions. If any of the above happen, stop taking Enalapril maleate Accord tablets and talk to a doctor straight away.

When you start taking this medicine you may feel faint or dizzy. If this happens, it will help to lie down. This is caused by your blood pressure lowering. It should improve as you continue to take the medicine. If you are worried, please talk to your doctor.

Other side effects include:

Very Common (may affect more than 1 in 10 people)

- feeling dizzy, weak or sick
- blurred vision
- cough.

Common (may affect up to 1 in 10 people)

- light-headedness due to low blood pressure, changes in heart rhythm, fast heartbeat, angina or chest pain
- headache, depression, fainting (syncope), change in sense of taste
- shortness of breath
- diarrhoea, abdominal pain
- tiredness (fatigue)
- rash, allergic reactions with swelling of the face, lips, tongue or throat with difficulty in swallowing or breathing
- high levels of potassium in the blood, increased levels of creatinine in your blood (both are usually detected by a test).

Uncommon (may affect up to 1 in 100 people)

- flushing
- sudden fall in blood pressure
- fast or uneven heart beats (palpitations)
- heart attack (possibly due to very low blood pressure in certain high-risk patients, including those with blood flow problems of the heart or brain)
- stroke (possibly due to very low blood pressure in high-risk patients)
- anaemia (including aplastic and haemolytic)
- confusion, sleeplessness or sleepiness, nervousness
- feeling your skin prickling or being numb
- vertigo (spinning sensation)
- ringing in your ears (tinnitus)
- runny nose, sore throat or hoarseness
- asthma-associated tightness in chest
- slow movement of food through your intestine (ileus), inflammation of your pancreas
- being sick (vomiting), indigestion, constipation, anorexia
- irritated stomach (gastric irritations), dry mouth, ulcer
- muscle cramps
- impaired kidney function, kidney failure
- increased sweating
- itching or nettle rash
- hair loss
- generally feeling unwell (malaise), high temperature (fever)
- impotence
- high level of proteins in your urine (measured in a test)
- low level of blood sugar or sodium, high level of blood urea (all measured in a blood test)

Rare (may affect up to 1 in 1,000 people)

- ‘Raynaud’s phenomenon’ where your hands and feet may become very cold and white due to low blood flow
- changes in blood values such as a lower number of white and red blood cells, lower haemoglobin, lower number of blood platelets
- bone marrow depression
- swollen glands in neck, armpit or groin
- autoimmune diseases
- strange dreams or sleep problems
- accumulation of fluid or other substances in the lungs (as seen on X-rays)
- inflammation of your nose
- inflammation of the lungs causing difficulty breathing (pneumonia)
- inflammation of the cheeks, gums, tongue, lips, throat
- reduced amount of urine
- rash that looks like targets (erythema multiforme)
- ‘Stevens-Johnson syndrome’ and ‘toxic epidermal necrolysis’ (serious skin conditions where you have reddening and scaling of your skin, blistering or raw sores), exfoliative dermatitis/erythroderma (severe skin rash with flaking or peeling of the skin), pemphigus (small fluid-filled bumps on the skin)
- liver or gallbladder problems such as lower liver function, inflammation of your liver, jaundice (yellowing of the skin or eyes), high levels of liver enzymes or bilirubin (measured in a blood test)
- enlargement of breasts in males (gynaecomastia)

Very Rare (may affect up to 1 in 10,000 people)

- swelling in your intestine (intestinal angioedema)

Not known (frequency cannot be estimated from the available data)

- overproduction of antidiuretic hormone, which causes fluid retention, resulting in weakness, tiredness or confusion
- A symptom complex has been reported which may include some or all of the following: fever, inflammation of the blood vessels (serositis/vasculitis), muscle pain (myalgia/myositis), joint pain (arthralgia/arthritis). Rash, photosensitivity or other skin manifestations may occur.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via

HPRA Pharmacovigilance

Website: www.hpra.ie

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Enalapril maleate Accord tablets

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date, which is stated on the blister and carton after “EXP”. The expiry date refers to the last day of that month.
- This medicinal product does not require any special storage conditions.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist

how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Enalapril maleate Accord tablets contain

- The active substance is Enalapril maleate Accord (either 5 mg, 10 mg, or 20 mg).
- The other ingredients are microcelac 100 (contains: lactose monohydrate, microcrystalline cellulose), pregelatinized starch, croscarmellose sodium, sodium hydrogen carbonate (“See section 2”), magnesium stearate. The 10 mg/20 mg tablets also contains iron oxide red (E172).

What Enalapril maleate Accord tablets look like and contents of the pack

Enalapril maleate Accord 5 mg tablets

White to off-white, coloured, triangular shaped (6.50 mm diameter), biconvex uncoated tablets debossed with “IS2” on one side and **breakline** on other side.

Enalapril maleate Accord 10 mg tablets

Light Pink to Pink coloured, triangular shaped (6.50 mm diameter), biconvex uncoated tablets with mottled appearance, debossed with “IS3” on one side and **breakline** on other side.

Enalapril maleate Accord 20 mg tablets

Light Pink to Pink coloured, triangular shaped (8.80 mm diameter), biconvex uncoated tablets with mottled appearance, debossed with “IS5” on one side and **breakline** on other side.

Enalapril maleate Accord 5 mg tablets are available in aluminium-aluminium blister packs containing 20, 28, 30, 56, 60, 90, 100 tablets

Enalapril maleate Accord 10 mg tablets are available in aluminium-aluminium blister packs containing 20, 28, 30, 56, 60, 90, 100 tablets

Enalapril maleate Accord 20 mg tablets are available in aluminium-aluminium blister packs containing 14, 20, 28, 30, 60, 90, 100 tablets

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Accord Healthcare Ireland Limited
Euro House, Euro Business Park,
Little Island, Cork, T45 K857,
Ireland

Manufacturer

Accord Healthcare Polska Sp.
z.o.o., ul. Lutomiarska 50,
Pabianice, Poland

Accord Healthcare B.V.,
Winthontlaan 200, 3526 KV,
Netherlands

Accord Healthcare Limited,
Sage House, 319 Pinner Road,

North Harrow, Middlesex
HA1 4HF
United Kingdom

This medicine is authorized in the Member States of the European Economic Area (EEA) and in the United Kingdom (Northern Ireland) under the following names:

Name of member state	Name of medicinal product
Austria	Enalapril Accord 5mg Tabletten Enalapril Accord 10mg Tabletten Enalapril Accord 20mg Tabletten
Denmark	Enalapril Accord
Sweden	Enalapril Accord
Netherlands	Enalaprilmaleaat Accord 5mg tabletten Enalaprilmaleaat Accord 10mg tabletten Enalaprilmaleaat Accord 20mg tabletten
France	Enalapril Accord 5mg comprimé Enalapril Accord 20mg comprimé
Romania	Enalapril Accord 5mg comprimate Enalapril Accord 10mg comprimate Enalapril Accord 20mg comprimate
Italy	Enalapril Accord
Cyprus	Enalapril Accord 5mg Tablets Enalapril Accord 10mg Tablets
Portugal	Enalapril Accord
Malta	Enalapril Accord 5mg Tablets Enalapril Accord 20mg Tablets
Ireland	Enalapril maleate Accord 5mg Tablets Enalapril maleate Accord 10mg Tablets Enalapril maleate Accord 20mg Tablets
United Kingdom	Enalapril maleate 2.5mg Tablets Enalapril maleate 5mg Tablets Enalapril maleate 10mg Tablets Enalapril maleate 20mg Tablets

This leaflet was last revised in November 2024.