

Package leaflet: Information for the User
Lamotrigine SyriMed 5mg/ml Oral Suspension
lamotrigine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- ▶ Keep this leaflet. You may need to read it again.
- ▶ If you have any further questions, ask your doctor or pharmacist.
- ▶ This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- ▶ If you get any of the side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of your medicine is Lamotrigine SyriMed 5mg/ml Oral Suspension but it will be referred to as 'Lamotrigine' throughout this leaflet.

What is in this leaflet

1. What Lamotrigine is and what it is used for
2. What you need to know before you take Lamotrigine
3. How to take Lamotrigine
4. Possible side effects
5. How to store Lamotrigine
6. Contents of the pack and other information

1. What Lamotrigine is and what it is used for

Lamotrigine belongs to a group of medicines called *anti-epileptics*. It is used to treat two conditions - **epilepsy** and **bipolar disorder**.

Lamotrigine treats epilepsy by blocking the signals in the brain that trigger epileptic seizures (fits).

- ▶ For adults and children aged 13 years and over, Lamotrigine can be used on its own or with other medicines, to treat epilepsy. Lamotrigine can also be used with other medicines to treat the seizures that occur with a condition called Lennox-Gastaut syndrome.
- ▶ For children aged between 2 and 12 years, Lamotrigine can be used with other medicines, to treat those conditions. It can be used on its own to treat a type of epilepsy called typical absence seizures.

Lamotrigine also treats bipolar disorder.

People with bipolar disorder (sometimes called *manic depression*) have extreme mood swings, with periods of mania (excitement or euphoria) alternating with periods of depression (deep sadness or despair). For adults aged 18 years and over, Lamotrigine can be used on its own or with other medicines, to prevent the periods of depression that occur in bipolar disorder. It is not yet known how Lamotrigine works in the brain to have this effect.

2. What you need to know before you take Lamotrigine

Do not take Lamotrigine :

- ▶ **if you are allergic** (*hypersensitive*) to lamotrigine or any of the other ingredients of this medicine (listed in Section 6).

If this applies to you:

- ▶ **Tell your doctor**, and don't take Lamotrigine.

Warnings and precautions

Take special care with Lamotrigine

Talk to your doctor or pharmacist before taking Lamotrigine:

- ▶ **if you have any kidney problems**

- ▶ **if you have ever developed a rash** after taking lamotrigine or other medicines for bipolar disorder or epilepsy
- ▶ **if you experience a rash or sunburn after taking lamotrigine and having been exposed to sun or artificial light (e.g. solarium).** Your doctor will check your treatment and may advise you to avoid sunlight or protect yourself against the sun (e.g. use of a sunscreen and/or to wear protective clothing)
- ▶ **if you have ever developed meningitis after taking lamotrigine** (*read the description of these symptoms in Section 4 of this leaflet: Rare side effects*)
- ▶ **if you are already taking medicine that contains lamotrigine**
- ▶ **if you have a condition called Brugada syndrome, or other heart problems.** Brugada syndrome is a genetic disease that results in abnormal electrical activity within the heart. ECG abnormalities which may lead to arrhythmias (abnormal heart rhythm) can be triggered by lamotrigine.

If any of these applies to you:

- **Tell your doctor**, who may decide to lower the dose, or that Lamotrigine is not suitable for you.

Important information about potentially life-threatening reactions

A small number of people taking Lamotrigine get an allergic reaction or potentially life-threatening skin reaction, which may develop into more serious problems if they are not treated. These can include Stevens- Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) and Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS). You need to know the symptoms to look out for while you are taking Lamotrigine. This risk may be associated with a variant in genes in people from Asian origin (mainly Han Chinese and Thai). If you are of such origin and have been tested previously carrying this genetic variant (HLA-B* 1502), discuss this with your doctor before taking lamotrigine.

- ▶ **Read the description of these symptoms in Section 4 of this leaflet** under '*Potentially life-threatening reactions: get a doctor's help straight away*'.

Haemophagocytic lymphohistiocytosis (HLH)

There have been reports of a rare but very serious immune system reaction, in patients taking lamotrigine.

- ▶ **Contact your doctor or pharmacist immediately** if you experience any of the following symptoms while taking lamotrigine: fever, rash, neurological symptoms (e.g. shaking or tremor, confusional state, disturbances of brain function).

Thoughts of harming yourself or suicide

Anti-epileptic medicines are used to treat several conditions, including epilepsy and bipolar disorder. People with bipolar disorder can sometimes have thoughts of harming themselves or committing suicide. If you have bipolar disorder, you may be more likely to think like this:

- ▶ when you first start treatment
- ▶ if you have previously had thoughts about harming yourself or about suicide
- ▶ if you are under 25 years old.

If you have distressing thoughts or experiences, or if you notice that you feel worse or develop new symptoms while you're taking Lamotrigine:

- ▶ **See a doctor as soon as possible or go to the nearest hospital for help.**

You may find it helpful to tell a family member, caregiver or close friend that you can become depressed or have significant changes in mood, and ask them to read this leaflet. You might ask them to tell you if they are worried about your depression or other changes in your behaviour.

A small number of people being treated with anti-epileptics such as Lamotrigine have also had thoughts of harming or killing themselves. If at any time you have these thoughts, immediately contact your doctor.

If you're taking Lamotrigine for epilepsy

The seizures in some types of epilepsy may occasionally become worse or happen more often while you're taking Lamotrigine. Some patients may experience severe seizures, which may cause serious health problems. If your seizures happen more often, or if you experience a severe seizure while you're taking Lamotrigine:

- ▶ **See a doctor as soon as possible.**

Children and adolescents

Lamotrigine should not be given to people aged under 18 years to treat bipolar disorder.

Medicines to treat depression and other mental health problems increase the risk of suicidal thoughts and behaviour in children and adolescents aged under 18 years.

Other medicines and Lamotrigine

Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines including herbal medicines or other medicines bought without a prescription.

Your doctor needs to know if you are taking other medicines to treat epilepsy or mental health problems. This is to make sure you take the correct dose of Lamotrigine. These medicines include:

- ▶ **oxcarbazepine, felbamate, gabapentin, levetiracetam, pregabalin, topiramate or zonisamide**, used to treat **epilepsy**
- ▶ **lithium, olanzapine or aripiprazole** used to treat **mental health problems**
- ▶ **bupropion**, used to treat **mental health problems** or to **stop smoking**
- ▶ **paracetamol**, used to treat **pain** and **fever**.

Tell your doctor if you are taking any of these.

Some medicines interact with Lamotrigine or make it more likely that people will have side effects. These include:

- ▶ **valproate**, used to treat **epilepsy** and **mental health problems**
- ▶ **carbamazepine**, used to treat **epilepsy** and **mental health problems**
- ▶ **phenytoin, primidone or phenobarbitone**, used to treat **epilepsy**
- ▶ **risperidone**, used to treat **mental health problems**
- ▶ **rifampicin**, which is an **antibiotic**
- ▶ medicines used to treat **Human Immunodeficiency Virus (HIV) infection** (a combination of lopinavir and ritonavir or atazanavir and ritonavir)
- ▶ **hormonal contraceptives**, such as **the Pill** (*see below*).

Tell your doctor if you are taking any of these, or if you start or stop taking any.

Hormonal contraceptives (such as the Pill) can affect the way Lamotrigine works

Your doctor may recommend that you use a particular type of hormonal contraceptive, or another method of contraception, such as condoms, a cap or coil. If you are using a hormonal contraceptive like the Pill, your doctor may take samples of your blood to check the level of Lamotrigine. If you are using a hormonal contraceptive, or if you plan to start using one:

- ▶ **Talk to your doctor**, who will discuss suitable methods of contraception with you.

Lamotrigine can also affect the way hormonal contraceptives work, although it's unlikely to make them less effective. If you are using a hormonal contraceptive, and you notice any changes in your menstrual pattern, such as breakthrough bleeding or spotting between periods:

- ▶ **Tell your doctor**. These may be signs that Lamotrigine is affecting the way your contraceptive is working.

Pregnancy and breast-feeding

- ▶ **If you are pregnant, think you may be pregnant, or are planning to have a baby ask your doctor or pharmacist for advice before taking this medicine.**
 - **You should not stop treatment without discussing this with your doctor.** This is particularly important if you have epilepsy.

- Pregnancy may alter the effectiveness of Lamotrigine, so you may need blood tests and your dose of Lamotrigine may be adjusted.
- There may be a small increased risk of birth defects, including a cleft lip or cleft palate, if Lamotrigine is taken during the first 3 months of pregnancy.
- Your doctor may advise you to take extra **folic acid** if you're planning to become pregnant and while you're pregnant.

► **If you are breast-feeding or planning to breast-feed ask your doctor or pharmacist for advice before taking this medicine.** The active ingredient of Lamotrigine passes into breast milk and may affect your baby. Your doctor will discuss the risks and benefits of breast-feeding while you're taking Lamotrigine, and will check your baby from time to time, whether drowsiness, rash or poor weight gain occurs, if you decide to breast-feed. Inform your doctor if you observe any of these symptoms in your baby.

Driving and using machines

Lamotrigine can cause dizziness and double vision.

- **Don't drive or use machines unless you are sure you're not affected.**

If you have epilepsy, talk to your doctor about driving and using machines.

Lamotrigine contains:

Methyl parahydroxybenzoate (E218) and propyl parahydroxybenzoate (E216): May cause allergic reactions (possibly delayed).

Sodium: This medicine contains 0.676 mg sodium (main component of cooking/table salt) in each ml. This is equivalent to 0.03% of the recommended maximum daily dietary intake of sodium for an adult.

Propylene glycol (E1520): This medicine contains 0.940 mg propylene glycol in each ml.

Benzoic acid (E210): This medicine contains 0.0025 mg benzoic acid in each ml.

3. How to take Lamotrigine

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much Lamotrigine to take

It may take a while to find the best dose of Lamotrigine for you. The dose you take will depend on:

- your age
- whether you are taking Lamotrigine with other medicines
- whether you have any kidney or liver problems.

Your doctor will prescribe a low dose to start, and gradually increase the dose over a few weeks until you reach a dose that works for you (called the *effective dose*). **Never take more Lamotrigine than your doctor tells you to.**

Use in children and adolescents

The usual effective dose of Lamotrigine for adults and children aged 13 years or over is between 100 mg and 400 mg each day. For doses greater than 200mg, due to the volume that needs to be administered, other suitable formulations should be considered.

For children aged 2 to 12 years, the effective dose depends on their body weight - usually, it's between 1 mg and 15 mg (0.2 ml and 3 ml) for each kilogram of the child's weight, up to a maximum maintenance dose of 200mg (40ml) daily.

Lamotrigine is not recommended for children aged under 2 years.


Route and Method of administration:

- This medicinal product must be taken orally.
- Shake well before use for at least 10 seconds.


- Take your dose of Lamotrigine once or twice a day, as your doctor advises. It can be taken with or without food.
- Your doctor may also advise you to start or stop taking other medicines, depending on what condition you're being treated for and the way you respond to treatment.
- **Always take the full dose** that your doctor has prescribed. Never take only part of a dose.
- Use the measuring syringes provided in the pack to deliver the required dose. Lamotrigine is available in two presentations, 150ml pack with a 1ml and a 3ml syringe and 300ml pack with a 10ml syringe and a 20ml cup. If your dose is smaller than 3ml, pack of 150ml is recommended. If your dose is larger than 3ml, pack of 300ml is recommended.
- Your doctor or pharmacist may advise you to take a required dose using 20ml cup provided with 300ml pack. Use of syringe adaptor is not recommended while taking dose using 20ml cup.

Instructions for the use of syringe:

- a) Open the bottle: press the cap and turn it anticlockwise (figure 1).
- b) Separate the adaptor from the syringe (figure 2). Insert the adaptor into the bottle neck (figure 3). Ensure it is properly fixed. Take the syringe and put it in the adaptor opening (figure 4).



- c) Turn the bottle upside down. Fill the syringe with a small amount of suspension by pulling the piston down (figure 5A) and then push the piston up in order to remove any possible air bubbles (figure 5B). Pull the piston down to the graduation mark corresponding to the quantity in millilitres (ml) prescribed by your doctor (figure 5C).



- d) Turn the bottle the right way up (figure 6A). Remove the syringe from the adaptor (figure 6B).
- e) Empty the contents of the syringe into the mouth by pushing the piston to the bottom of the syringe (figure 7). The contents of the syringe should be emptied into the side cheek of the patient's mouth to avoid a choking hazard. Close the bottle with the plastic screw cap. Wash the syringe with water (figure 8).



If you take more Lamotrigine than you should

- **Contact a doctor or nearest hospital emergency department immediately.** If possible, show them the Lamotrigine packet.

If you take too much **Lamotrigine** you may be more likely to have serious side effects which may be fatal.

Someone who has taken too much Lamotrigine may have any of these symptoms:

- ▶ rapid, uncontrollable eye movements (*nystagmus*)
- ▶ clumsiness and lack of co-ordination, affecting their balance (*ataxia*)
- ▶ heart rhythm changes (detected usually on ECG)
- ▶ loss of consciousness, fits (convulsions) or coma.

If you forget to take a single dose of Lamotrigine

- **Don't take an extra dose to make up for your missed dose. Just take your next dose at the usual time.**

In case you forget to take multiple doses of Lamotrigine

- **Ask your doctor for advice on how to start taking it again.** It's important that you do this.

Don't stop taking Lamotrigine without advice

Lamotrigine must be taken for as long as your doctor recommends. Don't stop unless your doctor advises you to.

If you're taking Lamotrigine for epilepsy

To stop taking Lamotrigine, **it is important that the dose is reduced gradually**, over about 2 weeks. If you suddenly stop taking Lamotrigine, your epilepsy may come back or get worse.

If you're taking Lamotrigine for bipolar disorder

Lamotrigine may take some time to work, so you are unlikely to feel better straight away. If you stop taking Lamotrigine, your dose will not need to be reduced gradually. But you should still talk to your doctor first, if you want to stop taking Lamotrigine.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Potentially life-threatening reactions: get a doctor's help straight away

A small number of people taking Lamotrigine get an allergic reaction or potentially life-threatening skin reaction, which may develop into more serious problems if they are not treated.

These symptoms are more likely to happen during the first few months of treatment with Lamotrigine, especially if the starting dose is too high or if the dose is increased too quickly, or if Lamotrigine is taken with another medicine called *valproate*. Some of the symptoms are more common in children, so parents should be especially careful to watch out for them.

Symptoms of these reactions include:

- ▶ **skin rashes or redness**, which may develop into serious or sometimes life-threatening skin reactions including rash with target lesions (*Erythema multiforme*), widespread rash with blisters and peeling skin, particularly occurring around the mouth, nose, eyes and genitals (*Stevens-Johnson syndrome*), extensive peeling of the skin (more than 30% of the body surface - *toxic epidermal necrolysis*) or extended rashes with liver, blood and other body organs involvement (Drug Reaction with Eosinophilia and Systemic Symptoms which is also known as DRESS hypersensitivity syndrome)
- ▶ **ulcers in the mouth, throat, nose or genitals**
- ▶ **a sore mouth or red or swollen eyes** (*conjunctivitis*)
- ▶ **a high temperature** (*fever*), flu-like symptoms or drowsiness
- ▶ **swelling around your face, or swollen glands** in your neck, armpit or groin
- ▶ **unexpected bleeding or bruising**, or the fingers turning blue
- ▶ **a sore throat**, or more infections (such as colds) than usual
- ▶ increased levels of liver enzymes seen in blood tests
- ▶ an increase in a type of white blood cell (eosinophils)

- ▶ enlarged lymph nodes
- ▶ involvement of the organs of the body including liver and kidneys.

In many cases, these symptoms will be signs of less serious side effects. **But you must be aware that they are potentially life-threatening and can develop into more serious problems**, such as organ failure, if they are not treated. If you notice any of these symptoms:

- **Contact a doctor immediately.** Your doctor may decide to carry out tests on your liver, kidneys or blood, and may tell you to stop taking Lamotrigine. In case you have developed Stevens-Johnson syndrome or toxic epidermal necrolysis your doctor will tell you that you must never use lamotrigine again.

Haemophagocytic lymphohistiocytosis (HLH) (see section 2: What you need to know before you take Lamotrigine).

Very common side effects

These may affect **more than 1 in 10** people:

- ▶ headache
- ▶ skin rash.

Common side effects

These may affect **up to 1 in 10** people:

- ▶ aggression or irritability
- ▶ feeling sleepy or drowsy
- ▶ feeling dizzy
- ▶ shaking or tremors
- ▶ difficulty in sleeping (*insomnia*)
- ▶ feeling agitated
- ▶ diarrhoea
- ▶ dry mouth
- ▶ feeling sick (*nausea*) or being sick (*vomiting*)
- ▶ feeling tired
- ▶ pain in your back or joints, or elsewhere.

Uncommon side effects

These may affect **up to 1 in 100** people:

- ▶ clumsiness and lack of co-ordination (*ataxia*)
- ▶ double vision or blurred vision
- ▶ unusual hair loss or thinning (*alopecia*)
- ▶ skin rash or sunburn after exposure to sun or artificial light (*photosensitivity*).

Rare side effects

These may affect **up to 1 in 1,000** people:

- ▶ skin reaction that causes red spots or patches on the skin, that may look like a target or “bulls-eye” with a dark red centre surrounded by paler red rings (*erythema multiforme*)
- ▶ a life-threatening skin reaction (*Stevens-Johnson syndrome*): (see also the information at the beginning of Section 4)
- ▶ a group of symptoms together including:
fever, nausea, vomiting, headache, stiff neck and extreme sensitivity to bright light.
This may be caused by an inflammation of the membranes that cover the brain and spinal cord (*meningitis*). These symptoms usually disappear once treatment is stopped however if the symptoms continue or get worse **contact your doctor**
- ▶ rapid, uncontrollable eye movements (*nystagmus*)
- ▶ itchy eyes, with discharge and crusty eyelids (*conjunctivitis*).

Very rare side effects

These may affect **up to 1 in 10,000** people:

- ▶ a life-threatening skin reaction (*toxic epidermal necrolysis*): (see also the information at the beginning of Section 4)
- ▶ Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS): (see also the information at the beginning of Section 4)
- ▶ a high temperature (*fever*): (see also the information at the beginning of Section 4)
- ▶ swelling around the face (*oedema*) or swollen glands in the neck, armpit or groin (*lymphadenopathy*): (see also the information at the beginning of Section 4)
- ▶ changes in liver function, which will show up in blood tests, or liver failure: (see also the information at the beginning of Section 4)
- ▶ a serious disorder of blood clotting, which can cause unexpected bleeding or bruising (*disseminated intravascular coagulation*): (see also the information at the beginning of Section 4)
- ▶ haemophagocytic lymphohistiocytosis (HLH) (see Section 2: What you need to know before you take Lamotrigine)
- ▶ changes which may show up in blood tests - including reduced numbers of red blood cells (*anaemia*), reduced numbers of white blood cells (*leucopenia*, *neutropenia*, *agranulocytosis*), reduced numbers of platelets (*thrombocytopenia*), reduced numbers of all these types of cell (*pancytopenia*), and a disorder of the bone marrow called *aplastic anaemia*
- ▶ hallucinations ('seeing' or 'hearing' things that aren't really there)
- ▶ confusion
- ▶ feeling 'wobbly' or unsteady when you move about
- ▶ uncontrollable repeated body movements and/or sounds or words (*tics*), uncontrollable muscle spasms affecting the eyes, head and torso (*choreoathetosis*), or other unusual body movements such as jerking, shaking or stiffness
- ▶ in people who already have epilepsy, seizures happening more often
- ▶ in people who already have Parkinson's disease, worsening of the symptoms
- ▶ lupus-like reaction (symptoms may include: back or joint pain which sometimes may be accompanied by fever and/or general ill-health).

Other side effects

Other side effects have occurred in a small number of people but their exact frequency is unknown:

- ▶ There have been reports of bone disorders including osteopenia and osteoporosis (thinning of the bone) and fractures. Check with your doctor or pharmacist if you are on long-term anti-epileptic medication, have a history of osteoporosis, or take steroids
- ▶ inflammation of the kidney (*tubulointerstitial nephritis*), or inflammation of both the kidney and the eye (*tubulointerstitial nephritis* and *uveitis syndrome*)
- ▶ Nightmares
- ▶ Lower immunity because of lower levels of antibodies called immunoglobulins in the blood which help protect against infection.
- ▶ red nodules or patches on the skin (pseudolymphoma)

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance Website: www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Lamotrigine

- ▶ Keep this medicine out of the sight and reach of children.
- ▶ Do not use after the expiry date which is stated on the carton or bottle after 'EXP'. The expiry date refers to the last day of that month.
- ▶ Do not store above 25°C.
- ▶ Discard 60 days after first opening.

- ▶ Do not use this medicine if you notice that the suspension becomes discoloured or shows any signs of deterioration. Seek the advice of your pharmacist.
- ▶ Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Lamotrigine contains

The active substance is Lamotrigine.

Each ml of suspension contains 5mg Lamotrigine.

The other ingredients are methyl parahydroxybenzoate (E218), propyl parahydroxybenzoate (E216), simethicone emulsion [contains benzoic acid (E210)], carbomer, glycerol, saccharin sodium, sodium hydroxide (for pH adjustment), strawberry flavour [contains propylene glycol (E1520)] and purified water.

What Lamotrigine looks like and contents of the pack

Lamotrigine is a white to off-white colour suspension with odour of strawberry which is supplied in amber glass bottles with a tamper evident, child resistant white plastic screw cap.

The 150ml glass bottle is packed in a cardboard box containing a 1ml oral syringe with 0.01ml graduation mark and a 3ml oral syringe with 0.1ml graduation mark and an adaptor for the syringe.

The 300ml glass bottle is packed in a cardboard box containing a 10ml oral syringe with 0.2ml graduation mark, a 20ml dosing cup with 2.5ml, 5ml, 10ml and 15ml graduation marks and an adaptor for the syringe.

Not all pack sizes may be marketed.

Marketing Authorizations Holder:

Syri Pharma Limited t/a Thame Laboratories,
Floor 0, 1 WML,
1 Windmill Lane,
Dublin 2,
D02 F206, Ireland.

Manufacturer:

Pharmadox Healthcare Ltd.
KW20A Kordin Industrial Park,
Paola PLA 3000, Malta.

This medicine is authorised in the Member States of the European Economic Area and in the United Kingdom (Northern Ireland) under the following names:

IE and UK(NI): Lamotrigine SyriMed 5mg/ml Oral Suspension

DE: Lamotrigin Syri Pharma 5mg/ml Suspension zum Einnehmen

If this leaflet is hard to see or read, please call +353 1 437 0882 for help.

This leaflet was last revised in 06/2025.