

These are not the only side effects of JEMPERLI. If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet or in the information provided as part of an informed consent for early access treatment with JEMPERLI. By reporting side effects you can help provide more information on the safety of this medicine.

NP-IE-DST-CRD-230001 September 2023

Important Contact Information

Name of Specialist

Specialist's Phone number

After-hours Phone

Patient Name

Patient Phone Number

Patient Emergency Contact (Name)

Patient Emergency Contact (Phone)

Important Information for Health Care Providers. This patient is being treated with JEMPERLI which can cause immune-related adverse reactions that may appear any time during treatment or after treatment. Immune-related adverse reactions may occur in any organ or tissue, and may affect more than one body system simultaneously. Assess patients for signs and symptoms of immune-related adverse reactions including but not limited to those listed on this card. Early diagnosis and appropriate management are essential to minimise any consequences of immune-related adverse reactions. For suspected immune-related adverse reactions, ensure adequate evaluation to confirm aetiology or exclude other causes. Based on the severity of the adverse reaction, JEMPERLI should be withheld or permanently discontinued and corticosteroids or other appropriate therapy administered. Specific guidelines for managing immune-related adverse reactions are available in the Summary of Product Characteristics for JEMPERLI. Consultation with an oncologist or other medical specialist may be helpful for the management of immune-related adverse reactions. Please consult the Summary of Product Characteristics for JEMPERLI at www.medicines.ie or call GSK Medical Information on Tel: 1800 244 255 for more information.

Patient Card

JEMPERLI ▼ (dostarlimab)

Important safety information to minimise the risk of immune-related side effects

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. You can also report side effects directly to the HPRA Pharmacovigilance on their website: www.hpra.ie.

Contact your specialist immediately if you have any of the listed symptoms, if your symptoms get worse, or if you experience other symptoms that are not listed on this card. Getting medical treatment right away may keep the problem from becoming more serious. Your doctor may give you other medicines to treat your symptoms. Your doctor may also withhold the next dose of JEMPERLI or stop treatment.

For further information, consult the Package Information Leaflet that was provided by your treating physician or contact your treating physician.

NP-IE-DST-CRD-230001 September 2023

JEMPERLI can cause side effects that can sometimes become serious or life-threatening and can lead to death. These side effects may happen at any time during treatment or even after your treatment has ended. You may experience side effects in more than one part of your body at the same time.

IMPORTANT

- Do not attempt to diagnose or treat side effects yourself.
- Take this card with you at all times, including when you travel, if you go to the Accident and Emergency department, or when you see another doctor.
- Carry this card for at least 4 months after the last dose of JEMPERLI.
- Be sure to notify any health care professional you see that you are being treated with JEMPERLI and show them this card.

Below is a helpful checklist of Signs and Symptoms for you to closely monitor. Call your Oncology Care Team if you have any of these signs or symptoms, or if they get worse.

Lungs

Shortness of breath
Chest pain
Cough

Intestines

Diarrhea (loose, frequent bowel movements)
Black, tarry, sticky bowel movements
Blood or mucus in bowel movements
Severe stomach-area pain or tenderness

Liver

Severe nausea or vomiting
Loss of appetite
Pain on right side of the stomach
Yellowing of skin or whites of eyes
Dark urine (tea colored)
Bleeding or bruising more easily than normal

Skin

Rash, itching, peeling or blistering of skin
Sores or ulcers in the mouth, nose, throat or genital area

Kidneys

Decrease in the amount of urine
Swelling of the ankles
Blood in the urine

Hormone glands

Rapid heartbeat
Weight loss or weight gain
Changes in eyesight
Increased sweating
Hair loss
Feeling cold
Extreme tiredness
Feeling more hungry or thirsty than usual
Needing to urinate more often
Constipation
Deeper voice
Dizziness or fainting
Headaches that will not go away or unusual headache

Other

Severe or persistent muscle pain or weakness
Confusion or memory problems
Seizures
Fever or flu-like symptoms
Changes in mood or behavior
Tingling or numbness or arms or legs
Irregular heartbeat