

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Eletriptan Hormosan 20 mg film-coated tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Eletriptan Hormosan 20 mg film-coated tablets

Each film-coated tablet contains 20 mg eletriptan (as hydrobromide).

Excipients with known effect

Each film-coated tablet contains 21.85 mg lactose (as monohydrate) and 0.0602 mg sunset yellow FCF aluminium lake (E110)..

3 PHARMACEUTICAL FORM

Film-coated tablet.

Eletriptan Hormosan 20 mg film-coated tablets

Orange, round, convex shaped film-coated tablets plain on one side and embossed with "20" on the other side, approximately 6 mm in diameter.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Eletriptan Hormosan is indicated in adults for the acute treatment of the headache phase of migraine attacks, with or without aura.

4.2 Posology and method of administration

Posology

Eletriptan Hormosan film-coated tablets should be taken as early as possible after the onset of migraine headache but they are also effective if taken at a later stage during a migraine attack.

Eletriptan Hormosan, if taken during the aura phase, has not been demonstrated to prevent migraine headache and therefore Eletriptan Hormosan should only be taken during the headache phase of migraine.

Eletriptan Hormosan tablets should not be used prophylactically.

Adults (18-65 years of age)

The recommended initial dose is 40 mg.

If headache returns within 24 hours: If the migraine headache recurs within 24 hours of an initial response, a second dose of the same strength of Eletriptan Hormosan has been shown to be effective in treating the recurrence. If a second dose is required, it should not be taken within 2 hours of the initial dose.

If no response is obtained: If a patient does not achieve a headache response to the first dose of Eletriptan Hormosan within 2 hours, a second dose should not be taken for the same attack as clinical trials have not adequately established efficacy with the second dose. Clinical trials show that patients who do not respond to the treatment of an attack are still likely to respond to the treatment of a subsequent attack.

Patients who do not obtain satisfactory efficacy after an appropriate trial of 40 mg, (e.g. good tolerability and failure to respond in 2 out of 3 attacks), may be effectively treated with 80 mg (2 x 40 mg) in subsequent migraine attacks (see section 5.1). A second dose of 80 mg should not be taken within 24 hours.

The maximum daily dose should not exceed 80 mg (see section 4.8).

Elderly patients

The safety and effectiveness of eletriptan in patients over 65 years of age have not been systematically evaluated due to the small number of such patients in clinical trials. Use of Eletriptan Hormosan in the elderly is therefore not recommended.

Paediatric population

Adolescents (12-17 years of age)

The efficacy of Eletriptan Hormosan in adolescents aged 12 to 17 years has not been established. Current available data are described in section 5.2 but no recommendation on a posology can be made.

Children (6-11 years of age)

The safety and efficacy of eletriptan in children aged 6 to 11 years has not been established. Current available data are described in section 5.2 but no recommendation on a posology can be made.

Patients with hepatic impairment

No dose adjustment is required in patients with mild or moderate hepatic impairment. As eletriptan has not been studied in patients with severe hepatic impairment, it is contraindicated in these patients.

Patients with renal impairment

As the blood pressure effects of Eletriptan Hormosan are amplified in renal impairment (see section 4.4), a 20 mg initial dose, is recommended in patients with mild or moderate renal impairment. The maximum daily dose should not exceed 40 mg. Eletriptan Hormosan is contraindicated, in patients with severe renal impairment.

Method of administration

The tablets should be swallowed whole with water.

4.3 Contraindications

Eletriptan Hormosan is contraindicated in patients with

- hypersensitivity to eletriptan hydrobromide or to any of the excipients listed in 6.1.
- severe hepatic or severe renal impairment.
- moderately severe or severe hypertension, or untreated mild hypertension.
- confirmed coronary heart disease, including ischaemic heart disease (angina pectoris, previous myocardial infarction or confirmed silent ischaemia), coronary artery vasospasm (Prinzmetal's angina), objective or subjective symptoms of ischaemic heart disease.
- significant arrhythmias or heart failure.
- peripheral vascular disease.
- a history of cerebrovascular accident (CVA) or transient ischaemic attack (TIA).
- administration of ergotamine, or derivatives of ergotamine (including methysergide), within 24hr before or after treatment with eletriptan (see section 4.5).
- concomitant administration of other 5-HT₁ receptor agonists with eletriptan

4.4 Special warnings and precautions for use

Eletriptan Hormosan should not be used together with potent CYP3A4 inhibitors e.g., ketoconazole, itraconazole, erythromycin, clarithromycin, josamycin and protease inhibitors (ritonavir, indinavir and nelfinavir).

Eletriptan Hormosan should only be used where a clear diagnosis of migraine has been established.

Eletriptan Hormosan is not indicated for the management of hemiplegic, ophthalmoplegic, or basilar migraine.

Eletriptan Hormosan should not be given for the treatment of 'atypical' headaches, i.e. headaches, which may be related to a possibly serious condition (stroke, aneurysm rupture) where cerebrovascular vasoconstriction may be harmful.

Eletriptan can be associated with transient symptoms including chest pain and tightness, which may be intense and involve the throat (see section 4.8). Where such symptoms are thought to indicate ischaemic heart disease, no further dose should be taken and appropriate evaluation should be carried out.

Patients with cardiac failure

Eletriptan Hormosan should not be given without prior evaluation, to patients in whom unrecognised cardiac disease is likely, or to patients at risk of coronary artery disease (CAD) [e.g., patients with hypertension, diabetes, smokers or users of nicotine substitution therapy, men over 40 years of age, post-menopausal women and those with a strong family history of CAD]. Cardiac evaluations may not identify every patient who has cardiac disease and, in very rare cases, serious cardiac events have occurred, in patients without underlying cardiovascular disease when 5-HT₁ agonists have been administered. Patients in whom CAD is established, should not be given Eletriptan Hormosan (see section 4.3). 5-HT₁ receptor agonists have been associated with coronary vasospasm. In rare cases, myocardial ischaemia or infarction, have been reported with 5-HT₁ receptor agonists.

Undesirable effects may be more common during concomitant use of triptans and herbal preparations containing St. John's wort (*Hypericum perforatum*).

Within the clinical dose range, slight and transient increases in blood pressure have been seen with eletriptan doses of 60 mg or greater. However, these increases have not been associated with clinical sequelae in the clinical trial programme. The effect was much more pronounced in renally impaired and elderly subjects. In renally impaired subjects, the range of mean maximum increases in systolic blood pressure was 14 -17mmHg (normal 3mmHg) and for diastolic blood pressure was 14 -21mmHg (normal 4mmHg). In elderly subjects, the mean maximum increase in systolic blood pressure was 23mmHg compared with 13mmHg in young adults (placebo 8mmHg). Post-marketing reports of increases in blood pressure have also been received for patients taking 20 and 40 mg doses of eletriptan, and in non-renally impaired and non-elderly patients.

Medication overuse headache (MOH)

Prolonged use of any painkiller for headaches can make them worse. If this situation is experienced or suspected, medical advice should be obtained and treatment should be discontinued. The diagnosis of MOH should be suspected in patients who have frequent or daily headaches despite (or because of) the regular use of headache medications.

Serotonin syndrome

Serotonin syndrome (including altered mental status, autonomic instability and neuromuscular abnormalities) has been reported following concomitant treatment with triptans and selective serotonin reuptake inhibitors (SSRIs) or serotonin noradrenaline reuptake inhibitors (SNRIs).

These reactions can be severe. If concomitant treatment with eletriptan and an SSRI or SNRI is clinically warranted, appropriate observation of the patient is advised, particularly during treatment initiation, with dose increases, or with addition of another serotonergic medication (see section 4.5).

Information about excipients

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

This medicinal product contains sunset yellow, which may cause allergic reactions.

4.5 Interaction with other medicinal products and other forms of interactions

Effect of other medicinal products on eletriptan

In the pivotal clinical trials of eletriptan no evidence of interaction with beta-blockers, tricyclic antidepressants, selective serotonin reuptake inhibitors and flunarizine was reported but data from formal clinical interaction studies with these medicinal products are not available (other than propranolol, see below).

Population pharmacokinetic analysis of clinical studies has suggested that the following medicinal products (beta-blockers, tricyclic antidepressants, selective serotonin re-uptake inhibitors, oestrogen based hormone replacement therapy, oestrogen containing oral contraceptives and calcium channel blockers) are unlikely to have an effect on the pharmacokinetic properties of eletriptan.

Eletriptan is not a substrate for MAO. Therefore, there is no expectation of an interaction between eletriptan and MAO inhibitors. Therefore, no formal interaction study has been undertaken.

In clinical studies with propranolol (160 mg), verapamil (480 mg) and fluconazole (100 mg) the C_{max} of eletriptan was increased 1.1 fold, 2.2 fold and 1.4 fold respectively. The increase in eletriptan's AUC being 1.3 fold, 2.7 fold and 2.0 fold respectively. These effects are not considered clinically significant, as there were no associated increases in blood pressure or adverse events compared to administering eletriptan alone.

In clinical studies with erythromycin (1000 mg) and ketoconazole (400 mg), specific and potent inhibitors of CYP3A4, significant increases in eletriptan C_{max} (2 and 2.7- fold) and AUC (3.6 and 5.9- fold) respectively, were observed. This increased exposure was associated with an increase in eletriptan $t_{1/2}$ from 4.6 to 7.1 hours for erythromycin and from 4.8 to 8.3 hours for ketoconazole (see section 5.2). Therefore, Eletriptan Hormosan should not be used together with potent CYP3A4 inhibitors e.g., ketoconazole, itraconazole, erythromycin, clarithromycin, josamycin and protease inhibitors (ritonavir, indinavir and nelfinavir).

In clinical studies with oral (caffeine/ergotamine) administered 1 and 2 hours after eletriptan, minor though additive increases in blood pressure were observed which are predictable based on the pharmacology of the two drugs. Therefore, it is recommended that either ergotamine containing or ergot-type medications (e.g., dihydroergotamine) should not be taken within 24 hours of eletriptan dosing. Conversely, at least 24 hours should elapse after the administration of an ergotamine-containing preparation before eletriptan is given.

Effect of eletriptan on other medicinal products

There is no *in vitro* or *in vivo* evidence that clinical doses (and associated concentrations) of eletriptan will inhibit or induce cytochrome P450 enzymes including CYP3A4 drug metabolising enzymes and therefore it is considered that eletriptan is unlikely to cause clinically important drug interactions mediated by these enzymes.

Selective Serotonin Reuptake Inhibitors (SSRIs) /Serotonin Norepinephrine Reuptake Inhibitors (SNRIs) and Serotonin Syndrome

There have been reports describing patients with symptoms compatible with serotonin syndrome (including altered mental status, autonomic instability and neuromuscular abnormalities) following the use of selective serotonin reuptake inhibitors (SSRIs) or serotonin noradrenaline reuptake inhibitors (SNRIs) and triptans (see section 4.4).

4.6 Fertility, pregnancy and lactation

Pregnancy

For eletriptan no clinical data on exposed pregnancies are available. Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/fetal development, parturition or postnatal development. Eletriptan Hormosan should be used during pregnancy only if clearly needed.

Breast-feeding

Eletriptan is excreted in human breast milk. In one study of 8 women given a single dose of 80 mg, the mean total amount of eletriptan in breast milk over 24 hours in this group was 0.02% of the dose. Nevertheless, caution should be exercised when considering the administration of Eletriptan Hormosan to women who are breast-feeding. Infant exposure can be minimised by avoiding breast-feeding for 24 hours after treatment.

4.7 Effects on ability to drive and use machines

Eletriptan Hormosan has moderate influence on the ability to drive and use machines. Migraine or treatment with Eletriptan Hormosan may cause drowsiness or dizziness in some patients. Patients should be advised to evaluate their ability to perform complex tasks such as driving during migraine attacks and following administration of Eletriptan Hormosan

4.8 Undesirable effects

Summary of the safety profile

Eletriptan Hormosan has been administered in clinical trials to over 5000 subjects, taking one or two doses of eletriptan 20 or 40 or 80 mg. The most common adverse reactions noted were asthenia, somnolence, nausea and dizziness. In randomised clinical studies using doses of 20, 40 and 80 mg, a trend for a dose-dependency of the incidence of adverse events has been shown.

Tabulated list of adverse reactions

The following adverse reactions (with an incidence $\geq 1\%$ and higher than placebo) were reported in patients treated with therapeutic doses in clinical trials. Events are categorized by frequency as common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), or rare ($\geq 1/10,000$ to $< 1/1,000$).

System Organ Class	Common	Uncommon	Rare
Infections and infestations	pharyngitis, and rhinitis		respiratory tract infection
Blood and the lymphatic system disorders			lymphadenopathy
Metabolism and nutrition disorders		anorexia	
Psychiatric disorders		thinking abnormal, agitation, confusion, depersonalisation, euphoria, depression, and insomnia	emotional lability
Nervous system disorders	somnolence, headache, dizziness, tingling or abnormal sensation, hypertonia, hypoaesthesia, and myasthenia	tremor, hyperaesthesia, ataxia, hypokinesia, speech disorder, stupor, and taste perversion	
Eye disorders		abnormal vision, eye pain, photophobia, and lacrimation disorder	conjunctivitis
Ear and labyrinth disorders	vertigo	ear pain, tinnitus	
Cardiac disorders	palpitation, and tachycardia		bradycardia
Vascular disorders	flushing	peripheral vascular disorder	shock
Respiratory, thoracic and mediastinal disorders	throat tightness	dyspnea, respiratory disorder and yawning	asthma and voice alteration
Gastrointestinal disorders	abdominal pain, nausea, dry mouth, and dyspepsia	diarrhoea, and glossitis	constipation, oesophagitis, tongue oedema and eructation
Hepato-biliary disorders			hyperbilirubinaemia, and increased AST
Skin and subcutaneous tissue disorders	sweating	rash and pruritis	skin disorder and urticaria

Musculoskeletal, connective tissue and bone disorders	back pain, myalgia	arthralgia, arthrosis and bone pain	arthritis, myopathy and twitching
Renal and urinary disorders		increased urinary frequency, urinary tract disorder and polyuria	
Reproductive system and breast disorders			breast pain and menorrhagia
General disorders and administration site conditions	feeling hot, asthenia, chest symptoms (pain, tightness, pressure), chills and pain	malaise, face oedema, thirst, oedema and peripheral oedema	

The common adverse events seen with eletriptan are typical of adverse events reported with 5-HT₁ agonists as a class.

In post-marketing experience, the following undesirable effects have been reported:

Immune system disorders: allergic reactions, some of which may be serious, including angioedema

Nervous system disorders: serotonin syndrome, rare cases of syncope, cerebrovascular accident

Vascular disorders: hypertension

Cardiac disorders: myocardial ischaemia or infarction, arteriospasm coronary

Gastrointestinal disorders: as with some other 5HT 1B/1D agonists, rare reports of ischaemic colitis have been received, vomiting.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRC Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2, Tel: +353 1 6764971, Fax: +353 1 6762517, Website: www.hpra.ie, e-mail: medsafety@hpra.ie.

4.9 Overdose

Subjects have received single doses of 120 mg without significant adverse effects. However based on the pharmacology of this class, hypertension or other more serious cardiovascular symptoms could occur on overdose.

In cases of overdose, standard supportive measures should be adopted as required. The elimination half-life of eletriptan is about 4 hours, and therefore monitoring of patients and provision of general supportive therapy after overdose with eletriptan should continue for at least 20 hours or while signs and symptoms persist.

It is unknown what effect haemodialysis or peritoneal dialysis has on the serum concentrations of eletriptan.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Selective Serotonin (5HT₁) receptor agonists, ATC code: NO2CC06

Mechanism of action

Eletriptan is a selective agonist at the vascular 5-HT_{1B} and neuronal 5-HT_{1D} receptors.

Eletriptan also exhibits high affinity for the 5-HT_{1F} receptor which may contribute to its antimigraine mechanism of action. Eletriptan has modest affinity for the human recombinant 5-HT_{1A}, 5-HT_{2B}, 5-HT_{1E} and 5-HT₇ receptors.

Clinical efficacy and safety

The efficacy and safety of eletriptan in the acute treatment of migraine has been evaluated in 10 placebo-controlled trials involving more than 6000 patients (all treatment groups) at doses of 20 to 80 mg. Headache relief occurred as early as 30 minutes following oral dosing. Response rates (i.e. reduction of moderate or severe headache pain to no or mild pain) 2 hours after dosing were 59-77% for the 80 mg dose, 54-65% for the 40 mg dose, 47-54% for the 20 mg dose, and 19-40% following placebo. Eletriptan was also effective in the treatment of associated symptoms of migraine such as vomiting, nausea, photophobia and phonophobia.

The recommendation for dose titration to 80 mg is derived from open label long term studies and from a short term double blind study, where only a trend towards statistical significance was observed.

Eletriptan remains effective in menstrually associated migraine. Eletriptan, if taken during the aura phase, has not been demonstrated to prevent migraine headache and therefore eletriptan should only be taken during the headache phase of migraine.

In a non-placebo controlled pharmacokinetic study of patients with renal impairment, larger elevations in blood pressure were recorded after an 80 mg dose of eletriptan than with normal volunteers (see Section 4.4). This cannot be explained by any pharmacokinetic changes and so may represent a specific pharmacodynamic response to eletriptan in patients with renal impairment.

5.2 Pharmacokinetic properties

Absorption

Eletriptan is rapidly and well absorbed across the gastro-intestinal tract (at least 81%) after oral administration. Absolute oral bioavailability across males and females is approximately 50%. The median T_{max} is 1.5 hours after oral dosing. Linear pharmacokinetics were demonstrated over the clinical dose range (20-80 mg).

The AUC and C_{max} of eletriptan were increased by approximately 20-30% following oral administration with a high fat meal. Following oral administration during a migraine attack, there was a reduction of approximately 30% in AUC and T_{max} was increased to 2.8 hours.

Following repeated doses (20 mg three times daily) for 5-7 days, the pharmacokinetics of eletriptan remained linear and accumulation was predictable. On multiple dosing of larger doses (40 mg three times daily and 80 mg two times daily), the accumulation of eletriptan over 7 days was greater than predicted (approximately 40%).

Distribution

The volume of distribution of eletriptan following IV administration is 138L indicating distribution into the tissues. Eletriptan is only moderately protein bound (approximately 85%).

Biotransformation

In vitro studies indicate that eletriptan is primarily metabolised by hepatic cytochrome P-450 enzyme CYP3A4. This finding is substantiated by increased plasma concentrations of eletriptan following co-administration with erythromycin and ketoconazole, known selective and potent CYP3A4 inhibitors. *In vitro* studies also indicate a small involvement of CYP2D6 although clinical studies do not indicate any evidence of polymorphism with this enzyme.

There are two major circulating metabolites identified that significantly contribute to plasma radioactivity following administration of C¹⁴-labelled eletriptan. The metabolite formed by N-oxidation, has demonstrated no activity in animal *in vitro* models. The metabolite formed by N-demethylation, has been demonstrated to have similar activity to eletriptan in animal *in vitro* models. A third area of radioactivity in plasma has not been formally identified, but is most likely to be a mixture of hydroxylated metabolites which have also been observed excreted in urine and faeces.

The plasma concentrations of the N-demethylated active metabolite are only 10-20% of those of parent and so would not be expected to significantly contribute to the therapeutic action of eletriptan.

Elimination

Mean total plasma clearance of eletriptan following IV administration is 36 L/h with a resultant plasma half-life of approximately 4 hours. The mean renal clearance following oral administration is approximately 3.9 L/h. Non-renal clearance accounts for approximately 90% of the total clearance indicating that eletriptan is eliminated primarily by metabolism.

Pharmacokinetics in Special Patient Groups

Gender

A meta-analysis across clinical pharmacology studies and a population pharmacokinetic analysis of clinical trial data indicate that gender does not have any clinically significant influence on plasma concentrations of eletriptan.

Elderly (over 65 years of age)

Though not statistically significant, there is a small reduction (16%) in clearance associated with a statistically significant increased half-life (from approximately 4.4 hours to 5.7 hours) between elderly (65-93 years) and younger adult subjects.

Adolescents (12-17 years of age)

The pharmacokinetics of eletriptan (40 mg and 80 mg) in adolescent migraine patients dosed between attacks, were similar to those seen in healthy adults.

Children (6-11 years of age)

The clearance of eletriptan is unchanged in children relative to adolescents. However, the volume of distribution is lower in children resulting in higher plasma levels than would be predicted following the same dose in adults.

Patients with hepatic impairment

Subjects with hepatic impairment (Child-Pugh A and B) demonstrated a statistically significant increase in both AUC (34%) and half-life. There was a small increase in C_{max} (18%). This small change in exposure is not considered clinically relevant.

Patients with renal impairment

Subjects with mild (creatinine clearance 61-89 ml/min), moderate (creatinine clearance 31-60 ml/min) or severe (creatinine clearance <30 ml/min) renal impairment did not have any statistically significant alterations in their eletriptan pharmacokinetics or plasma protein binding. Blood pressure elevations were observed in this group.

5.3 Preclinical safety data

Preclinical data, revealed no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenicity and toxicity to reproduction.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Core tablet:

Microcrystalline cellulose
Lactose monohydrate
Croscarmellose sodium
Magnesium stearate

Film-coat:

Polyvinyl alcohol
Titanium dioxide (E171)
Macrogol
Talc
Sunset yellow FCF aluminium lake (E110)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

Eletriptan Hormosan 20 mg film-coated tablets

Opaque PVC/PCTFE/Aluminium blister packs containing 1, 2, 3, 4, 6, 10 and 18 tablets.

Eletriptan Hormosan 40 mg film-coated tablets

Opaque PVC/PCTFE/Aluminium blister packs containing 1, 2, 3, 4, 6, 10, 18 and 20 tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

No special requirements.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER

Hormosan Pharma GmbH

Hanauer Landstrasse 139 - 143

Ostend

Frankfurt Am Main

Hesse

60314

Germany

8 MARKETING AUTHORISATION NUMBER

PA23052/001/001

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of First Authorisation: 18/12/2020