

# Summary of Product Characteristics

## 1 NAME OF THE MEDICINAL PRODUCT

Clarithromycin 500 mg powder for concentrate for solution for infusion

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial contains 500 mg clarithromycin as clarithromycin lactobionate.

When reconstituted and diluted as directed, the final diluted solution contains 2 mg/ml of clarithromycin as clarithromycin lactobionate.

For the full list of excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Powder for concentrate for solution for infusion.

White to off white cake or powder.

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Clarithromycin is indicated in adults and children 12 years and older.

Clarithromycin is indicated when parenteral therapy is required for treatment of infections caused by susceptible organisms in the following conditions (see sections 4.4 and 5.1);

- Acute exacerbation of chronic bronchitis
- Community acquired pneumonia
- Acute bacterial sinusitis (adequately diagnosed)
- Streptococcal pharyngitis and tonsillitis
- Skin and soft tissue infections

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

### 4.2 Posology and method of administration

#### Posology

Intravenous therapy may be given for 2 to 5 days in the very ill patient and should be changed to oral clarithromycin therapy whenever possible as determined by the physician. The total duration of treatment with clarithromycin should not extend 14 days.

#### *Adults*

The recommended dosage of Clarithromycin 500 mg powder for concentrate for solution for infusion is 1.0 gram daily, divided into two 500 mg doses, appropriately diluted as described below.

#### *Paediatric population*

Children older than 12 years: As for adults.

Children under 12 years: Use of Clarithromycin 500 mg powder for concentrate for solution for infusion is not recommended for children younger than 12 years.

Clinical trials have been conducted using clarithromycin paediatric suspension in children 6 months to 12 years of age. Therefore, children under 12 years of age should use clarithromycin paediatric suspension (granules for oral suspension). There are insufficient data to recommend a dosage regimen for use of the clarithromycin IV formulation in patients less than 18 years of age.

### *Special populations*

#### *Elderly*

As for adults.

#### *Renal impairment*

In patients with renal impairment who have creatinine clearance less than 30 ml/min, the dosage of clarithromycin should be reduced to one half of the normal recommended dose.

#### Method of administration

For intravenous administration only.

For instructions on reconstitution/dilution of the medicinal product before administration, see section 6.6.

### **4.3 Contraindications**

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Clarithromycin 500 mg powder for concentrate for solution for infusion is contra-indicated in patients with known hypersensitivity to macrolide antibiotic.
- Concomitant administration of clarithromycin and ergot alkaloids (e.g. ergotamine or dihydroergotamine) is contraindicated, as this may result in ergot toxicity (see section 4.5).
- Concomitant administration of clarithromycin and oral midazolam is contraindicated (see section 4.5).
- Concomitant administration of clarithromycin and any of the following medicinal products is contraindicated: astemizole, cisapride, domperidone, pimozone and terfenadine as this may result in QT prolongation and cardiac arrhythmias, including ventricular tachycardia, ventricular fibrillation, and torsades de pointes (see section 4.4 and 4.5).
- Clarithromycin must not be given to patients with history of QT prolongation (congenital or documented acquired QT prolongation) or ventricular cardiac arrhythmia, including torsades de pointes (see sections 4.4 and 4.5).
- Concomitant administration with ticagrelor, ivabradine or ranolazine is contraindicated.
- Clarithromycin must not be used concomitantly with HMG-CoA reductase inhibitors (statins) that are extensively metabolized by CYP3A4, (lovastatin or simvastatin), due to the increased risk of myopathy, including rhabdomyolysis (see section 4.5).
- Concomitant administration of clarithromycin and lomitapide is contraindicated (see section 4.5).
- As with other strong CYP3A4 inhibitors, Clarithromycin must not be used in patients taking colchicine (see sections 4.4 and 4.5).
- Clarithromycin must not be given to patients with electrolyte disturbances (hypokalaemia or hypomagnesaemia, due to the risk of prolongation of the QT interval).
- Clarithromycin must not be used in patients who suffer from severe hepatic failure in combination with renal impairment (see section 4.4).

### **4.4 Special warnings and precautions for use**

#### Pregnancy:

The physician should not prescribe clarithromycin to pregnant women without carefully weighing the benefits against risk, particularly during the first and second trimester of pregnancy (see section 4.6).

#### Renal and hepatic impairment:

Clarithromycin is principally metabolised by the liver. Therefore, caution should be exercised in administering this antibiotic to patients with impaired hepatic function. Caution should also be exercised when administering clarithromycin to patients with moderate to severe renal impairment (see section 4.2). Clinical trials in patients with hepatic impairment have shown that in patients with moderate to severe hepatic impairment and normal renal function, no dose adjustment is needed.

Hepatic dysfunction, including increased liver enzymes, and hepatocellular and/or cholestatic hepatitis, with or without jaundice, has been reported with clarithromycin. This hepatic dysfunction may be severe and is usually reversible. Cases of fatal hepatic failure (see section 4.8) have been reported. Some patients may have had pre-existing hepatic disease or may have been taking other hepatotoxic medicinal products. Patients should be advised to stop treatment and contact their doctor if signs and symptoms of hepatic disease develop, such as anorexia, jaundice, dark urine, pruritus, or tender abdomen.

#### Antibiotic-associated diarrhoea and colitis:

Pseudomembranous colitis has been reported with nearly all antibacterial agents, including macrolides, and may range in severity from mild to life-threatening. *Clostridioides difficile*- associated diarrhoea (CDAD) has been reported with use of nearly all antibacterial agents including clarithromycin, and may range in severity from mild diarrhoea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon, which may lead to overgrowth of *C. difficile*. CDAD must be considered in all patients who present with diarrhoea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents. Therefore, discontinuation of clarithromycin therapy should be considered regardless of the indication. Microbial testing should be performed and adequate treatment initiated. Medicinal product inhibiting peristalsis should be avoided.

#### Interaction with medicinal products:

There have been post-marketing reports of colchicine toxicity with concomitant use of clarithromycin and colchicine, especially in the elderly, some of which occurred in patients with renal insufficiency. Deaths have been reported in some such patients (see section 4.5). Concomitant administration of clarithromycin and colchicine is contraindicated (see section 4.3).

Caution is advised regarding concomitant administration of clarithromycin and triazolobenzodiazepines, such as triazolam, and intravenous or oromucosal midazolam (see section 4.5).

Cardiovascular events: Prolongation of the QT interval, reflecting effects on cardiac repolarisation imparting a risk of developing cardiac arrhythmia and *torsade de pointes*, have been seen in patients treated with macrolides including clarithromycin (see section 4.8). Due to increased risk of QT prolongation and ventricular arrhythmias (including *torsade de pointes*), the use of clarithromycin is contraindicated: in patients taking any of astemizole, cisapride, domperidone, pimozide and terfenadine; in patients with electrolyte disorders such as hypokalaemia or hypomagnesaemia; and in patients with a history of QT prolongation or ventricular cardiac arrhythmia (see section 4.3).

Furthermore, clarithromycin should be used with caution in the following patients:

- Patients with coronary artery disease, severe cardiac insufficiency, conduction disturbances or clinically relevant bradycardia.
- Patients concomitantly taking other medicinal products associated with QT prolongation other than those which are contraindicated.

Epidemiological studies investigating the risk of adverse cardiovascular outcomes with macrolides have shown variable results. Some observational studies have identified a rare short-term risk of arrhythmia, myocardial infarction and cardiovascular mortality associated with macrolides including clarithromycin. Consideration of these findings should be balanced with treatment benefits when prescribing clarithromycin.

Pneumonia: In view of the emerging resistance of *Streptococcus pneumoniae* to macrolides, it is important that sensitivity testing be performed when prescribing clarithromycin for community-acquired pneumonia. In hospital-acquired pneumonia, clarithromycin should be used in combination with additional appropriate antibiotics.

Caution is advised when co-administering clarithromycin with other ototoxic medicinal products, especially with aminoglycosides. Monitoring of vestibular and auditory function should be performed during and after treatment.

Skin and soft tissue infections of mild to moderate severity: These infections are most often caused by *Staphylococcus aureus* and *Streptococcus pyogenes*, both of which may be resistant to macrolides. Therefore, it is important that sensitivity testing be performed. In cases where beta-lactam antibiotics cannot be used (e.g. allergy), other antibiotics, such as clindamycin, may be the medicinal product of first choice. Currently, macrolides are only considered to play a role in some skin and soft tissue

infections, such as those caused by *Corynebacterium minutissimum*, acne vulgaris, and erysipelas and in situations where penicillin treatment cannot be used.

In the event of severe acute hypersensitivity reactions, such as anaphylaxis, severe cutaneous adverse reactions (SCAR) (e.g. Acute generalised exanthematous pustulosis (AGEP) Stevens-Johnson Syndrome, toxic epidermal necrolysis, and drug rash with eosinophilia and systemic symptoms (DRESS)), clarithromycin therapy should be discontinued immediately and appropriate treatment should be urgently initiated.

Clarithromycin should be used with caution when administered concurrently with medications that induce the cytochrome CYP3A4 enzyme (see section 4.5). Clarithromycin is a CYP3A4 inhibitor and concomitant use with other medicinal products that are extensively metabolized by this enzyme should be limited to situations where clearly indicated (see section 4.5).

HMG-CoA Reductase Inhibitors (statins): Concomitant use of clarithromycin with lovastatin or simvastatin is contraindicated (see section 4.3). Caution should be exercised when prescribing clarithromycin with other statins. Rhabdomyolysis has been reported in patients taking clarithromycin and statins. Patients should be monitored for signs and symptoms of myopathy. In situations where the concomitant use of clarithromycin with statins cannot be avoided, it is recommended to prescribe the lowest registered dose of the statin. Use of the statin that is not dependent on CYP3A metabolism (e.g. fluvastatin) can be considered (see section 4.5).

Oral hypoglycaemic agents/Insulin: The concomitant use of clarithromycin and oral hypoglycaemic agents (such as sulfonylurias) and/or insulin can result in significant hypoglycaemia. Careful monitoring of glucose is recommended (see section 4.5).

Oral anticoagulants: There is a risk of serious haemorrhage and significant elevations in International Normalized Ratio (INR) and prothrombin time when clarithromycin is co-administered with warfarin (see section 4.5). INR and prothrombin times should be frequently monitored while patients are receiving clarithromycin and oral anticoagulants concurrently. Caution should be exercised when clarithromycin is co-administered with direct acting oral anticoagulants such as dabigatran, rivaroxaban, apixaban and edoxaban, particularly to patients at high risk of bleeding (see section 4.5).

The use of antimicrobial therapy of any kind, such as clarithromycin, to treat *H. pylori* infection can lead to drug-resistant microorganisms.

Long-term use may, as with other antibiotics, result in colonisation with increased numbers of non-susceptible bacteria and fungi. If superinfections occur, appropriate therapy should be initiated.

Attention should also be paid to the possibility of cross resistance between clarithromycin and other macrolide antibiotic, as well as lincomycin and clindamycin.

This medicinal product contains less than 1 mmol sodium (23 mg) per 500 mg, that is to say essentially 'sodium-free'.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

**The use of the following medicinal products is strictly contraindicated due to the potential for severe drug interaction effects:**

Astemizole, cisapride, domperidone, pimozone and terfenadine

Elevated cisapride levels have been reported in patients receiving clarithromycin and cisapride concomitantly. This may result in QT prolongation and cardiac arrhythmias including ventricular tachycardia, ventricular fibrillation and torsades de pointes. Similar effects have been observed in patients taking clarithromycin and pimozone concomitantly (see section 4.3).

Macrolides have been reported to alter the metabolism of terfenadine resulting in increased levels of terfenadine which has occasionally been associated with cardiac arrhythmias, such as QT prolongation, ventricular tachycardia, ventricular fibrillation and torsades de pointes. Concomitant use is contraindicated (see section 4.3). In one study in 14 healthy volunteers, the concomitant administration of clarithromycin and terfenadine resulted in 2- to 3-fold increase in the serum level of the acid metabolite of terfenadine and in prolongation of the QT interval which did not lead to any clinically detectable effect. Similar effects have been observed with concomitant administration of astemizole and other macrolides.

Ergot alkaloids

Post-marketing reports indicate that co-administration of clarithromycin with ergotamine or dihydroergotamine has been associated with acute ergot toxicity characterized by vasospasm, and ischaemia of the extremities and other tissues including the central nervous system. Concomitant administration of clarithromycin and ergot alkaloids is contraindicated (see section 4.3).

#### Oral Midazolam

When midazolam was co-administered with clarithromycin tablets (500 mg twice daily), midazolam AUC was increased 7-fold after oral administration of midazolam. Concomitant administration of oral midazolam and clarithromycin is contraindicated (see section 4.3).

#### HMG-CoA Reductase Inhibitors (statins)

Concomitant use of clarithromycin with lovastatin or simvastatin is contraindicated (see section 4.3) as these statins are extensively metabolized by CYP3A4 and concomitant treatment with clarithromycin increases their plasma concentration, which increases the risk of myopathy, including rhabdomyolysis. Reports of rhabdomyolysis have been received for patients taking clarithromycin concomitantly with these statins. If treatment with clarithromycin cannot be avoided, therapy with lovastatin or simvastatin must be suspended during the course of treatment.

Caution should be exercised when prescribing clarithromycin with statins. In situations where the concomitant use of clarithromycin with statins cannot be avoided, it is recommended to prescribe the lowest registered dose of the statin. Use of a statin that is not dependent on CYP3A metabolism (e.g. fluvastatin) can be considered. Patients should be monitored for signs and symptoms of myopathy.

Concomitant administration of clarithromycin with colchicine, ticagrelor and ranolazine is contraindicated (see section 4.3).

#### **The concomitant use of the following preparations is not recommended.**

The following medicinal products are known or suspected to alter the plasma concentration of clarithromycin. Dose adjustment of clarithromycin or consideration of alternative treatments may be required.

Concomitant use of clarithromycin and antifungals, azoles (ketoconazole) increases the risk of cardiac toxicity (prolonged QT interval, torsades de pointes, cardiac arrest), therefore combination therapy is not recommended.

Concomitant use of clarithromycin and nifedipine increases the risk of cardiac toxicity (prolonged QT interval, torsades de pointes, cardiac arrest), therefore combination therapy is not recommended.

Hypotension and bradycardia have been observed in a patient treated with clarithromycin and verapamil concomitantly.

Lactic acidosis has been reported with concomitant use of verapamil and a ketolide antibiotic, telithromycin (1 case), due to the possibility of CYP3A inhibition. Since clarithromycin is also a macrolide and a CYP3A inhibitor, a similar interaction with verapamil may occur.

#### **Effects of Other Medicinal Products on Clarithromycin**

Medicinal products that are inducers of CYP3A (e.g. rifampicin, phenytoin, carbamazepine, phenobarbital, St John's wort) may induce the metabolism of clarithromycin. This may result in sub-therapeutic levels of clarithromycin leading to reduced efficacy. Furthermore, it might be necessary to monitor the plasma levels of the CYP3A inducer, which could be increased owing to the inhibition of CYP3A by clarithromycin (see also the relevant product information for the CYP3A4 inducer administered).

Concomitant administration of rifabutin and clarithromycin resulted in an increase in rifabutin, and decrease in clarithromycin serum levels together with an increased risk of uveitis.

The following medicinal products are known or suspected to affect circulating concentrations of clarithromycin; clarithromycin dosage adjustment or consideration of alternative treatments may be required.

#### Efavirenz, nevirapine, rifampicin, rifabutin and rifapentine

Strong inducers of the cytochrome P450 metabolism system such as efavirenz, nevirapine, rifampicin, rifabutin, and rifapentine may accelerate the metabolism of clarithromycin and thus lower the plasma levels of clarithromycin, while increasing those of 14-OH-clarithromycin, a metabolite that is also microbiologically active. Since the microbiological activities of clarithromycin and 14-OH-clarithromycin are different for different bacteria, the intended therapeutic effect could be impaired during concomitant administration of clarithromycin and enzyme inducers.

Etravirine

Clarithromycin exposure was decreased by etravirine; however, concentrations of the active metabolite, 14-OH-clarithromycin, were increased. Because 14-OH-clarithromycin has reduced activity against *Mycobacterium avium* complex (MAC), overall activity against this pathogen may be reduced; therefore, alternatives to clarithromycin should be considered for the treatment of MAC.

Fluconazole

Concomitant administration of fluconazole 200 mg daily and clarithromycin 500 mg twice daily to 21 healthy volunteers led to increases in the mean steady-state minimum clarithromycin concentration ( $C_{min}$ ) and area under the curve (AUC) of 33% and 18% respectively. Steady state concentrations of the active metabolite 14-OH-clarithromycin were not significantly affected by concomitant administration of fluconazole. No clarithromycin dose adjustment is necessary.

Ritonavir

A pharmacokinetic study demonstrated that the concomitant administration of ritonavir 200 mg every eight hours and clarithromycin 500 mg every 12 hours resulted in a marked inhibition of the metabolism of clarithromycin. The clarithromycin  $C_{max}$  increased by 31%,  $C_{min}$  increased 182% and AUC increased by 77% with concomitant administration of ritonavir. An essentially complete inhibition of the formation of 14-OH-clarithromycin was noted. Because of the large therapeutic window for clarithromycin, no dosage reduction should be necessary in patients with normal renal function. However, for patients with renal impairment, the following dosage adjustments should be considered: For patients with  $CL_{CR}$  30 to 60 ml/min the dose of clarithromycin should be reduced by 50%. For patients with  $CL_{CR} < 30$  ml/min the dose of clarithromycin should be decreased by 75%. Doses of clarithromycin greater than 1 g/day should not be co-administered with ritonavir.

Similar dose adjustments should be considered in patients with reduced renal function when ritonavir is used as a pharmacokinetic enhancer with other HIV protease inhibitors including atazanavir and saquinavir (see section below, Bi-directional drug interactions).

Protease inhibitors (ritonavir, nelfinavir) increase the bioavailability of clarithromycin, therefore combination therapy may require dose adjustment. Similar dose adjustments should be considered in patients with renal impairment when ritonavir is used as a pharmacokinetic enhancer with other HIV protease inhibitors including atazanavir and saquinavir (see section below, Bi-directional drug interactions).

**Effect of Clarithromycin on Other Medicinal Products**CYP3A-based interactions

Co-administration of clarithromycin, which is known to inhibit CYP3A, and a medicinal product primarily metabolised by CYP3A may be associated with elevations in medicinal product concentrations that could increase or prolong both therapeutic and adverse effects of the concomitant medicinal product.

The use of clarithromycin is contraindicated in patients receiving the CYP3A substrates astemizole, cisapride, domperidone, pimozide and terfenadine due to the risk of QT prolongation and cardiac arrhythmias, including ventricular tachycardia, ventricular fibrillation, and torsades de pointes (see sections 4.3 and 4.4).

The use of clarithromycin is also contraindicated with ergot alkaloids, oral midazolam, HMG CoA reductase inhibitors metabolised mainly by CYP3A4 (e.g. lovastatin and simvastatin), colchicine, ticagrelor, ivabradine and ranolazine (see section 4.3).

Concomitant administration of clarithromycin with lomitapide is contraindicated due the potential for markedly increased transaminases (see section 4.3).

Caution is required if clarithromycin is co-administered with other medicinal products known to be CYP3A enzyme substrates, especially if the CYP3A substrate has a narrow safety margin (e.g. carbamazepine) and/or the substrate is extensively metabolised by this enzyme.

Dosage adjustments may be considered, and when possible, serum concentrations of medicinal products primarily metabolised by CYP3A should be monitored closely in patients concurrently receiving clarithromycin. Medicinal products or drug classes that are known or suspected to be metabolised by the same CYP3A isozyme include (but this list is not comprehensive) alprazolam, carbamazepine, cilostazole, ciclosporin, disopyramide, ibrutinib, methylprednisolone, midazolam (intravenous), omeprazole, oral anticoagulants (e.g. warfarin, rivaroxaban, apixaban), atypical antipsychotics (e.g. quetiapine), quinidine, rifabutin, sildenafil, sirolimus, tacrolimus, triazolam and vinblastine.

Medicinal products interacting by similar mechanisms through other isozymes within the cytochrome P450 system include phenytoin, theophylline and valproate.

### Corticosteroids

Caution should be exercised in concomitant use of clarithromycin with systemic and inhaled corticosteroids that are primarily metabolised by CYP3A due to the potential for increased systemic exposure to corticosteroids. If concomitant use occurs, patients should be closely monitored for systemic corticosteroid undesirable effects.

### Antiarrhythmics

There have been post-marketed reports of torsades de pointes occurring with the concurrent use of clarithromycin and quinidine or disopyramide. Electrocardiograms should be monitored for QT prolongation during co-administration of clarithromycin with these medicinal products. Serum levels of quinidine and disopyramide should be monitored during clarithromycin therapy.

There have been post marketing reports of hypoglycaemia with the concomitant administration of clarithromycin and disopyramide. Therefore, blood glucose levels should be monitored during concomitant administration of clarithromycin and disopyramide.

### Oral anticoagulants

Clarithromycin decreases the clearance of warfarin, therefore monitoring and dose adjustment of warfarin may be necessary. Spontaneous post-marketing reports suggest that co-administration of clarithromycin and oral anticoagulants may increase the effect of the anticoagulants. Prothrombin times should be closely monitored while patients are receiving clarithromycin and oral anticoagulants concomitantly (see section 4.4).

### Direct acting oral anticoagulants (DOACs)

The DOACs dabigatran and edoxaban are substrates for the efflux transporter P-gp. Rivaroxaban and apixaban are metabolised via CYP3A4 and are also substrates for P-gp. Caution should be exercised when clarithromycin is co-administered with these agents particularly to patients at high risk of bleeding (see section 4.4).

### Oral hypoglycemic agents/Insulin

With certain hypoglycaemic medicinal products such as nateglinide, and repaglinide, inhibition of CYP3A enzyme by clarithromycin may be involved and could cause hypoglycaemia when used concomitantly. Careful monitoring of glucose is recommended.

### Omeprazole

Clarithromycin (500 mg every 8 hours) was given in combination with omeprazole (40 mg daily) to healthy adult subjects. The steady-state plasma concentrations of omeprazole were increased ( $C_{max}$ , AUC<sub>0-24</sub>, and  $t_{1/2}$  increased by 30%, 89%, and 34%, respectively), by the concomitant administration of clarithromycin. The mean 24-hour gastric pH value was 5.2 when omeprazole was administered alone and 5.7 when omeprazole was co-administered with clarithromycin.

### Sildenafil, tadalafil and vardenafil

Each of these phosphodiesterase inhibitors is metabolised, at least in part, by CYP3A, and CYP3A may be inhibited by concomitantly administered clarithromycin. Co-administration of clarithromycin with sildenafil, tadalafil or vardenafil would likely result in increased phosphodiesterase inhibitor exposure. Reduction of sildenafil, tadalafil and vardenafil dosages should be considered when these drugs are co-administered with clarithromycin.

### Theophylline, carbamazepine

Results of clinical studies indicate that there was a modest but statistically significant ( $p \leq 0.05$ ) increase of circulating theophylline or carbamazepine levels when either of these medicinal products were administered concomitantly with clarithromycin. Dose reduction may need to be considered.

### Tolterodine

The primary route of metabolism for tolterodine is via the 2D6 isoform of cytochrome P450 (CYP2D6). However, in a subset of the population devoid of CYP2D6, the identified pathway of metabolism is via CYP3A. In this population subset, inhibition of CYP3A results in significantly higher serum concentrations of tolterodine. A reduction in tolterodine dosage may be necessary in the presence of CYP3A inhibitors, such as clarithromycin in the CYP2D6 poor metaboliser population.

### Triazolobenzodiazepines (e.g., alprazolam, midazolam, triazolam)

When midazolam was co-administered with clarithromycin tablets (500 mg twice daily), midazolam AUC was increased 2.7-fold after intravenous administration of midazolam. If intravenous midazolam is co-administered with clarithromycin, the patient must be closely monitored to allow dose adjustment. Drug delivery of midazolam via oromucosal route, which could bypass pre-systemic elimination of the medicinal product, will likely result in a similar interaction to that observed after intravenous

midazolam rather than oral administration. The same precautions should also apply to other benzodiazepines that are metabolised by CYP3A, including triazolam and alprazolam. For benzodiazepines which are not dependent on CYP3A for their elimination (temazepam, nitrazepam, lorazepam), a clinically important interaction with clarithromycin is unlikely.

Clarithromycin thus increases the bioavailability of benzodiazepines and benzodiazepine derivatives (midazolam, alprazolam, diazepam, triazolam). Combination therapy may require dose adjustment.

There have been post-marketing reports of drug interactions and central nervous system (CNS) effects (e.g., somnolence and confusion) with the concomitant use of clarithromycin and triazolam. Monitoring the patient for increased CNS pharmacological effects is suggested.

### ***Other drug interactions***

#### Colchicine

Colchicine is a substrate for both CYP3A and the efflux transporter, P-glycoprotein (Pgp). Clarithromycin and other macrolides are known to inhibit CYP3A and Pgp. When clarithromycin and colchicine are administered together, inhibition of Pgp and/or CYP3A by clarithromycin may lead to increased exposure to colchicine (see section 4.3 and 4.4).

#### Digoxin

Digoxin is thought to be a substrate for the efflux transporter, P-glycoprotein (Pgp). Clarithromycin is known to inhibit Pgp. When clarithromycin and digoxin are administered together, inhibition of Pgp by clarithromycin may lead to increased exposure to digoxin. Elevated digoxin serum concentrations in patients receiving clarithromycin and digoxin concomitantly have also been reported in post marketing surveillance. Some patients have shown clinical signs consistent with digoxin toxicity, including potentially fatal arrhythmias. Serum digoxin concentrations should be carefully monitored while patients are receiving digoxin and clarithromycin simultaneously.

#### Zidovudine

Simultaneous oral administration of clarithromycin tablets and zidovudine to HIV-infected adult patients may result in decreased steady-state zidovudine concentrations. Because clarithromycin appears to interfere with the absorption of simultaneously administered oral zidovudine, this interaction can be largely avoided by staggering the doses of clarithromycin and zidovudine to allow for a 4-hour interval between each medication. This interaction does not appear to occur in paediatric HIV-infected patients taking clarithromycin suspension with zidovudine or dideoxyinosine. This interaction is unlikely when clarithromycin is administered via intravenous infusion.

#### Phenytoin and Valproate

There have been spontaneous or published reports of interactions of CYP3A inhibitors, including clarithromycin with medicinal products not thought to be metabolised by CYP3A (e.g. phenytoin and valproate). Serum level determinations are recommended for these medicinal products when administered concomitantly with clarithromycin. Increased serum levels have been reported.

#### Hydroxychloroquine and chloroquine

Clarithromycin should be used with caution in patients receiving these medicines known to prolong the QT interval due to the potential to induce cardiac arrhythmia and serious adverse cardiovascular events.

#### Fluoxetine

Co-administration of clarithromycin and fluoxetine increases the risk of psychiatric side effects.

### ***Bi-directional drug interactions***

#### Atazanavir

Both clarithromycin and atazanavir are substrates and inhibitors of CYP3A, and there is evidence of a bi-directional drug interaction. Co-administration of clarithromycin (500 mg twice daily) with atazanavir (400 mg once daily) resulted in a 2-fold increase in exposure to clarithromycin and a 70% decrease in exposure to 14-OH-clarithromycin, with a 28% increase in the AUC of atazanavir. Because of the large therapeutic window for clarithromycin, no dosage reduction should be necessary in patients with normal renal function. For patients with moderate renal function (creatinine clearance 30 to 60 ml/min), the dose of clarithromycin should be decreased by 50%. For patients with creatinine clearance <30 ml/min, the dose of clarithromycin should be decreased by 75% using an appropriate clarithromycin formulation. Doses of clarithromycin greater than 1000 mg per day should not be co-administered with protease inhibitors.

### Calcium Channel Blockers

Caution is advised regarding the concomitant administration of clarithromycin and calcium channel blockers metabolized by CYP3A4 (e.g. verapamil, amlodipine, diltiazem) due to the risk of hypotension. Plasma concentrations of clarithromycin as well as calcium channel blockers may increase due to the interaction. Hypotension, bradyarrhythmias and lactic acidosis have been observed in patients taking clarithromycin and verapamil concomitantly.

Concomitant use of clarithromycin and verapamil increases the risk of cardiac toxicity (prolonged QT interval, torsades de pointes, cardiac arrest).

### Itraconazole

Both clarithromycin and itraconazole are substrates and inhibitors of CYP3A, leading to a bidirectional drug interaction. Clarithromycin may increase the plasma levels of itraconazole, while itraconazole may increase the plasma levels of clarithromycin. Patients taking itraconazole and clarithromycin concomitantly should be monitored closely for signs or symptoms of increased or prolonged pharmacologic effect.

### Saquinavir

Both clarithromycin and saquinavir are substrates and inhibitors of CYP3A, and there is evidence of a bi-directional drug interaction. Concomitant administration of clarithromycin (500 mg twice daily) and saquinavir (soft gelatin capsules, 1200 mg three times daily) to 12 healthy volunteers resulted in steady-state AUC and  $C_{max}$  values of saquinavir which were 177% and 187% higher than those seen with saquinavir alone. Clarithromycin AUC and  $C_{max}$  values were approximately 40% higher than those seen with clarithromycin alone. No dose adjustment is required when the two medicinal products are co-administered for a limited time at the doses/formulations studied. Observations from drug interaction studies using the soft gelatin capsule formulation may not be representative of the effects seen using the saquinavir hard gelatin capsule. Observations from drug interaction studies performed with saquinavir alone may not be representative of the effects seen with saquinavir/ritonavir therapy. When saquinavir is co-administered with ritonavir, consideration should be given to the potential effects of ritonavir, consideration should be given to the potential effects of ritonavir on clarithromycin (see section 4.5: Ritonavir)

Patients taking oral contraceptives should be warned that if diarrhoea vomiting or breakthrough bleeding occur there is a possibility of contraceptive failure.

## **4.6 Fertility, pregnancy and lactation**

### Pregnancy

The safety of clarithromycin for use in pregnancy has not been established. Based on variable results obtained from animal studies and experience in humans, the possibility of adverse effects on embryofetal development cannot be excluded. Some observational studies evaluating exposure to clarithromycin during the first and second trimester have reported an increased risk of miscarriage compared to no antibiotic use or other antibiotic use during the same period. The available epidemiological studies on the risk of major congenital malformations with use of macrolides including clarithromycin during pregnancy provide conflicting results.

Therefore, use during pregnancy is not advised without carefully weighing the benefits against risks.

### Breast-feeding

Clarithromycin is excreted into human breast milk in small amounts. It has been estimated that an exclusively breastfed infant would receive about 1.7% of the maternal weight-adjusted dose of clarithromycin. Therefore, diarrhoea and fungus infection of the mucous membranes could occur in the breast-fed infant, so that nursing might have to be discontinued. The possibility of sensitisation should be considered. The benefit of treatment of the mother should be weighed against the potential risk for the infant.

### Fertility

There is no data available on the effect of clarithromycin on fertility in humans. In rats, the limited data available do not indicate any effects on fertility.

## **4.7 Effects on ability to drive and use machines**

There are no data on the effect of clarithromycin on the ability to drive or use machines. The potential for dizziness, vertigo, confusion and disorientation, which may occur with the medication, should be taken into account before patients drive or use machines.

## 4.8 Undesirable effects

### a. Summary of the safety profile

The most frequent and common adverse reactions related to clarithromycin therapy for both adult and paediatric populations are abdominal pain, diarrhoea, nausea, vomiting and taste perversion. These adverse reactions are usually mild in intensity and are consistent with the known safety profile of macrolide antibiotics (see section b of section 4.8).

There was no significant difference in the incidence of these gastrointestinal adverse reactions during clinical trials between the patient population with or without pre-existing mycobacterial infections.

### b. Tabulated summary of adverse reactions

The following table displays adverse reactions reported in clinical trials and from post-marketing experience with clarithromycin immediate-release tablets, granules for oral suspension, powder for solution for injection, extended-release tablets and modified-release tablets.

The reactions considered at least possibly related to clarithromycin are displayed by system organ class and frequency using the following convention: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ) and not known (adverse reactions from post-marketing experience; cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness when the seriousness could be assessed.

System Organ Class	Very common $\geq 1/10$	Common $\geq 1/100$ to $< 1/10$	Uncommon $\geq 1/1,000$ to $< 1/100$	Rare $\geq 1/10,000$ to $< 1/1,000$	Very Rare $< 1/10,000$	Not Known* (cannot be estimated from the available data)
Infections and infestations		Oral candidiasis	Cellulitis <sup>1</sup> , candidiasis, gastroenteritis <sup>2</sup> , infection <sup>3</sup> , vaginal infection			Pseudomembranous colitis, erysipelas,
Blood and lymphatic system			Leukopenia, neutropenia <sup>4</sup> , thrombocythaemia <sup>3</sup> , eosinophilia <sup>4</sup>			Agranulocytosis, thrombocytopenia
Immune system disorders			Anaphylactoid reaction <sup>1</sup> , hypersensitivity			Anaphylactic reaction. angioedema
Metabolism and nutrition disorders			Anorexia, decreased appetite			
Psychiatric disorders		Insomnia	Anxiety, nervousness <sup>3</sup>			Psychotic disorder, confusional state, depersonalisation, depression, disorientation, hallucination, abnormal dreams, mania
Nervous system disorders		Dysgeusia, headache, altered sense of smell	Loss of consciousness <sup>1</sup> , dyskinesia <sup>1</sup> , dizziness, somnolence <sup>6</sup> , tremor		Brain fog	Convulsion, ageusia, parosmia, anosmia, paraesthesia, cramp
Ear and labyrinth disorders			Vertigo, hearing impaired, tinnitus	Reversible hearing loss		Deafness
Cardiac disorders			Cardiac arrest <sup>1</sup> , atrial fibrillation <sup>1</sup> , electrocardiogram QT prolonged <sup>7</sup> , extrasystoles <sup>1</sup> ,			Torsades de pointes <sup>7</sup> , ventricular tachycardia <sup>7</sup> , ventricular fibrillation

			palpitations			
Vascular disorders		Vasodilation <sup>1</sup>				Haemorrhage <sup>8</sup>
Respiratory, thoracic and mediastinal disorder			Asthma <sup>1</sup> , epistaxis <sup>2</sup> , pulmonary embolism <sup>1</sup>			
Gastrointestinal disorders		Diarrhoea <sup>9</sup> , vomiting, dyspepsia, nausea, abdominal pain	Oesophagitis <sup>1</sup> , gastrooesophageal reflux disease <sup>2</sup> , gastritis, proctalgia <sup>2</sup> , stomatitis, glossitis, abdominal distension <sup>4</sup> , constipation, dry mouth, eructation, flatulence,			Pancreatitis acute, tongue discolouration, tooth discolouration
Hepatobiliary disorders		Liver function test abnormal	Hepatic dysfunction (usually transient and reversible), Jaundice, Cholestasis <sup>4</sup> , hepatitis <sup>4</sup> , alanine aminotransferase increased, aspartate aminotransferase increased, gamma-glutamyltransferase increased <sup>4</sup>			Hepatic failure, jaundice hepatocellular
Skin and subcutaneous tissue disorders		Rash, hyperhidrosis	Dermatitis bullous <sup>1</sup> , pruritus, urticaria, rash maculo-papular <sup>3</sup> , exanthema			Severe cutaneous adverse reactions (SCAR) (e.g Acute generalised exanthematous pustulosis (AGEP), Stevens-Johnson syndrome <sup>5</sup> , toxic epidermal necrolysis <sup>5</sup> , drug rash with eosinophilia and systemic symptoms (DRESS)), acne
Musculoskeletal and connective tissue disorders			Muscle spasms <sup>3</sup> , musculoskeletal stiffness <sup>1</sup> , myalgia <sup>2</sup> , arthralgia			Rhabdomyolysis <sup>2,11</sup> , myopathy
Renal and urinary disorders			Blood creatinine increased <sup>1</sup> , blood urea increased <sup>1</sup>		Renal insufficiency	Renal failure, nephritis interstitial
General disorders and administration site conditions	Injection site phlebitis <sup>1</sup>	Injection site pain <sup>1</sup> , injection site inflammation <sup>1</sup>	Malaise <sup>4</sup> , pyrexia <sup>3</sup> , asthenia, chest pain <sup>4</sup> , chills <sup>4</sup> , fatigue <sup>4</sup>			
Investigations		Elevated plasma urea	Elevated plasma creatinine, Elevated liver transaminases, Albumin globulin ratio abnormal <sup>1</sup> , blood alkaline phosphatase increased <sup>4</sup> , blood lactate dehydrogenase			International normalised ratio increased <sup>8</sup> , prothrombin time prolonged <sup>8</sup> , urine colour abnormal

<sup>1</sup> ADRs reported only for the powder for concentrate for solution for infusion

<sup>2</sup> ADRs reported only for the extended-release tablets

<sup>3</sup> ADRs reported only for the granules for oral suspension

<sup>4</sup> ADRs reported only for the immediate-release tablets

<sup>5, 7, 9, 10</sup> See section a)

<sup>6, 8, 11</sup> See section c)

\* *Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. Patient exposure is estimated to be greater than 1 billion patient treatment days for clarithromycin.*

#### c. Description of selected adverse reactions

Injection site phlebitis, injection site pain, and injection site inflammation are specific to the clarithromycin intravenous formulation.

In some of the reports of rhabdomyolysis, clarithromycin was administered concomitantly with statins, fibrates, colchicine or allopurinol (see section 4.3 and 4.4).

There have been post-marketing reports of drug interactions and central nervous system (CNS) effects (e.g. somnolence and confusion) with the concomitant use of clarithromycin and triazolam. Monitoring the patient for increased CNS pharmacological effects is suggested (see section 4.5).

There have been rare reports of clarithromycin extended-release tablets in the stool, many of which have occurred in patients with anatomic (including ileostomy or colostomy) or functional gastrointestinal disorders with shortened GI transit times. In several reports, tablet residues have occurred in the context of diarrhoea. It is recommended that patients who experience tablet residue in the stool and no improvement in their condition should be switched to a different clarithromycin formulation (e.g. suspension) or another antibiotic.

Special population: Adverse Reactions in Immunocompromised Patients (see section e).

#### d. Paediatric populations

Clinical trials have been conducted using clarithromycin paediatric suspension in children 6 months to 12 years of age. Therefore, children under 12 years of age should use clarithromycin paediatric suspension.

Frequency, type and severity of adverse reactions in children are expected to be the same as in adults.

#### e. Other special populations

##### *Immunocompromised patients*

In AIDS and other immunocompromised patients treated with the higher doses of clarithromycin over long periods of time for mycobacterial infections, it was often difficult to distinguish adverse events possibly associated with clarithromycin administration from underlying signs of Human Immunodeficiency Virus (HIV) disease or intercurrent illness.

In adult patients, the most frequently reported adverse reactions by patients treated with total daily doses of 1000 mg and 2000 mg of clarithromycin were: nausea, vomiting, taste perversion, abdominal pain, diarrhoea, rash, flatulence, headache, constipation, hearing disturbance, Serum Glutamic Oxaloacetic Transaminase (SGOT) and Serum Glutamic Pyruvate Transaminase (SGPT) elevations. Additional low-frequency events included dyspnoea, insomnia and dry mouth. The incidences were comparable for patients treated with 1000 mg and 2000 mg, but were generally about 3 to 4 times as frequent for those patients who received total daily doses of 4000 mg of clarithromycin.

In these immunocompromised patients, evaluations of laboratory values were made by analysing those values outside the seriously abnormal level (i.e. the extreme high or low limit) for the specified test. On the basis of these criteria, about 2% to 3% of those patients who received 1000 mg or 2000 mg of clarithromycin daily had seriously abnormal elevated levels of SGOT and SGPT, and abnormally low white blood cell and platelet counts. A lower percentage of patients in these two dosage groups also had elevated Blood Urea Nitrogen levels. Slightly higher incidences of abnormal values were noted for patients who received 4000 mg daily for all parameters except White Blood Cell.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system:

HPRA Pharmacovigilance

Website: [www.hpra.ie](http://www.hpra.ie)

## 4.9 Overdose

### Symptoms

Reports indicate that the ingestion of large amounts of clarithromycin orally can be expected to produce gastro-intestinal symptoms. One patient who had a history of bipolar disorder ingested 8 grams of clarithromycin and showed altered mental status, paranoid behaviour, hypokalaemia and hypoxaemia.

### Treatment

Adverse reactions accompanying overdosage should be treated by the prompt elimination of unabsorbed medicinal product and supportive measures. As with other macrolides, clarithromycin serum levels are not expected to be appreciably affected by haemodialysis or peritoneal dialysis.

In the case of overdosage, treatment should be discontinued and all other appropriate supportive measures should be instituted.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

#### Pharmacotherapeutic group:

Antibacterial for systemic use, macrolide, ATC code: J01FA09

#### Mechanism of action

Clarithromycin is an antibiotic belonging to the macrolide antibiotic group. It exerts its antibacterial action by selectively binding to the 50S ribosomal sub-unit of susceptible bacteria preventing translocation of activated amino acids. It inhibits the intracellular protein synthesis of susceptible bacteria.

The 14-hydroxy metabolite of clarithromycin, a product of parent drug metabolism also has anti-microbial activity. The metabolite is less active than the parent compound for most organisms, including *Mycobacterium* spp. An exception is *Haemophilus influenzae* where the 14-hydroxy metabolite is two-fold more active than the parent compound.

#### Resistance

The mechanisms of acquired resistance in macrolides are: efflux of active substance by an active pump mechanism, inducible or constitutive production of a methylase enzyme that modifies the ribosomal target, hydrolysis of macrolides by esterases, chromosomal mutations that alter a 50S ribosomal protein. Cross-resistance between clarithromycin and other macrolides and clindamycin and lincomycin may therefore occur. Methicillin-resistant and oxacillin-resistant staphylococci (MRSA) and penicillin-resistant *Streptococcus pneumoniae* are resistant to all currently available Beta-lactam antibiotics and macrolides such as clarithromycin.

#### Susceptibility testing breakpoints

MIC (minimum inhibitory concentration) interpretive criteria for susceptibility testing have been established by the European Committee on Antimicrobial Susceptibility Testing (EUCAST) for clarithromycin and are listed here:

[https://www.ema.europa.eu/documents/other/minimum-inhibitory-concentration-mic-breakpoints\\_en.xlsx](https://www.ema.europa.eu/documents/other/minimum-inhibitory-concentration-mic-breakpoints_en.xlsx).

#### Susceptibility

The prevalence of acquired resistance may vary geographically and over time for selected species and local information on resistance is desirable, particularly when treating severe infections. Expert advice should be sought as needed when the local prevalence of resistance is such that the usefulness of a medicinal product, at least for some types of infections, is questionable. Clarithromycin has a pronounced effect on a wide range of aerobic, anaerobic, Gram-positive, Gram-negative and acid-resistant bacteria.

The activity of 14(R)-hydroxyclearithromycin is greater than clarithromycin against *Haemophilus influenzae*. In vitro studies suggest that 14(R)-hydroxyclearithromycin -and its parent molecule have additive activity against *H. influenzae*.

<b>Commonly susceptible species</b>
<b>Aerobic, Gram-positive microorganisms</b>
<i>Streptococcus agalactiae</i>
<i>Streptococcus pyogenes</i> (Group A beta-haemolytic streptococci)
<i>Streptococcus viridans</i>
<i>Streptococcus pneumoniae</i>
<i>Staphylococcus aureus</i> (methicillin susceptible)

<i>Listeria monocytogenes</i>
<b>Aerobic, Gram-negative microorganisms</b>
<i>Bordetella pertusis</i>
<i>Haemophilus influenzae</i>
<i>Haemophilus parainfluenzae</i>
<i>Helicobacter pylori</i>
<i>Campylobacter jejuni</i>
<i>Moraxella catarrhalis</i>
<i>Neisseria gonorrhoeae</i>
<i>Legionella</i> spp.
<b>Anaerobic microorganisms</b>
<i>Clostridium perfringens</i>
<i>Bacterioides fragilis</i>
<i>Peptococcus/Peptostreptococcus</i> spp.
<i>Propionibacterium acnes</i>
<b>Other microorganisms</b>
<i>Mycoplasma pneumoniae</i>
<i>Chlamydia trachomatis</i>
<i>Chlamydia pneumoniae</i>
<i>Ureaplasma urealyticum</i>
<i>Mycobacterium</i> spp.

#### Other information

Susceptibility and resistance of *Streptococcus pneumoniae* and *Streptococcus* spp. to clarithromycin can be predicted by testing erythromycin.

## 5.2 Pharmacokinetic properties

### Distribution

Following IV administration, the blood levels of clarithromycin achieved are well in excess of the MIC<sub>90s</sub> for the common pathogens and the levels of 14-hydroxyclearithromycin exceed the necessary concentrations for important pathogens, e.g. *H. influenzae*. The microbiologically active metabolite 14-hydroxyclearithromycin is formed by first pass metabolism as indicated by lower bioavailability of the metabolite following IV administration.

Clarithromycin gives good penetration into different compartments. Clarithromycin provides tissue concentrations that are several times higher than the circulating active substance levels. Increased levels have been found in both tonsillar and lung tissue. Clarithromycin also penetrates the gastric mucus.

Clarithromycin is 80% bound to plasma proteins at therapeutic levels.

### Serum half-life

The serum half-life of the active 14-(R)-hydroxy metabolite ranges between 5 to 6 hours.

### Biotransformation and elimination

Clarithromycin is rapidly and extensively metabolised in the liver. Metabolism involves mainly N-dealkylation, oxidation and stereospecific hydroxylation at position C14.

### Linearity

The pharmacokinetics of clarithromycin and the 14-hydroxy metabolite are non-linear; steady state is achieved by day 3 of IV dosing. Following a single 500 mg IV dose over 60 minutes, about 33% clarithromycin and 11% 14-hydroxyclearithromycin is excreted in the urine at 24 hours.

## 5.3 Preclinical safety data

In 4-week-studies in animals, toxicity of clarithromycin was found to be related to the dose and to the duration of the treatment. In all species, the first signs of toxicity were observed in the liver, in which lesions were seen within 14 days in dogs and monkeys. The systemic levels of exposure, related to this toxicity, are not known in detail, but toxic doses (300 mg/kg/day) were clearly higher than the therapeutic doses recommended for humans. Other tissues affected included the stomach, thymus

and other lymphoid tissues as well as the kidneys. At near therapeutic doses conjunctival injection and lacrimation occurred only in dogs. At a dose of 400mg/kg/day some dogs and monkeys developed corneal opacities and/or oedema.

In vitro and in vivo studies showed that clarithromycin did not have genotoxic potential.

No mutagenic effects were found in in vitro- and in vivo -studies with clarithromycin.

Studies on reproduction toxicity showed that administration of clarithromycin at doses 2x the clinical dose in rabbit (iv) and 10x the clinical dose in monkey (po) resulted in an increased incidence of spontaneous abortions. These doses were related to maternal toxicity. No embryotoxicity or teratogenicity was generally noted in rat studies. However, cardiovascular malformations were observed in two studies in rats treated with doses of 150 mg/kg/d.

In mice at doses 70x the clinical dose, cleft palate occurred at varying incidence (3-30%).

Clarithromycin has been found in the milk of lactating animals.

In 3-day old mice and rats, the LD50 values were approximately half those in adult animals. Juvenile animals presented similar toxicity profiles to mature animals although enhanced nephrotoxicity in neonatal rats has been reported in some studies. Slight reductions in erythrocytes, platelets and leukocytes have also been found in juvenile animals.

Clarithromycin has not been tested for carcinogenicity.

## 6 PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

Sodium hydroxide (pH adjuster)

### 6.2 Incompatibilities

None known.

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

### 6.3 Shelf life

*Unopened:* 5 years

*After reconstitution:*

Chemical and physical in-use stability has been demonstrated for 24 hours at 25°C / 48 hours at 2-8°C for the reconstituted solution.

*After dilution:*

Chemical and physical in-use stability has been demonstrated 6 hours at 25°C / 48 hours at 2-8°C for the final infusion solution.

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8°C, unless reconstitution / dilution has taken place in controlled and validated aseptic conditions.

### 6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

For storage conditions after reconstitution/dilution of the medicinal product, see section 6.3.

### 6.5 Nature and contents of container

Uncoloured clear glass vial type I (15 ml), sealed with bromobutyl rubber stopper and aluminium cap with plastic flip-off seal.

The following pack sizes are available for Clarithromycin 500 mg powder for concentrate for solution for infusion:

1, 5 or 10 vial(s) containing 500 mg of clarithromycin.

Not all pack sizes may be marketed.

## 6.6 Special precautions for disposal and other handling

Clarithromycin 500 mg powder for concentrate for solution for infusion should be administered into one of the larger proximal veins as an IV infusion over 60 minutes, using a solution concentration of about 2 mg/ml. Clarithromycin should not be given as a bolus or an intramuscular injection.

### *Preparation for use*

Prepare all solutions using aseptic techniques.

#### Reconstitution (Step 1)

Clarithromycin 500 mg powder for concentrate for solution for infusion is reconstituted with 10 ml of sterile water for injections, a solution with a concentration of 50 mg/ml is obtained. Shake until the vial contents have dissolved. The reconstitution time is not more than 7 minutes and the solution is clear to slightly opalescent, colourless to slightly yellow. Use only water for injections, as other diluents may cause precipitation during reconstitution. Do not use diluents containing preservatives or inorganic salts.

For storage conditions after reconstitution of the medicinal product, see section 6.3.

#### Dilution (Step 2)

The reconstituted solution should be added to 250 ml of a suitable intravenous diluent prior to infusion: 0.9% Sodium chloride, 5% Dextrose, 5% Dextrose in 0.3% Sodium chloride, 5% Dextrose in 0.45% Sodium chloride, 5% Dextrose in Ringer's lactate solution and Ringer's lactate solution. The solution after dilution is clear to slightly opalescent, colourless to slightly yellow.

The concentration of Clarithromycin for the final solution for infusion is 2 mg/ml.

For storage conditions after dilution of the medicinal product, see section 6.3.

IMPORTANT: BOTH DILUENT STEPS (1 and 2) SHOULD BE COMPLETED BEFORE USE.

For single use only.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## 7 MARKETING AUTHORISATION HOLDER

hameln pharma gmbh  
Inselstraße 1  
31787 Hameln  
Germany

## 8 MARKETING AUTHORISATION NUMBER

PA2237/003/001

## 9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 19<sup>th</sup> March 2021

Date of last renewal: 26<sup>th</sup> April 2023

## 10 DATE OF REVISION OF THE TEXT

April 2025