

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Torsemide 10 mg Tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 10 mg of torsemide.

Excipient(s) with known effect: Each tablet contains 166.60 mg of lactose monohydrate

For a full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Tablets.

White to off white, round tablets, with the imprint "T 10" and scoreline on one side and plain on other side.

Tablet dimension: 8.00 ± 0.20 mm

The score line is only to facilitate breaking for ease of swallowing and not to divide into equal doses.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Oedema due to congestive heart failure; hepatic, pulmonary or renal oedema.

4.2 Posology and method of administration

Adults

Oedema: The usual dose is 5 mg. once daily. If necessary, the dose can be increased stepwise up to 20 mg once daily. In individual cases, as much as 40 mg torsemide/day has been administered.

A starting dose of 5 mg is not achievable with this formulation, therefore another formulation should be used.

Elderly

No special dosage adjustments are necessary.

Children

There is no experience of torsemide in children.

4.3 Contraindications

Renal failure with anuria; hepatic coma and pre-coma; hypotension; pre-existing hypovolaemia; pregnancy and lactation; hypersensitivity to torsemide and sulphonylureas or to any of the excipients listed in section 6.1; cardiac arrhythmias, simultaneous therapy with aminoglycosides or cephalosporins, or renal dysfunction due to drugs which cause renal damage.

4.4 Special warnings and precautions for use

Hypokalaemia, hyponatraemia, hypovolaemia and disorders of micturition must be corrected before treatment.

On long-term treatment with torsemide, regular monitoring of the electrolyte balance, glucose, uric acid, creatinine and lipids in the blood, is recommended.

Careful monitoring of patients with a tendency to hyperuricaemia and gout is recommended.

Carbohydrate metabolism in latent or manifest diabetes mellitus should be monitored.

As for other drugs which produce changes in blood pressure, patients taking torasemide should be warned not to drive or operate machinery if they experience dizziness or related symptoms.

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.

Difficulty with micturition

Particular caution is required in patients with difficulty with micturition including prostatic hypertrophy because they have an increased risk of developing acute urinary retention and require careful close monitoring.

4.5 Interaction with other medicinal products and other forms of interaction

When used simultaneously with cardiac glycosides, a potassium and/or magnesium deficiency may increase sensitivity of the cardiac muscle to such drugs. The kaliuretic effect of mineralo- and glucocorticoids and laxatives may be increased.

As with other diuretics, the effect of antihypertensive drugs given concomitantly may be potentiated.

Torasemide, especially at high doses, may potentiate the toxicity of aminoglycoside antibiotics, cisplatin preparations, the nephrotoxic effects of cephalosporins, and the cardio- and neurotoxic effect of lithium. The action of curare-containing muscle relaxants and of theophylline can be potentiated. In patients receiving high doses of salicylates, salicylate toxicity may be increased. The action of anti-diabetic drugs may be reduced.

Sequential or combined treatment, or starting a new co-medication with an ACE inhibitor may result in transient hypotension. This may be minimised by lowering the starting dose of the ACE inhibitor and/or reducing or stopping temporarily the dose of torasemide. Torasemide may decrease arterial responsiveness to pressor agents e.g. adrenaline, noradrenaline.

Non-steroidal anti-inflammatory drugs (eg. Indometacin) and probenecid may reduce the diuretic and hypotensive effect of torasemide.

Concomitant use of torasemide and colestyramine has not been studied in humans, but in an animal study co-administration of colestyramine decreased absorption of oral torasemide.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no data from experience in humans of the effect of torasemide on the embryo and foetus. Animal studies have shown reproductive toxicity (see section 5.3). Consequently, torasemide is contra-indicated in pregnancy and lactation.

Lactation

No studies have been conducted on excretion in breast milk.

Fertility

No effects on fertility have been seen in animal studies.

4.7 Effects on ability to drive and use machines

As for other drugs which produce changes in blood pressure, patients taking torasemide should be warned not to drive or operate machinery if they experience dizziness or related symptoms.

4.8 Undesirable effects

Within the system organ classes, adverse reactions are listed under headings of frequency (number of patients expected to experience the reaction), using the following categories:

Very common ($\geq 1/10$)

Common ($\geq 1/100$ to $< 1/10$)

Uncommon ($\geq 1/1,000$ to $< 1/100$)

Rare ($\geq 1/10,000$ to $< 1/1,000$)

Very rare ($< 1/10,000$)

Not known (cannot be estimated from available data)

The following undesirable effects were observed whereas the frequency of undesirable effect is not known:

Blood and lymphatic system disorders

Frequency not known: Thrombocytopenia, Leukopenia, Anaemia

Immune system disorders

Very rare: Allergic skin reactions (e.g. Pruritus, Exanthema), Photosensitivity reaction

Frequency not known: Serious skin reactions (e.g. Stevens-Johnson syndrome, Toxic epidermal necrolysis)

Metabolism and nutrition disorders

Common: Metabolic alkalosis, Fluid and electrolyte imbalance (e.g. Hypovolaemia, Hyponatraemia)

Nervous system disorders

Common: Headache, Dizziness

Frequency not known: Cerebral ischaemia, Paresthesia, confusional state

Eye disorders

Frequency not known: Visual impairment

Ear and labyrinth disorders

Frequency not known: tinnitus, Deafness

Cardiac disorders

Frequency not known: Acute myocardial infarction, Myocardial ischaemia, Angina pectoris, Syncope, Hypotension

Vascular disorders

Frequency not known: Embolism

Gastrointestinal disorders

Common: Gastrointestinal disorder, (e.g. Loss of appetite, abdominal pain upper, Nausea, Vomiting, Diarrhoea, Constipation)

Frequency not known: Dry mouth, Pancreatitis

Hepatobiliary disorders

Uncommon: Hepatic enzyme increased (e.g. Gamma-glutamyltransferase increased)

Skin and subcutaneous tissue disorders

Very rare: Allergic skin reactions (e.g. Pruritus, Exanthema), Photosensitivity reaction

Frequency not known: Serious skin reactions (e.g. Stevens-Johnson syndrome, Toxic epidermal necrolysis)

Musculoskeletal and connective tissue disorders

Common: Muscle spasms

Renal and urinary disorders

Uncommon: Urinary retention, Bladder dilatation

Rare: Blood urea increased, Blood creatinine increased

General disorders and administration site conditions

Common: Fatigue, Asthenia

Investigations

Uncommon: Blood uric acid increased, Blood glucose increased, Lipids increased (e.g. Blood triglycerides increased, Blood cholesterol increased)

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRa Pharmacovigilance, Website: www.hpra.ie.

4.9 Overdose

Symptoms and signs

No typical picture of intoxication is known. If overdosage occurs, then there may be marked diuresis with the danger of loss of fluid and electrolytes which may lead to somnolence, confusion, hypotension, hyponatremia, hypokalemia, hypochloremic alkalosis, hemoconcentration dehydration and circulatory collapse. Gastrointestinal disturbances may occur.

Treatment

No specific antidote is known. Symptoms and signs of overdosage require the reduction of the dose or withdrawal of torasemide, and simultaneous replacement of fluid and electrolytes.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: High ceiling diuretics, sulphonamide monodrugs, ATC code: C03CA04

Torasemide is a loop diuretic. However, at low doses its pharmacodynamic profile resembles that of the thiazide class regarding the level and duration of diuresis. At higher doses, torasemide induces a brisk diuresis in a dose dependant manner with a high ceiling of effect.

Torasemide acts as a salidiuretic by inhibition of renal sodium and chloride reabsorption in the ascending limb of the loop of Henle. After oral administration the onset of diuresis is within the 1st hour with a peak action within 2 to 3h. The action may last up to 12h.

In healthy subjects an increase in dose results in a linear increase in urine excretion corresponding to the logarithm of the dose (high-ceiling activity) within the 5 to 100 mg dose range. An increase in diuresis may also take place if other diuretics are no longer active, eg in the presence of impaired renal function.

In renal failure endogenous organic acids compete with loop diuretics for the acid secretion mechanism in the proximal tubule. Therefore, the torasemide dose has to be adequately increased in order to achieve effective amounts of drug at the site of action.

Torasemide leads to a gentle removal of edema and especially to an improvement of the working condition of the heart failure by reducing the preload and afterload. In patients with severe to endstage chronic renal failure there is a reduction of arterial blood pressure in addition to removal of edema and maintenance of residual diuresis.

5.2 Pharmacokinetic properties

Absorption

Torasemide is absorbed rapidly and almost completely after oral administration, and peak serum levels are reached after one to two hours.

Serum protein binding

More than 99% of torasemide is bound to plasma proteins.

Distribution

The apparent distribution volume is 16 litres.

Metabolism

Torasemide is metabolised to three metabolites, M1, M3 and M5 by stepwise oxidation, hydroxylation or ring hydroxylation. Further metabolites (M2 and M4) have been found in animal experiments, but not in humans.

Elimination

The terminal half-life of torasemide and its metabolites is three to four hours in healthy subjects. Total clearance of torasemide is 40ml/min and renal clearance about 10ml/min. About 80% of the dose administered is excreted as torasemide and metabolites into the renal tubule - torasemide 24%, M1 12%, M3 3%, M5 41%.

In patients with congestive heart failure and disorders of liver function, the elimination half-lives of torasemide and metabolite M5 are only slightly increased compared with those in healthy volunteers. The amounts of torasemide and metabolites excreted in the urine are similar to those in healthy subjects; therefore no accumulation is to be expected.

In the presence of renal failure, elimination half-life of torasemide is unchanged.

5.3 Preclinical safety data

Preclinical data reveal no special hazard for humans based on single dose toxicity, genotoxicity and carcinogenicity studies.

In repeat dose toxicity studies in dogs and rats weight reduction, increases in creatinine and urea and renal alterations such as tubular dilatation and interstitial nephritis were observed. These changes, which occurred at high doses are attributable, to an excess pharmacodynamic action (diuresis) and were shown to be reversible.

Reproduction toxicology studies in the rat have shown no teratogenic effect, but foetal and maternal toxicity have been observed after high doses in pregnant rabbits and rats. No effects on fertility have been seen.

Torasemide showed no mutagenic potential. Carcinogenicity studies in rats and mice showed no tumourigenic potential.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Lactose monohydrate
Pregelatinised Starch
Colloidal Anhydrous silica
Magnesium stearate

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

30 months.

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

PVC/PVDC Clear Alu-blister pack, containing 10, 14, 28 and 30 tablets
PVC Clear Alu-blister pack, containing 10, 14, 28 and 30 tablets

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Not applicable

7 MARKETING AUTHORISATION HOLDER

Azure Pharmaceuticals Ltd.
12 Hamilton Drive
The Rock Road
Blackrock
Co. Louth

A91 T997

Ireland

8 MARKETING AUTHORISATION NUMBER

PA22871/014/003

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date for first authorisation: 11th October 2024

10 DATE OF REVISION OF THE TEXT