

Package leaflet: Information for the user

Duloxetine Accord 30 mg hard gastro-resistant capsules

Duloxetine Accord 60 mg hard gastro-resistant capsules

duloxetine (as hydrochloride)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any of the side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Duloxetine Accord is and what it is used for
2. What you need to know before you take Duloxetine Accord
3. How to take Duloxetine Accord
4. Possible side effects
5. How to store Duloxetine Accord
6. Contents of the pack and other information

1. What Duloxetine Accord is and what it is used for

Duloxetine Accord contains the active substance duloxetine. Duloxetine Accord increases the levels of serotonin and noradrenaline in the nervous system.

Duloxetine Accord is used in adults to treat:

- depression
- generalised anxiety disorder (chronic feeling of anxiety or nervousness)
- diabetic neuropathic pain (often described as burning, stabbing, stinging, shooting or aching or like an electric shock. There may be loss of feeling in the affected area, or sensations such as touch, heat, cold or pressure may cause pain)

Duloxetine Accord starts to work in most people with depression or anxiety within two weeks of starting treatment, but it may take 2-4 weeks before you feel better. Tell your doctor if you do not start to feel better after this time. Your doctor may continue to give you Duloxetine Accord when you are feeling better to prevent your depression or anxiety from returning.

In people with diabetic neuropathic pain, it can take some weeks before you feel better. Talk to your doctor if you do not feel better after 2 months.

2. What you need to know before you take Duloxetine Accord

Do not take Duloxetine Accord if you:

- are allergic to duloxetine or any of the other ingredients of this medicine (listed in section 6)
- have liver disease
- have severe kidney disease
- are taking or have taken within the last 14 days, another medicine known as a monoamine oxidase inhibitor (MAOI) (see ‘Other medicines and Duloxetine Accord’)

- are taking fluvoxamine, which is usually used to treat depression, ciprofloxacin or enoxacin which are used to treat some infections
- are taking other medicines containing duloxetine (see ‘Other medicines and Duloxetine Accord’)

Talk to your doctor if you have high blood pressure or heart disease. Your doctor will tell you if you should be taking Duloxetine Accord.

Warnings and precautions

The following are reasons why Duloxetine Accord may not be suitable for you. Talk to your doctor before taking Duloxetine Accord if you:

- are taking other medicines to treat depression or buprenorphine. The use of these medicines together with Duloxetine Accord can lead to serotonin syndrome, a potentially life-threatening condition (see “Other medicines and Duloxetine Accord”).
- are taking St. John’s Wort, a herbal treatment (*Hypericum perforatum*)
- have kidney disease
- have had seizures (fits)
- have had mania
- suffer from bipolar disorder
- have eye problems, such as certain kinds of glaucoma (increased pressure in the eye)
- have a history of bleeding disorders (tendency to develop bruises), especially if you are pregnant (see 'Pregnancy and breast-feeding')
- are at risk of low sodium levels (for example if you are taking diuretics, especially if you are elderly)
- are currently being treated with another medicine which may cause liver damage
- are taking other medicines containing duloxetine (see ‘Other medicines and Duloxetine Accord’)

Duloxetine Accord may cause a sensation of restlessness or an inability to sit or stand still. You should tell your doctor if this happens to you.

Medicines like Duloxetine Accord (so called SSRIs/SNRIs) may cause symptoms of sexual dysfunction (see section 4). In some cases, these symptoms have continued after stopping treatment.

Thoughts of suicide and worsening of your depression or anxiety disorder

If you are depressed and/or have anxiety disorders, you can sometimes have thoughts of harming or killing yourself. These may be increased when first starting antidepressants, since these medicines all take time to work, usually about two weeks but sometimes longer.

You may be more likely to think like this if you:

- have previously had thoughts about killing or harming yourself
- are a young adult. Information from clinical trials has shown an increased risk of suicidal behaviour in adults aged less than 25 years with psychiatric conditions who were treated with an antidepressant

If you have thoughts of harming or killing yourself at any time, contact your doctor or go to a hospital straight away.

You may find it helpful to tell a relative or close friend that you are depressed or have an anxiety disorder and ask them to read this leaflet. You might ask them to tell you if they think your depression or anxiety is getting worse, or if they are worried about changes in your behaviour.

You should also contact your doctor:

If you experience signs and symptoms of restlessness, hallucinations, loss of coordination, fast heart beat, increased body temperature, fast changes in blood pressure, overactive reflexes, diarrhoea, coma, nausea, vomiting, as you might be suffering a serotonin syndrome.

In its most severe form, serotonin syndrome can resemble Neuroleptic Malignant Syndrome (NMS). Signs and symptoms of NMS may include a combination of fever, fast heart beat, sweating, severe muscle stiffness, confusion, increased muscle enzymes (determined by a blood test).

Children and adolescents under 18 years of age

Duloxetine Accord should normally not be used for children and adolescents under 18 years. Also, you should know that patients under 18 have an increased risk of side-effects such as suicide attempt, suicidal thoughts and hostility (predominantly aggression, oppositional behaviour and anger) when they take this class of medicines. Despite this, your doctor may prescribe Duloxetine Accord for patients under 18 because he/she decides that this is in their best interests. If your doctor has prescribed Duloxetine Accord for a patient under 18 and you want to discuss this, please go back to your doctor. You should inform your doctor if any of the symptoms listed above develop or worsen when patients under 18 are taking Duloxetine Accord. Also, the long-term safety effects concerning growth, maturation, and cognitive and behavioural development of Duloxetine Accord in this age group have not yet been demonstrated.

Other medicines and Duloxetine Accord

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

The active substance of Duloxetine Accord, duloxetine, is used in other medicines for other conditions:

- diabetic neuropathic pain, depression, anxiety and urinary incontinence

Using more than one of these medicines at the same time should be avoided. Check with your doctor if you are already taking other medicines containing duloxetine.

Your doctor should decide whether you can take Duloxetine Accord with other medicines. Do not start or stop taking any medicines, including those bought without a prescription and herbal remedies, before checking with your doctor.

You should also tell your doctor if you are taking any of the following:

Monoamine oxidase inhibitors (MAOIs): You should not take Duloxetine Accord if you are taking, or have recently taken (within the last 14 days) another antidepressant medicine called a monoamine oxidase inhibitor (MAOI). Examples of MAOIs include moclobemide (an antidepressant) and linezolid (an antibiotic). Taking a MAOI together with many prescription medicines, including Duloxetine Accord, can cause serious or even life-threatening side effects. You must wait at least 14 days after you have stopped taking an MAOI before you can take Duloxetine Accord. Also, you need to wait at least 5 days after you stop taking Duloxetine Accord before you take a MAOI.

Medicines that cause sleepiness: These include medicines prescribed by your doctor including benzodiazepines, strong painkillers, antipsychotics, phenobarbital and antihistamines.

Medicines that increase the level of serotonin: Triptans, tryptophan, SSRIs (such as paroxetine and fluoxetine), SNRIs (such as venlafaxine), tricyclic antidepressants (such as clomipramine, amitriptyline), St John's Wort, MAOIs (such as moclobemide and linezolid), buprenorphine, tramadol and pethidine. These medicines may interact with Duloxetine Accord, and you may experience symptoms such as involuntary, rhythmic contractions of muscles, including the muscles that control movement of the eye, agitation, hallucinations, coma, excessive sweating, tremor, exaggeration of reflexes, increased muscle tension, body temperature above 38°C. Contact

your doctor when experiencing such symptoms, as these might indicate a potentially life-threatening condition known as serotonin syndrome.

Oral anticoagulants or antiplatelet agents: Medicines which thin the blood or prevent the blood from clotting. These medicines might increase the risk of bleeding.

Duloxetine Accord with food, drink and alcohol

Duloxetine Accord may be taken with or without food.

Care should be taken if you drink alcohol while you are being treated with Duloxetine Accord.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

- Tell your doctor if you become pregnant, or you are trying to become pregnant, while you are taking Duloxetine Accord. You should use Duloxetine Accord only after discussing the potential benefits and any potential risks to your unborn child with your doctor.

Make sure your midwife and/or doctor knows you are on Duloxetine Accord. When taken during pregnancy, similar drugs (SSRIs) may increase the risk of a serious condition in babies, called persistent pulmonary hypertension of the newborn (PPHN), making the baby breathe faster and appear bluish. These symptoms usually begin during the first 24 hours after the baby is born. If this happens to your baby, you should contact your midwife and/or doctor immediately.

If you take Duloxetine Accord near the end of your pregnancy, your baby might have some symptoms when it is born. These usually begin at birth or within a few days of your baby being born. These symptoms may include floppy muscles, trembling, jitteriness, not feeding properly, trouble with breathing and fits. If your baby has any of these symptoms when it is born, or you are concerned about your baby's health, contact your doctor or midwife who will be able to advise you.

If you take Duloxetine Accord near the end of your pregnancy there is an increased risk of excessive vaginal bleeding shortly after birth, especially if you have a history of bleeding disorders. Your doctor or midwife should be aware that you are taking duloxetine so they can advise you.

Available data from the use of Duloxetine Accord during the first three months of pregnancy do not show an increased risk of overall birth defects in general in the child. If Duloxetine Accord is taken during the second half of pregnancy, there may be an increased risk that the infant will be born early (6 additional premature infants for every 100 women who take Duloxetine Accord in the second half of pregnancy), mostly between weeks 35 and 36 of pregnancy.

Tell your doctor if you are breast-feeding. The use of Duloxetine Accord while breastfeeding is not recommended. You should ask your doctor or pharmacist for advice.

Driving and using machines

Duloxetine Accord may make you feel sleepy or dizzy. Do not drive or use any tools or machines until you know how Duloxetine Accord affects you.

Duloxetine Accord contains sucrose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Duloxetine Accord contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

3. How to take Duloxetine Accord

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Duloxetine Accord is for oral use. You should swallow your capsule whole with a drink of water.

For depression and diabetic neuropathic pain:

The usual dose of Duloxetine Accord is 60 mg once a day, but your doctor will prescribe the dose that is right for you.

For generalised anxiety disorder:

The usual starting dose of Duloxetine Accord is 30 mg once a day after which most patients will receive 60 mg once a day, but your doctor will prescribe the dose that is right for you. The dose may be adjusted up to 120 mg a day based on your response to Duloxetine Accord.

To help you remember to take Duloxetine Accord, you may find it easier to take it at the same times every day.

Talk with your doctor about how long you should keep taking Duloxetine Accord. Do not stop taking Duloxetine Accord, or change your dose, without talking to your doctor. Treating your disorder properly is important to help you get better. If it is not treated, your condition may not go away and may become more serious and difficult to treat.

If you take more Duloxetine Accord than you should

Call your doctor or pharmacist immediately if you take more than the amount of Duloxetine Accord prescribed by your doctor. Symptoms of overdose include sleepiness, coma, serotonin syndrome (a rare reaction which may cause feelings of great happiness, drowsiness, clumsiness, restlessness, feeling of being drunk, fever, sweating or rigid muscles), fits, vomiting and fast heart rate.

If you forget to take Duloxetine Accord

If you miss a dose, take it as soon as you remember. However, if it is time for your next dose, skip the missed dose and take only a single dose as usual. Do not take a double dose to make up for a forgotten dose. Do not take more than the daily amount of Duloxetine Accord that has been prescribed for you in one day.

If you stop taking Duloxetine Accord

DO NOT stop taking your capsules without the advice of your doctor even if you feel better. If your doctor thinks that you no longer need Duloxetine Accord, he or she will ask you to reduce your dose over at least 2 weeks before stopping treatment altogether.

Some patients who stop taking Duloxetine Accord suddenly have had symptoms such as:

- dizziness, tingling feelings like pins and needles or electric shock-like feelings (particularly in the head), sleep disturbances (vivid dreams, nightmares, inability to sleep), fatigue, sleepiness, feeling restless or agitated, feeling anxious, feeling sick (nausea) or being sick (vomiting), shaking (tremor), headaches, muscle pain, feeling irritable, diarrhoea, excessive sweating or vertigo.

These symptoms are usually not serious and disappear within a few days, but if you have symptoms that are troublesome you should ask your doctor for advice.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. These effects are normally mild to moderate and often disappear after a few weeks.

Very common side effects (may affect more than 1 in 10 people)

- headache, feeling sleepy
- feeling sick (nausea), dry mouth

Common side effects (may affect up to 1 in 10 people)

- lack of appetite
- trouble sleeping, feeling agitated, less sex drive, anxiety, difficulty or failure to experience orgasm, unusual dreams
- dizziness, feeling sluggish, tremor, numbness, including numbness, pricking or tingling of the skin
- blurred eyesight
- tinnitus (hearing sound in the ear when there is no external sound)
- feeling the heart pumping in the chest,
- increased blood pressure, flushing
- increased yawning
- constipation, diarrhoea, stomach pain, being sick (vomiting), heartburn or indigestion, breaking wind
- increased sweating, (itchy) rash
- muscle pain, muscle spasm
- painful urination, frequent urination
- problems getting an erection, changes in ejaculation
- falls (mostly in elderly people), fatigue
- weight loss

Children and adolescents under 18 years of age with depression treated with this medicine had some weight loss when they first start taking this medicine. Weight increased to match other children and adolescents of their age and sex after 6 months of treatment.

Uncommon side effects (may affect up to 1 in 100 people)

- throat inflammation that causes a hoarse voice
- suicidal thoughts, difficulty sleeping, grinding or clenching the teeth, feeling disorientated, lack of motivation
- sudden involuntary jerks or twitches of the muscles, sensation of restlessness or an inability to sit or stand still, feeling nervous, difficulty concentrating, changes in sense of taste, difficulty controlling movement e.g. lack of coordination or involuntary movements of the muscles, restless legs syndrome, poor sleep quality
- large pupils (the dark centre of the eye), problems with eyesight
- feeling of dizziness or “spinning” (vertigo), ear pain
- fast and/or irregular heartbeat
- fainting, dizziness, lightheadedness or fainting on standing up, cold fingers and/or toes
- throat tightness, nose bleeds
- vomiting blood, or black tarry stools (faeces), gastroenteritis, burping, difficulty swallowing
- inflammation of the liver that may cause abdominal pain and yellowing of the skin or whites of the eyes

- night sweats, hives, cold sweats, sensitivity to sunlight, increased tendency to bruise
- muscle tightness, muscle twitching
- difficulty or inability to pass urine, difficulty to start urinating, needing to pass urine during the night, needing to pass more urine than normal, having a decreased urine flow
- abnormal vaginal bleeding, abnormal periods, including heavy, painful, irregular or prolonged periods, unusually light or missed periods, pain in the testicles or scrotum
- chest pain, feeling cold, thirst, shivering, feeling hot, abnormal gait
- weight gain
- Duloxetine Accord may cause effects that you may not be aware of, such as increases in liver enzymes or blood levels of potassium, creatine phosphokinase, sugar, or cholesterol

Rare side effects (may affect up to 1 in 1000 people)

- serious allergic reaction which causes difficulty in breathing or dizziness with swollen tongue or lips, allergic reactions
- decreased thyroid gland activity which can cause tiredness or weight gain
- dehydration, low levels of sodium in the blood (mostly in elderly people; the symptoms may include feeling dizzy, weak, confused, sleepy or very tired, or feeling or being sick, more serious symptoms are fainting, fits or falls), syndrome of inappropriate secretion of anti-diuretic hormone (SIADH)
- suicidal behaviour, mania (over activity, racing thoughts and decreased need for sleep), hallucinations, aggression and anger
- “Serotonin syndrome” (a rare reaction which may cause feelings of great happiness, drowsiness, clumsiness, restlessness, feeling of being drunk, fever, sweating or rigid muscles), fits
- increased pressure in the eye (glaucoma)
- inflammation of the mouth, passing bright red blood in your stools, bad breath, inflammation of the large intestine (leading to diarrhoea)
- liver failure, yellowing of the skin or whites of the eyes (jaundice)
- Stevens-Johnson syndrome (serious illness with blistering of the skin, mouth, eyes and genitals), serious allergic reaction which causes swelling of the face or throat (angioedema)
- contraction of the jaw muscle
- abnormal urine odour
- menopausal symptoms, abnormal production of breast milk in men or women
- coughing, wheezing and shortness of breath which may be accompanied by a high temperature
- excessive vaginal bleeding shortly after birth (postpartum haemorrhage)

Very rare side effects (may affect up to 1 in 10,000 people)

- inflammation of the blood vessels in the skin (cutaneous vasculitis)

Frequency not known (cannot be estimated from the available data)

- Signs and symptoms of a condition called “stress cardiomyopathy” which may include chest pain, shortness of breath, dizziness, fainting, irregular heart beat

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via

HPRA Pharmacovigilance,

Website: www.hpra.ie

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Duloxetine Accord

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and carton. The expiry date refers to the last day of that month.

Blister PVC/PVDC-/Alu: Do not store above 30°C. Keep the blister in the outer carton. Store in the original package in order to protect from light.

Blister PVC/PVDC-Alu: Do not store above 30°C.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Duloxetine Accord contains

- The active substance is duloxetine.
30 mg capsules: Each gastro-resistant capsule, hard contains 30 mg of duloxetine (as hydrochloride).
60 mg capsules: Each gastro-resistant capsule, hard contains 60 mg of duloxetine (as hydrochloride).

- The other ingredients are:
Capsule content: Hypromellose, talc, microcrystalline cellulose, methacrylic acid-ethyl acrylate copolymer (1:1) dispersion 30%, triethyl citrate, sugar spheres, cysteine hydrochloride monohydrate and sodium hydroxide solution..

Capsule shell: gelatin, titanium dioxide (E171), Indigo carmine (E132), printing ink (black iron oxide (E-172), potassium hydroxide and shellac), yellow iron oxide (E172) (60 mg only).

What Duloxetine Accord looks like and contents of the pack

Duloxetine Accord is a gastro-resistant capsule, hard. Each capsule of Duloxetine Accord contains pellets of duloxetine hydrochloride with a covering to protect them from stomach acid.

The 30 mg capsules are printed (cap E/body 127), opaque blue and opaque white.

The 60 mg capsules are printed (cap E/body 129) opaque blue and opaque green.

Duloxetine Accord is available in:

PVC-PVDC/Alu blister:

7, 10, 14, 20, 28, 30, 50, 56, 60, 84, 98 and 504 (4x126) (hospital pack) hard gastro-resistant capsules

PVC-PVDC/Alu blister:

7, 10, 14, 20, 28, 30, 50, 56, 60, 84, 98 and 504 (4x126) (hospital pack) hard gastro-resistant capsules

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:

Accord Healthcare Ireland Ltd,
Euro House,
Euro Business Park,

Little Island,
Cork T45 K857,
Ireland

Manufacturer:

TOWA Pharmaceutical Europe, S.L.
C/ de Sant Martí, 75-97, Martorelles, 08107 Barcelona, Spain

This medicinal product is authorised in the Member States of the European Economic Area under the following names:

Name of the Member State	Name of the medicine
Germany	Dutilox 30 mg magensaftresistente Hartkapseln Dutilox 60 mg magensaftresistente Hartkapseln
Austria	Dutilox 30 mg magensaftresistente Hartkapseln Dutilox 60 mg magensaftresistente Hartkapseln
Netherland	Dutilox 30 mg maagsapresistente capsules, harde Dutilox 60 mg maagsapresistente capsules, harde
Belgium	{FR}: Dutilox 30 mg, gélules gastro-résistantes {NL}: Dutilox 30 mg maagsapresistente capsules, hard {DE}: Dutilox 30 mg magensaftresistente Hartkapseln {FR}: Dutilox 60 mg, gélules gastro-résistantes {NL}: Dutilox 60 mg maagsapresistente capsules, hard {DE}: Dutilox 60 mg magensaftresistente Hartkapseln
Poland	Dutilox
Romania	Duloxetina Terapia 30 mg capsule gastrorezistente Duloxetina Terapia 60 mg capsule gastrorezistente
Ireland	Duloxetine Accord 30 mg hard gastro-resistant capsules Duloxetine Accord 60 mg hard gastro-resistant capsules

This leaflet was last revised in ~~July 2026~~ August 2026