

Package Leaflet: Information for the patient

Levetiracetam Clonmel 100 mg/ml oral solution levetiracetam

Read all of this leaflet carefully before you or your child start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Levetiracetam Clonmel is and what it is used for
2. What you need to know before you take Levetiracetam Clonmel
3. How to take Levetiracetam Clonmel
4. Possible side effects
5. How to store Levetiracetam Clonmel
6. Contents of the pack and other information

1. What Levetiracetam Clonmel is and what it is used for

Levetiracetam is an antiepileptic medicine (a medicine used to treat seizures in epilepsy).

Levetiracetam Clonmel is used:

- on its own in adults and adolescents from 16 years of age with newly diagnosed epilepsy, to treat a certain form of epilepsy. Epilepsy is a condition where the patients have repeated fits (seizures). Levetiracetam is used for the epilepsy form in which the fits initially affect only one side of the brain, but could thereafter extend to larger areas on both sides of the brain (partial onset seizure with or without secondary generalisation). Levetiracetam has been given to you by your doctor to reduce the number of fits.
- as an add-on to other antiepileptic medicines to treat:
 - partial onset seizures with or without generalisation in adults, adolescents, children and infants from one month of age
 - myoclonic seizures (short, shock-like jerks of a muscle or group of muscles) in adults and adolescents from 12 years of age with juvenile myoclonic epilepsy
 - primary generalised tonic-clonic seizures (major fits, including loss of consciousness) in adults and adolescents from 12 years of age with idiopathic generalised epilepsy (the type of epilepsy that is thought to have a genetic cause)

2. What you need to know before you take Levetiracetam Clonmel

DO NOT take Levetiracetam Clonmel:

- if you are allergic to levetiracetam, pyrrolidone derivatives or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor or pharmacist before taking Levetiracetam Clonmel

- if you suffer from kidney problems, follow your doctor's instructions. He/she may decide if your dose should be adjusted.
- if you notice any slowdown in the growth or unexpected puberty development of your child, please contact your doctor.
- A small number of people being treated with anti-epileptics such as Levetiracetam Clonmel have had thoughts of harming or killing themselves. If you have any symptoms of depression and/or suicidal ideation, please contact your doctor.
- If you have a family or medical history of irregular heart rhythm (visible on an electrocardiogram),

or if you have a disease and/or take a treatment that make(s) you prone to heartbeat irregularities or salt imbalances.

Tell your doctor or pharmacist if any of the following side effects gets serious or last longer than a few days:

- Abnormal thoughts, feeling irritable or reacting more aggressively than usually, or if you or your family and friends notice important changes in mood or behaviour.
- Aggravation of epilepsy
Your seizures may rarely become worse or happen more often, mainly during the first month after the start of the treatment or increase of the dose. If you experience any of these new symptoms while taking Levetiracetam Clonmel, see a doctor as soon as possible.

Children and adolescents

Levetiracetam Clonmel is not indicated in children and adolescents below 16 years on its own (monotherapy).

Other medicines and Levetiracetam Clonmel

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Do not take macrogol (a medicine used as laxative) for one hour before and one hour after taking levetiracetam as this may result in a reduction of its effect.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Levetiracetam can be used during pregnancy, only if after careful assessment it is considered necessary by your doctor.

You should not stop your treatment without discussing this with your doctor. A risk of birth defects for your unborn child cannot be completely excluded. Two studies do not suggest an increased risk of autism or intellectual disability in children born to mothers treated with levetiracetam during pregnancy. However, the available data regarding the impact of levetiracetam on neurodevelopment in children is limited.

Breast-feeding is not recommended during treatment.

Driving and using machines

Levetiracetam Clonmel may impair your ability to drive or operate any tools or machinery, as it may make you feel sleepy. This is more likely at the beginning of treatment or after an increase in the dose. You should not drive or use machines until it is established that your ability to perform such activities is not affected.

Levetiracetam Clonmel contains methyl parahydroxybenzoate (E218), liquid maltitol (E965), sodium and benzyl alcohol

Levetiracetam Clonmel oral solution contains methyl parahydroxybenzoate which may cause allergic reactions (possibly delayed).

Levetiracetam Clonmel oral solution also contains liquid maltitol. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per ml that is to say essentially ‘sodium-free’.

This medicine contains 0.0016 mg benzyl alcohol in each ml. Benzyl alcohol may cause allergic reactions.

Do not use for more than a week in young children (less than 3 years old), unless advised by your doctor or pharmacist.

Ask your doctor or pharmacist for advice if you are pregnant or breast-feeding. This is because large amounts of benzyl alcohol can build-up in your body and may cause side effects (called “metabolic acidosis”).

Ask your doctor or pharmacist for advice if you have a liver or kidney disease. This is because large amounts of benzyl alcohol can build-up in your body and may cause side effects (called “metabolic acidosis”).

3. How to take Levetiracetam Clonmel

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Levetiracetam Clonmel must be taken twice a day, once in the morning and once in the evening, at about the same time each day.

Take the oral solution following your doctor’s instructions.

Monotherapy (from 16 years of age)

Adults (≥ 18 years) and adolescents (from 16 years of age):

Measure the appropriate dosage using the 10 ml syringe included in the package for patients 4 years and above.

General dose: Levetiracetam Clonmel is taken twice daily, in two equally divided doses, each individual dose being measured between 5 ml (500 mg) and 15 ml (1,500 mg).

When you will first start taking Levetiracetam Clonmel, your doctor will prescribe you a **lower dose** during 2 weeks before giving you the lowest general dose.

Add-on therapy

Dose in adults and adolescents (12 to 17 years):

Measure the appropriate dosage using the 10 ml syringe included in the package for patients of 4 years and above.

General dose: Levetiracetam Clonmel is taken twice daily, in two equally divided doses, each individual dose being measured between 5 ml (500 mg) and 15 ml (1,500 mg).

Dose in children 6 months and older:

Your doctor will prescribe the most appropriate pharmaceutical form of levetiracetam according to the age, weight and dose.

For children 6 months to 4 years, measure the appropriate dosage using the **3 ml** syringe included in the package.

For children above 4 years, measure the appropriate dosage using the **10 ml** syringe included in the package.

General dose: Levetiracetam Clonmel is taken twice daily, in two equally divided doses, each individual dose being measured between 0.1 ml (10 mg) and 0.3 ml (30 mg), per kg bodyweight of the child (see table below for dose examples).

Dose in children 6 months and older:

Weight	Starting dose: 0.1 ml/kg twice daily	Maximum dose: 0.3 ml/kg twice daily
6 kg	0.6 ml twice daily	1.8 ml twice daily
8 kg	0.8 ml twice daily	2.4 ml twice daily
10 kg	1 ml twice daily	3 ml twice daily
15 kg	1.5 ml twice daily	4.5 ml twice daily
20 kg	2 ml twice daily	6 ml twice daily
25 kg	2.5 ml twice daily	7.5 ml twice daily
From 50 kg	5 ml twice daily	15 ml twice daily

Dose in infants (1 month to less than 6 months):

For infants 1 month to less than 6 months, measure the appropriate dosage using the **1 ml** syringe

included in the package.

General dose: Levetiracetam Clonmel is taken twice daily, in two equally divided doses, each individual dose being measured between 0.07 ml (7 mg) and 0.21 ml (21 mg), per kg bodyweight of the infant (see table below for dose examples).

Dose in infants (1 month to less than 6 months):


Weight	Starting dose: 0.07 ml/kg twice daily	Maximum dose: 0.21 ml/kg twice daily
4 kg	0.3 ml twice daily	0.85 ml twice daily
5 kg	0.35 ml twice daily	1.05 ml twice daily
6 kg	0.45 ml twice daily	1.25 ml twice daily
7 kg	0.5 ml twice daily	1.5 ml twice daily

Method of administration:

After measuring the correct dosage with an appropriate syringe, Levetiracetam Clonmel oral solution may be diluted in a glass of water or baby's bottle. You may take Levetiracetam Clonmel with or without food. After oral administration the bitter taste of levetiracetam may be experienced.

Instructions for use:

- Open the bottle: press the cap and turn it anticlockwise (figure 1)
- Separate the adaptor from the syringe (figure 2). Insert the adaptor into the bottle neck. Ensure it is well fixed.
- Take the syringe and put it in the adaptor opening. Turn the bottle upside down (figure 3)
- Fill the syringe with a small amount of solution by pulling the piston down (figure 4 a), then push the piston upward in order to remove any possible bubble (figure 4 b) Pull the piston down to the graduation mark corresponding to the quantity in millilitres (ml) prescribed by your doctor (figure 4 c).
- Turn the bottle the right way up. Remove the syringe from the adaptor.
- Empty the contents of the syringe in a glass of water or baby's bottle by pushing the piston to the bottom of the syringe (figure 5).
- Drink the whole contents of the glass/baby's bottle.
- Close the bottle with the child resistant closure.
- Wash the syringe with water only (figure 6)
-



Duration of treatment:

- Levetiracetam Clonmel is used as a chronic treatment. You should continue Levetiracetam Clonmel treatment for as long as your doctor has told you.
- Do not stop your treatment without your doctor's advice as this could increase your seizures.

If you take more Levetiracetam Clonmel than you should

The possible side effects of an overdose of Levetiracetam Clonmel are sleepiness, agitation, aggression, decrease of alertness, inhibition of breathing and coma.

Contact your doctor if you took more Levetiracetam Clonmel than you should.

Your doctor will establish the best possible treatment of overdose.

If you forget to take Levetiracetam Clonmel

Contact your doctor if you have missed one or more doses. Do not take a double dose to make up for a forgotten dose.

If you stop taking Levetiracetam Clonmel

If stopping treatment, Levetiracetam Clonmel should be discontinued gradually to avoid an increase of seizures. Should your doctor decide to stop your Levetiracetam Clonmel treatment, he/she will instruct you about the gradual withdrawal of Levetiracetam Clonmel.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Tell your doctor immediately, or go to your nearest emergency department, if you experience:

- weakness, feel light-headed or dizzy or have difficulty breathing, as these may be signs of a serious allergic (anaphylactic) reaction
- swelling of the face, lips, tongue and throat (Quincke's oedema)
- flu-like symptoms and a rash on the face followed by an extended rash with a high temperature, increased levels of liver enzymes seen in blood tests and an increase in a type of white blood cell (eosinophilia) and enlarged lymph nodes (Drug Reaction with Eosinophilia and Systemic Symptoms [DRESS])
- symptoms such as low urine volume, tiredness, nausea, vomiting, confusion and swelling in the legs, ankles or feet, as this may be a sign of sudden decrease of kidney function
- a skin rash which may form blisters and look like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge) (*erythema multiforme*)
- a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (*Stevens-Johnson syndrome*)
- a more severe form of rash causing skin peeling in more than 30 % of the body surface (*toxic epidermal necrolysis*)
- signs of serious mental changes or if someone around you notices signs of confusion, somnolence (sleepiness), amnesia (loss of memory), memory impairment (forgetfulness), abnormal behaviour or other neurological signs including involuntary or uncontrolled movements. These could be symptoms of an encephalopathy.

The most frequently reported adverse reactions were nasopharyngitis, somnolence (sleepiness), headache, fatigue and dizziness. At the beginning of the treatment or at dose increase side effects like sleepiness, tiredness and dizziness may be more common. These effects should however decrease over time.

Very common (may affect more than 1 in 10 people):

- nasopharyngitis;
- somnolence (sleepiness), headache.

Common (may affect up to 1 in 10 people):

- anorexia (loss of appetite);
- depression, hostility or aggression, anxiety, insomnia, nervousness or irritability;
- convulsion, balance disorder (equilibrium disorder), dizziness (sensation of unsteadiness), lethargy (lack of energy and enthusiasm), tremor (involuntary trembling);
- vertigo (sensation of rotation);
- cough;
- abdominal pain, diarrhoea, dyspepsia (indigestion), vomiting, nausea;
- rash;
- asthenia/fatigue (tiredness).

Uncommon (may affect up to 1 in 100 people):

- decreased number of blood platelets, decreased number of white blood cells;
- weight decrease, weight increase;
- suicide attempt and suicidal ideation, mental disorder, abnormal behaviour, hallucination, anger, confusion, panic attack, emotional instability/mood swings, agitation;
- amnesia (loss of memory), memory impairment (forgetfulness), abnormal coordination/ataxia (impaired coordinated movements), paraesthesia (tingling), disturbance in attention (loss of concentration);
- diplopia (double vision), vision blurred;
- elevated/abnormal values in a liver function test;

- hair loss, eczema, pruritus;
- muscle weakness, myalgia (muscle pain);
- injury.

Rare (may affect up to 1 in 1,000 people):

- infection;
- decreased number of all blood cell types;
- severe allergic reactions (DRESS, anaphylactic reaction [severe and important allergic reaction], Quincke's oedema [swelling of the face, lips, tongue and throat]);
- decreased blood sodium concentration;
- suicide, personality disorders (behavioural problems), thinking abnormal (slow thinking, unable to concentrate);
- delirium;
- encephalopathy (see sub-section "Tell your doctor immediately" for a detailed description of symptoms);
- seizures may become worse or happen more often;
- uncontrollable muscle spasms affecting the head, torso and limbs, difficulty in controlling movements, hyperkinesia (hyperactivity);
- change of the heart rhythm (Electrocardiogram);
- pancreatitis;
- liver failure, hepatitis;
- sudden decrease in kidney function;
- skin rash, which may form blisters and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge) (*erythema multiforme*), a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (*Stevens-Johnson syndrome*), and a more severe form causing skin peeling in more than 30 % of the body surface (*toxic epidermal necrolysis*).
- rhabdomyolysis (breakdown of muscle tissue) and associated blood creatine phosphokinase increase. Prevalence is significantly higher in Japanese patients when compared to non- Japanese patients.
- limp or difficulty walking

Reporting of side effects

If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via:

Ireland

HPRA Pharmacovigilance

Website: www.hpra.ie

Malta

ADR Reporting

Website: www.medicinesauthority.gov.mt/adrportal

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Levetiracetam Clonmel

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and bottle after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

After first opening: Do not store above 25 °C.

Do not use after 7 months from the first opening of the bottle.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to

throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Levetiracetam Clonmel contains

- The active substance is levetiracetam.
Each ml solution contains 100 mg of levetiracetam.
- The other ingredients are: sodium citrate (for pH adjustment), citric acid (for pH adjustment), methyl parahydroxybenzoate (E218), glycerol (E422), acesulfame potassium (E950), liquid maltitol (E965), raspberry liquid (containing benzyl alcohol).

What Levetiracetam Clonmel looks like and contents of the pack

Levetiracetam Clonmel 100 mg/ml oral solution is a clear liquid.

The 300 ml glass bottle of Levetiracetam Clonmel (for children aged 4 years and above, adolescents and adults) is packed in a cardboard box containing a 10 ml oral syringe (graduated every 0.25 ml) and an adaptor for the syringe.

The 150 ml glass bottle of Levetiracetam Clonmel (for infants and young children aged from 6 months to less than 4 years) is packed in a cardboard box containing a 3 ml oral syringe (graduated every 0.1 ml) and an adaptor for the syringe.

The 150 ml glass bottle of Levetiracetam Clonmel (for infants aged 1 month to less than 6 months) is packed in a cardboard box containing a 1 ml oral syringe (graduated every 0.05 ml) and an adaptor for the syringe.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Clonmel Healthcare Ltd, Waterford Road, Clonmel, Co. Tipperary, Ireland

Manufacturer

Remedica Ltd, Aharnon Street, Limassol Industrial Estate, Limassol 3056, Cyprus

STADA Arzneimittel AG, Stadastrasse 2-18, 61118 Bad Vilbel, Germany

Galenica Pharmaceutical Industry S.A., 3rd Km Old National Road Chalkida Athens, Chalcis, 341 00, Greece

This medicine is authorised in the Member States of the European Economic Area under the following names:

Germany:	Levetiracetam AL 100 mg/ml Lösung zum Einnehmen
Austria:	Levetiracetam STADA 100 mg/ml Lösung zum Einnehmen
Denmark:	Levetiracetam STADA
Finland:	Levetiracetam STADA 100 mg/ml oraaliliuos
France:	LEVETIRACETAM EG 100 mg/ml, solution buvable
Iceland:	Levetiracetam STADA 100 mg/ml mixtúra, lausn
Ireland:	Levetiracetam Clonmel 100 mg/ml oral solution
Italy:	LEVETIRACETAM EUROGENERICI
Malta:	Levetiracetam Clonmel 100 mg/ml oral solution
Poland:	Levetiracetam Stada
Portugal:	Levetiracetam Ciclum Farma
Spain:	Levetiracetam STADAGEN 100 mg/ml solución oral EFG
Sweden:	Levetiracetam STADA 100 mg/ml oral lösning

This leaflet was last revised in October 2025.