## Pomalidomide Prescription Authorisation Form (PAF)

A newly completed copy of this form MUST accompany EVERY pomalidomide prescription. Completion of this information is mandatory for ALL patients. The completed form should be retained in pharmacy.

Name of Treating Hospital:	Both signatures must be prese	nt prior to
Patient Date of Birth: DD MM YYYY Patient ID Number/Initials:	dispensing pomalidomide	
Prescriber: (print)	Prescriber's declaration	
Supervising Physician name: (print)	As the Prescriber, I have read and understood the Healthcare Professionals' Information Pack. I confirm the information provided on this PAF is	
Indication: (tick) Relapsed and Refractory Multiple Myeloma  Multiple Myeloma  If other please specify:	accurate, complete and in accordance with the requirements of the Pregnancy Prevention Programme for pomalidomide. I confirm treatment has been initiated and is monitored under the supervision of a physician with expertise in managing immunomodulatory or chemotherapeutic agents.	
Capsule strength prescribed: (tick)  Quantity of capsules per cycle prescribed*  Number of cycle(s) prescribed  *Do NOT enter number of packs	Sign	Print
Please tick all boxes that apply  Woman of non-childbearing potential  TICK	Date DD MM YYYY Bleep  Note to pharmacist – Prescription must be accompanied by a Prescription Authorisation Form	
Male	Pharmacist's declaration	
The patient has been counselled about the teratogenic risk of treatment with pomalidomide and understands the need to use a condom if involved in sexual activity with a woman of childbearing potential not using effective contraception or if their partner is pregnant (even if the patient has had a vasectomy).	I am satisfied that this Prescription Authorisation Form has been completed fully and that I have read and understood the Healthcare Professionals' Information Pack. For women of childbearing potential, dispensing will be taking place within 7 days of the date of prescription. I am dispensing no more than a 4-week supply to women of childbearing potential and 12-	
Note to pharmacist – do not dispense unless ticked and, for a male, Y selected	weeks for males and women of non-childbearing potential.	
Woman of childbearing potential	Sign	Print
The patient has been counselled about the teratogenic risk of treatment, the need to avoid pregnancy, and has been on effective contraception for at least 4 weeks or committed to absolute and continuous abstinence confirmed on a monthly basis.	Date DD MM YYYY Bleep  Name and postcode of dispensing pharmacy	
Date of last negative pregnancy test DD MM YYYY	disperising priarriacy	
Note to pharmacist – do not dispense unless ticked, Y selected for counselling and a negative test has been conducted within 3 days prior of the prescription date and dispensing is taking place within 7 days of the prescription date.	Product brand you have dispensed	