

Package leaflet: information for the user

Macitentan Clonmel 10 mg film-coated tablets macitentan

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Macitentan Clonmel is and what it is used for
2. What you need to know before you take Macitentan Clonmel
3. How to take Macitentan Clonmel
4. Possible side effects
5. How to store Macitentan Clonmel
6. Contents of the pack and other information

1. What Macitentan Clonmel is and what it is used for

Macitentan Clonmel contains the active substance macitentan, which belongs to the class of medicines called “endothelin receptor antagonists”.

Macitentan Clonmel is used for the long-term treatment of pulmonary arterial hypertension (PAH):

- in adults of WHO Functional Class (FC) II to III
- in children under 18 years and body weight of at least 40 kg with WHO Functional Class (FC) II to III

It can be used on its own or with other medicines for PAH. PAH is high blood pressure in the blood vessels that carry blood from the heart to the lungs (the pulmonary arteries). In people with PAH, these arteries get narrower, so the heart has to work harder to pump blood through them. This causes people to feel tired, dizzy, and short of breath.

Macitentan Clonmel widens the pulmonary arteries, making it easier for the heart to pump blood through them. This lowers the blood pressure, relieves the symptoms and improves the course of the disease.

2. What you need to know before you take Macitentan Clonmel

DO NOT take Macitentan Clonmel

- if you are allergic to macitentan or any of the other ingredients of this medicine (listed in section 6)
- if you are pregnant, if you are planning to become pregnant, or if you could become pregnant because you are not using reliable birth control (contraception). See section ‘Pregnancy and breast-feeding’
- if you are breast-feeding. See section ‘Pregnancy and breast-feeding’
- if you have liver disease or if you have very high levels of liver enzymes in your blood. Talk to your doctor, who will decide whether this medicine is suitable for you

If any of these apply to you, please tell your doctor.

Warnings and precautions

Talk to your doctor or pharmacist before taking Macitentan Clonmel.

You will need blood tests, as indicated by your doctor:

Your doctor will take blood test before you start treatment with Macitentan Clonmel and during treatment to test:

- whether you have anaemia (a reduced number of red blood cells)
- whether your liver is working properly

If you have anaemia (a reduced number of red blood cells), you may have the following signs:

- dizziness
- fatigue/malaise/weakness
- fast heart rate, palpitations
- pallor

If you notice any of these signs, **tell your doctor**.

Signs that your liver may not be working properly include:

- feeling sick (nausea)
- vomiting
- fever
- pain in your stomach (abdomen)
- yellowing of your skin or the whites of your eyes (jaundice)
- dark-coloured urine
- itching of your skin
- unusual tiredness or exhaustion (lethargy or fatigue)
- flu-like syndrome (joint and muscle pain with fever)

If you notice any of these signs, **tell your doctor immediately**.

If you have kidney problems, talk to your doctor before using Macitentan Clonmel. Macitentan may lead to more reduction of blood pressure and decrease in haemoglobin in patients with kidney problems.

In patients with pulmonary veno-occlusive disease (obstruction of the lung veins), the use of medicines for treatment of PAH, including Macitentan Clonmel, may lead to pulmonary oedema. If you have signs of pulmonary oedema when using Macitentan Clonmel, such as a sudden, important increase in breathlessness and low oxygen, **tell your doctor immediately**. Your doctor may perform additional tests, and will determine what treatment regimen is most suitable for you.

Children and adolescents

Do not give this medicine to children below 2 years of age because efficacy and safety have not been established.

Other medicines and Macitentan Clonmel

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicine. Macitentan Clonmel can affect other medicines.

If you take Macitentan Clonmel together with other medicines including those listed below, the effects of Macitentan Clonmel or the other medicines might be altered. Please talk to your doctor or pharmacist if you are taking any of the following medicines:

- rifampicin, clarithromycin, telithromycin, ciprofloxacin, erythromycin (antibiotics used to treat infections)
- phenytoin (a medicine used to treat seizures)

- carbamazepine (used to treat depression and epilepsy)
- St. John's Wort (an herbal preparation used to treat depression)
- ritonavir, saquinavir (used to treat HIV infections)
- nefazodone (used to treat depression)
- ketoconazole (except shampoo), fluconazole, itraconazole, miconazole, voriconazole (medicines used against fungal infections)
- amiodarone (to control the heartbeat)
- cyclosporine (used to prevent organ rejection after transplant)
- diltiazem, verapamil (to treat high blood pressure or specific heart problems)

Macitentan Clonmel with food

If you are taking piperine as a dietary supplement, this may alter how the body responds to some medicinal products, including Macitentan Clonmel. Please talk to your doctor or pharmacist should this be the case.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

Macitentan Clonmel may harm unborn babies conceived before, during or soon after treatment.

- If it is possible you could become pregnant, use a reliable form of birth control (contraception) while you are taking Macitentan Clonmel. Talk to your doctor about this.
- Do not take Macitentan Clonmel if you are pregnant or planning to become pregnant.
- If you become pregnant or think that you may be pregnant while you are taking Macitentan Clonmel, or shortly after stopping Macitentan Clonmel (up to 1 month), see your doctor immediately.

If you are a woman who could become pregnant, your doctor will ask you to take a pregnancy test before you start taking Macitentan Clonmel and regularly (once a month) while you are taking Macitentan Clonmel.

It is not known if macitentan is transferred to breast milk. Do not breast-feed while you are taking Macitentan Clonmel. Talk to your doctor about this.

Fertility

If you are a man taking Macitentan Clonmel, it is possible that this medicine may lower your sperm count. Talk to your doctor if you have any questions or concerns about this.

Driving and using machines

Macitentan Clonmel can cause side effects such as headaches and hypotension (listed in section 4), and the symptoms of your condition can also make you less fit to drive or use machines.

Macitentan Clonmel contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Macitentan Clonmel

Macitentan Clonmel should only be prescribed by a doctor experienced in the treatment of pulmonary arterial hypertension.

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

Adults and children aged less than 18 years weighing at least 40 kg

The recommended dose of Macitentan Clonmel is one 10 mg tablet, once a day. Swallow the whole tablet, with a glass of water. Do not chew or break the tablet in order to avoid damage to the film-coating. You can take Macitentan Clonmel with or without food. It is best to take the tablet at the same time each day.

For children weighing less than 40 kg, macitentan dispersible tablets with a lower strength are available under other brand names.

If you take more Macitentan Clonmel than you should

If you have taken more tablets than you have been told to take, you may experience headache, nausea, or vomiting. Ask your doctor for advice.

If you forget to take Macitentan Clonmel

If you forget to take Macitentan Clonmel, take a dose as soon as you remember, then continue to take your tablets at the usual times. Do not take a double dose to make up for a forgotten tablet.

If you stop taking Macitentan Clonmel

Macitentan Clonmel is a treatment that you will need to keep on taking to control your PAH. Do not stop taking Macitentan Clonmel unless you have agreed this with your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Uncommon serious side effects (may affect up to 1 in 100 people)

- Allergic reactions (swelling around the eyes, face, lips, tongue or throat, itching and/or rash)

If you notice any of these signs, tell your doctor immediately.

Very common side effects (may affect more than 1 in 10 people)

- Anaemia (low number of red blood cells) or reduced haemoglobin
- Headache
- Bronchitis (inflammation of the airways)
- Nasopharyngitis (inflammation of the throat and nasal passages)
- Oedema (swelling), especially of the ankles and feet

Common side effects (may affect up to 1 in 10 people)

- Pharyngitis (inflammation of the throat)
- Influenza (flu)
- Urinary tract infection (bladder infection)
- Hypotension (low blood pressure)
- Nasal congestion (blocked nose)
- Elevated liver tests
- Leukopenia (decreased white blood cell counts)
- Thrombocytopenia (decreased blood platelet counts)
- Flushing (redness of the skin)
- Increased uterine bleeding

Side effects in children and adolescents

The side effects listed above may also be seen in children. Additional side effects very commonly seen in children include upper respiratory tract infection (infected nose sinuses, or

throat) and gastroenteritis (inflamed stomach and gut). Rhinitis (itchy, runny, or blocked nose) was seen commonly in children.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via:

Ireland: HPRA Pharmacovigilance, Website: www.hpra.ie

Malta: ADR Reporting, Website: www.medicinesauthority.gov.mt/adrportal

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Macitentan Clonmel

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer require. These measures will help protect the environment.

6. Contents of the pack and other information

What Macitentan Clonmel contains

The active substance is macitentan. Each film-coated tablet contains 10 mg macitentan.

The other ingredients are maltodextrin (E1400), polysorbate 80 (E433), mannitol (E421), starch, pregelatinised (maize starch), croscarmellose sodium (E468), magnesium stearate (E572), poly(vinyl alcohol) (E1203), talc (E553b), titanium dioxide (E171), glycerol monocaprylocaprate, type I, sodium laurilsulfate (E487).

What Macitentan Clonmel looks like and contents of the pack

Macitentan Clonmel is a film-coated tablet.

White to off-white, round, biconvex film-coated tablets with a diameter of approximately 5.6 mm.

Macitentan Clonmel 10 mg is available in PVC/PE/PVdC-Al blisters containing 15 or 30 film-coated tablets or PVC/PE/PVdC-Al unit-dose blisters containing 15x1 or 30x1 film-coated tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Clonmel Healthcare Ltd, Waterford Road, Clonmel, Co. Tipperary, Ireland

Manufacturer

STADA Arzneimittel AG, Stadastrasse 2-18, 61118 Bad Vilbel, Germany

Clonmel Healthcare Ltd, Waterford Road, Clonmel, Co. Tipperary, Ireland

STADA Arzneimittel GmbH, Muthgasse 36/2, 1190 Wien, Austria

Centrafarm Services B.V., Van de Reijtstraat 31 E, Breda, 4814NE, Netherlands

This medicine is authorised in the Member States of the European Economic Area under the following names:

Austria: Macitenant STADA 10 mg Filmtabletten

Belgium:	Macitentan EG 10 mg filmomhulde tabletten
Denmark:	Macitentan STADA
Finland:	Macitentan STADA 10 mg kalvopäällysteiset tabletit
Germany:	Macitentan STADA 10 mg Filmtabletten
Hungary:	Macitentan Stada 10 mg filmtabletta
Iceland:	Macitentan STADA 10 mg filmuhúðaðar töflur
Ireland:	Macitentan Clonmel 10 mg film-coated tablets
Italy:	Macitentan EG
Luxembourg:	Macitentan EG 10 mg comprimés pelliculés
Malta:	Macitentan Clonmel 10 mg film-coated tablets
Netherlands:	Macitentan CF 10 mg, filmomhulde tabletten
Norway:	Macitentan STADA
Poland:	Macitentan Stada
Portugal:	Macitentano Stada
Romania:	Macitentan Stada 10 mg comprimate filmate
Slovakia:	Macitentan STADA 10 mg filmom obalené tablety
Spain:	Macitentan STADA 10 mg comprimidos recubiertos con película EFG
Sweden:	Macitentan STADA

This leaflet was last revised in June 2025.