

**B. PACKAGE LEAFLET**

## Package leaflet: Information for the user

### Linagliptin Accord 5 mg film-coated tablets

linagliptin

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

1. What Linagliptin Accord is and what it is used for
2. What you need to know before you take Linagliptin Accord
3. How to take Linagliptin Accord
4. Possible side effects
5. How to store Linagliptin Accord
6. Contents of the pack and other information.

#### **1. What Linagliptin Accord is and what it is used for**

Linagliptin Accord contains the active substance linagliptin which belongs to a group of medicines called “oral anti-diabetics”. Oral anti-diabetics are used to treat high blood sugar levels. They work by helping the body reduce the level of sugar in your blood.

Linagliptin Accord is used for ‘type 2 diabetes’ in adults, if the disease cannot be adequately controlled with one oral anti-diabetic medicine (metformin or sulphonylureas) or diet and exercise alone. Linagliptin Accord may be used together with other anti-diabetic medicines e.g. metformin, sulphonylureas (e.g. glimepiride, glipizide), empagliflozin, or insulin.

It is important to keep following the advice about diet and exercise that you have been given by your doctor or nurse.

#### **2. What you need to know before you take Linagliptin Accord**

##### **Do not take Linagliptin Accord**

- if you are allergic to linagliptin or any of the other ingredients of this medicine (listed in section 6).

##### **Warnings and precautions**

Talk to your doctor, pharmacist or nurse before taking Linagliptin Accord if you:

- have type 1 diabetes (your body does not produce any insulin) or diabetic ketoacidosis (a complication of diabetes with high blood sugar, rapid weight loss, nausea or vomiting). Linagliptin Accord should not be used to treat these conditions.
- are taking an anti-diabetic medicine known as a ‘sulphonylurea’ (e.g. glimepiride, glipizide), your doctor may want to reduce your dose of sulphonylurea when you take it together with Linagliptin Accord in order to avoid your blood sugar going too low.
- have had allergic reactions to any other medicines that you take to control the amount of sugar in your blood.

- have or have had a disease of the pancreas.

If you have symptoms of acute pancreatitis, like persistent, severe stomach ache (abdominal pain), you should consult your doctor.

If you encounter blistering of the skin it may be a sign for a condition called bullous pemphigoid. Your doctor may ask you to stop Linagliptin Accord.

Diabetic skin lesions are a common complication of diabetes. You are advised to follow the recommendations for skin and foot care that you are given by your doctor or nurse.

### **Children and adolescents**

Linagliptin Accord is not recommended for children and adolescents under 18 years. It is not effective in children and adolescents between the ages of 10 and 17 years. It is not known if this medicine is safe and effective when used in children younger than 10 years.

### **Other medicines and Linagliptin Accord**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

In particular, you should tell your doctor if you are using medicines containing any of the following active substances:

- Carbamazepine, phenobarbital or phenytoin. These may be used to control fits (seizures) or chronic pain.
- Rifampicin. This is an antibiotic used to treat infections such as tuberculosis.

### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

It is unknown if Linagliptin Accord is harmful to the unborn child. Therefore, it is preferable to avoid using Linagliptin Accord if you are pregnant.

It is not known if Linagliptin Accord passes into human breast milk. A decision must be made by your doctor whether to discontinue breast-feeding or to discontinue/abstain from Linagliptin Accord therapy.

### **Driving and using machines**

Linagliptin Accord has no or negligible influence on the ability to drive and use machines.

Taking Linagliptin Accord in combination with medicines called sulphonylureas and/or insulin can cause too low blood sugar levels (hypoglycaemia), which may affect your ability to drive and use machines or work without safe foothold. However, more frequent blood glucose testing might be recommended to minimise the risk for hypoglycaemia, especially when Linagliptin Accord is combined with sulphonylurea and/or insulin.

## **3. How to take Linagliptin Accord**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose of Linagliptin Accord is one 5 mg tablet once a day.

You can take Linagliptin Accord with or without food.

Your doctor may prescribe Linagliptin Accord together with another oral anti-diabetic medicine. Remember to take all medicines as directed by your doctor to achieve the best results for your health.

### **If you take more Linagliptin Accord than you should**

If you take more Linagliptin Accord than you should, talk to a doctor immediately.

### **If you forget to take Linagliptin Accord**

- If you forget to take a dose of Linagliptin Accord, take it as soon as you remember it. However, if it is nearly time for the next dose, skip the missed dose.
- Do not take a double dose to make up for a forgotten dose. Never take two doses on the same day.

### **If you stop taking Linagliptin Accord**

Do not stop taking Linagliptin Accord without first consulting your doctor. Your blood sugar levels may increase when you stop taking Linagliptin Accord.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

### Some symptoms need immediate medical attention

You should stop taking Linagliptin Accord and see your doctor immediately if you experience the following symptoms of low blood sugar: trembling, sweating, anxiety, blurred vision, tingling lips, paleness, mood change or confusion (hypoglycaemia). Hypoglycaemia (frequency: very common, may affect more than 1 in 10 people) is an identified side effect when Linagliptin Accord is taken together with metformin and a sulphonylurea.

Some patients have experienced allergic reactions (hypersensitivity; frequency uncommon, may affect up to 1 in 100 people) while taking Linagliptin Accord alone or in combination with other medicinal products for the treatment of diabetes, which may be serious, including wheezing and shortness of breath (bronchial hyperreactivity; frequency not known, frequency cannot be estimated from the available data). Some patients experienced rash (frequency uncommon), hives (urticaria; frequency rare, may affect up to 1 in 1000 people), and swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing (angioedema; frequency rare). If you experience any of the signs of illness mentioned above, stop taking Linagliptin Accord and call your doctor right away. Your doctor may prescribe a medicine to treat your allergic reaction and a different medicine for your diabetes.

Some patients have experienced inflammation of the pancreas (pancreatitis; frequency rare, may affect up to 1 in 1000 people) while taking Linagliptin Accord alone or in combination with other medicinal products for the treatment of diabetes.

STOP taking Linagliptin Accord and contact a doctor immediately if you notice any of the following serious side effects:

- Severe and persistent pain in the abdomen (stomach area) which might reach through to your back, as well as nausea and vomiting, as it could be a sign of an inflamed pancreas (pancreatitis).

Some patients have had the following side effects while taking Linagliptin Accord alone or in combination with other medicinal products for the treatment of diabetes:

- Common: level of lipase in the blood increased.

- Uncommon: inflamed nose or throat (nasopharyngitis), cough, constipation (in combination with insulin), level of amylase in the blood increased.
- Rare: blistering of skin (bullous pemphigoid).

### **Reporting of side effects**

If you get any side effects talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via

HPRA Pharmacovigilance

Website: [www.hpra.ie](http://www.hpra.ie)

By reporting side effects you can help provide more information on the safety of this medicine.

### **5. How to store Linagliptin Accord**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and the carton after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not use Linagliptin Accord if the package is damaged or shows signs of tampering.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

### **6. Contents of the pack and other information**

#### **What Linagliptin Accord contains**

- The active substance is linagliptin  
Each film-coated tablet contains 5 mg of linagliptin
- The other ingredients are  
Tablet core: Mannitol (E421), maize starch, copovidone, magnesium stearate (E470b)  
Film coating: Hypromellose (E464), titanium dioxide (E171), talc (E553b), macrogol (6000), iron oxide red (E172)

#### **What Linagliptin Accord looks like and contents of the pack**

- Linagliptin Accord 5 mg tablets are 8 mm diameter round, light red, biconvex, bevel-edged film coated tablet debossed with "L5" on one side.
- Linagliptin Accord is available in OPA/Al/PVC//Al unit dose blisters containing, 10 x 1, 14 x 1, 28 x 1, 30 x 1, 56 x 1, 60 x 1, 84 x 1, 90 x 1, 98 x 1, 100 x 1, 120 x 1 film-coated tablets.

Not all pack sizes may be marketed in your country.

#### **Marketing Authorisation Holder and Manufacturer**

##### Marketing Authorisation Holder:

Accord Healthcare Ireland Ltd  
Euro House  
Euro Business Park

Little Island  
Cork T45 K857  
Ireland

Manufacturer:

Accord Healthcare Polska Sp.z.o.o.  
ul. Lutomierska 50,  
95-200, Pabianice,  
Poland

Accord Healthcare B.V.  
Winthontlaan 200,  
3526 KV Utrecht, Netherlands

Pharmadox Healthcare Limited  
KW20A Kordin Industrial Park,  
PLA 3000, Paola  
Malta

**This medicine is authorised in the Member States of the EEA under the following names:**

<b>Name of the member state</b>	<b>Name of the medicinal product</b>
Denmark	Linagliptin Accord
Finland	Linagliptin Accord 5 mg kalvopäällysteiset tabletit
Latvia	Linagliptin Accord 5 mg apvalkotās tabletes
Norway	Linagliptin Accord
Netherlands	Linagliptine Accord 5 mg, filmomhulde tabletten
Sweden	Linagliptin Accord
Estonia	Linagliptin Accord
Austria	Linagliptin Accord 5 mg Filmtabletten
Bulgaria	Линаглиптин Акорд 5 mg филмирани таблетки
Czechia	Linagliptin Accord
Poland	Linagliptin Accord
Ireland	Linagliptin Accord 5 mg film-coated tablets
Greece	Linagliptin/Accord
Italy	Linagliptin Accord
Cyprus	Linagliptin Accord 5 mg επικαλυμμένα με λεπτό υμένιο δισκία

**This leaflet was last revised in 12/2025**