

Package leaflet: Information for the patient

Anastrozole Amarox 1 mg film-coated tablets anastrozole

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, or pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Anastrozole Amarox is and what it is used for
2. What you need to know before you take Anastrozole Amarox
3. How to take Anastrozole Amarox
4. Possible side effects
5. How to store Anastrozole Amarox
6. Contents of the pack and other information

1. What Anastrozole Amarox is and what it is used for

Anastrozole Amarox contains a substance called anastrozole. This belongs to a group of medicines called ‘aromatase inhibitors’. Anastrozole is used to treat breast cancer in women who have gone through the menopause.

Anastrozole works by cutting down the amount of the hormone called estrogen that your body makes. It does this by blocking a natural substance (an enzyme) in your body called ‘aromatase’.

2. What you need to know before you take Anastrozole Amarox

Do not take Anastrozole Amarox

- if you are allergic to anastrozole or any of the other ingredients of this medicine (listed in section 6).
- if you are pregnant or breast-feeding (see the section called ‘Pregnancy and breast-feeding’).

Do not take Anastrozole Amarox if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Anastrozole Amarox.

Warnings and precautions

Talk to your doctor, or pharmacist or nurse before taking Anastrozole Amarox

- if you still have menstrual periods and have not yet gone through the menopause.

- if you are taking a medicine that contains tamoxifen or medicines that contain estrogen (see the section called ‘Other medicines and Anastrozole AmaroX ’).
- if you have ever had a condition that affects the strength of your bones (osteoporosis).
- if you have problems with your liver or kidneys.

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking Anastrozole AmaroX.

If you go into the hospital, let the medical staff know you are taking Anastrozole AmaroX.

Other medicines and Anastrozole AmaroX

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes medicines that you buy without a prescription and herbal medicines. This is because Anastrozole AmaroX can affect the way some medicines work and some medicines can have an effect on Anastrozole AmaroX.

Do not take Anastrozole AmaroX if you are already taking any of the following medicines:

- Certain medicines used to treat breast cancer (selective estrogen receptor modulators), e.g. medicines that contain tamoxifen. This is because these medicines may stop Anastrozole AmaroX from working properly.
- Medicines that contain estrogen, such as hormone replacement therapy (HRT).

If this applies to you, ask your doctor or pharmacist for advice.

Tell your doctor or pharmacist if you are taking the following:

- A medicine known as an ‘LHRH analogue’. This includes gonadorelin, buserelin, goserelin, leuprorelin and triptorelin. These medicines are used to treat breast cancer, certain female health (gynaecological) conditions, and infertility.

Pregnancy and breast-feeding

Do not take Anastrozole AmaroX if you are pregnant or breast-feeding. Stop Anastrozole AmaroX if you become pregnant and talk to your doctor.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Anastrozole AmaroX is not likely to affect your ability to drive or use any tools or machines. However, some people may occasionally feel weak or sleepy while taking Anastrozole AmaroX. If this happens to you, ask your doctor or pharmacist for advice.

Anastrozole AmaroX contains lactose

Anastrozole AmaroX contains lactose which is a type of sugar. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

Anastrozole AmaroX contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially ‘sodium-free’.

3. How to take Anastrozole Amarox

Always take Anastrozole Amarox exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

- The recommended dose is one tablet once a day.
- Try to take your tablet at the same time each day.
- Swallow the tablet whole with a drink of water.
- It does not matter if you take Anastrozole Amarox before, with or after food.

Keep taking Anastrozole Amarox for as long as your doctor or pharmacist tells you to. It is a long-term treatment and you may need to take it for several years. Check with your doctor or pharmacist if you are not sure.

Use in children and adolescents

Anastrozole Amarox should not be given to children and adolescents.

If you take more Anastrozole Amarox than you should

If you take more Anastrozole Amarox than you should, talk to a doctor straight away.

If you forget to take Anastrozole Amarox

If you forget to take a dose, just take your next dose as normal.

Do not take a double dose (two doses at the same time) to make up for a forgotten dose.

If you stop taking Anastrozole Amarox

Do not stop taking your tablets unless your doctor tells you to.

If you have any further questions on the use of this medicine, ask your doctor, or pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop taking Anastrozole Amarox and seek urgent medical treatment, if you experience any of the following serious but very rare side effects:

- An extremely severe skin reaction with ulcers or blisters on the skin. This is known as ‘Stevens-Johnson syndrome’.
- Allergic (hypersensitivity) reactions with swelling of the throat that may cause difficulty in swallowing or breathing. This is known as ‘angioedema’.

Very common side effects (affect more than 1 in 10 people)

- Headache.
- Hot flushes.
- Feeling sick (nausea).
- Skin rash.
- Pain or stiffness in your joints.

- Inflammation of the joints (arthritis).
- Feeling weak.
- Bone loss (osteoporosis).
- Depression

Common side effects (affect 1 to 10 people in 100)

- Loss of appetite.
- Raised or high levels of a fatty substance known as cholesterol in your blood. This would be seen in a blood test.
- Feeling sleepy.
- Carpal tunnel syndrome (tingling, pain, coldness, weakness in parts of the hand).
- Tickling, tingling or numbness of skin, loss/lack of taste
- Diarrhoea.
- Being sick (vomiting).
- Changes in blood tests that show how well your liver is working.
- Thinning of your hair (hair loss).
- Allergic (hypersensitivity) reactions including face, lips, or tongue.
- Bone pain.
- Vaginal dryness.
- Bleeding from the vagina (usually in the first few weeks of treatment – if the bleeding continues, talk to your doctor).
- Muscle pain.

Uncommon side effects (affect 1 to 10 people in 1,000)

- Changes in special blood tests that show how your liver is working (gamma-GT and bilirubin).
- Inflammation of the liver (hepatitis).
- Hives or nettle rash.
- Trigger finger (a condition in which your finger or thumb catches in a bent position).
- Increased amounts of calcium in your blood. If you experience nausea, vomiting and thirst, you should tell your doctor, or pharmacist or nurse as you may need to have blood tests.

Rare side effects (affect 1 to 10 people in 10,000)

- Rare inflammation of your skin that may include red patches or blisters.
- Skin rash caused by hypersensitivity (this can be from allergic or anaphylactoid reaction).
- Inflammation of the small blood vessels causing red or purple colouring of the skin. Very rarely symptoms of joint, stomach, and kidney pain may occur; this is known as ‘Henoch-Schönlein purpura’.

Side effects with frequency not known (frequency cannot be estimated from the available data)

- Dry eye
- Lichenoid eruption (small, red or purple itchy bumps on the skin)
- Inflammation of a tendon or tendonitis (connective tissues that connect muscles to bones)
- Rupture of a tendon (connective tissues that connect muscles to bones)
- Memory impairment

Effects on your bones

Anastrozole AmaroX lowers the amount of the hormone called estrogen that is in your body. This may lower the mineral content of your bones. Your bones may be less strong and may be more likely to fracture. Your doctor will manage these risks according to treatment guidelines for managing bone health in women who have gone through the menopause. You should talk to your doctor about the risks and treatment options.

If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance Website: www.hpra.ie

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Anastrozole AmaroX

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the outer carton and blister after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater <or household waste>. Ask your pharmacist how to throw away of medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Anastrozole AmaroX contains

The active substance is anastrozole.

Each film-coated tablet contains 1 mg of anastrozole.

The other ingredients are:

Tablet core: Lactose monohydrate, povidone, Sodium starch glycolate (Type A) and magnesium stearate

Film coat: Hypromellose (E 464), macrogol and titanium dioxide (E 171).

What Anastrozole AmaroX looks like and contents of the pack

White, round, biconvex (6.0 mm in length and 6.0 mm in width), film-coated tablet, debossed with "1" on one side and "H" on the other side.

Anastrozole AmaroX is presented in blisters containing 28 film-coated tablets.

Marketing Authorisation Holder

Amarox Pharma B.V.

Roubloslaan 32
2252 TR Voorschoten
Netherlands

Manufacturer

Amarox Pharma B.V.
Roubloslaan 32
2252 TR Voorschoten
Netherlands

This leaflet was last revised in