

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Colistimethate sodium Accord 2 million IU powder for solution for injection/infusion

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial contains 2 million International Units (IU) which is approximately equivalent to 160 mg of colistimethate sodium.

3 PHARMACEUTICAL FORM

Powder for solution for injection/infusion

White lyophilised powder.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Colistimethate sodium Accord is indicated in adults and children including neonates for the treatment of serious infections due to selected aerobic Gram-negative pathogens in patients with limited treatment options (see sections 4.2, 4.4, 4.8 and 5.1).

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

The dose to be administered and the treatment duration should take into account the severity of the infection as well as the clinical response. Therapeutic guidelines should be adhered to.

The dose is expressed in IU of colistimethate sodium (CMS). A conversion table from CMS in IU to mg of CMS as well as to mg of colistin base activity (CBA) is included at the end of this section.

Posology

The following dose recommendation are made based on limited population-pharmacokinetics data in critically ill patients (see section 4.4).

Adults and adolescents

Maintenance dose 9 million IU/day in 2-3 divided doses

In patients who are critically ill, a loading dose of 9 MIU should be administered.

The most appropriate time interval to the first maintenance dose has not been established.

Modelling suggests that loading and maintenance doses of up to 12 MIU may be required in patients with good renal function in some cases. Clinical experience with such doses is however extremely limited, and safety has not been established.

The loading dose applies to patients with normal and impaired renal functions including those on renal replacement therapy.

Renal impairment

Dose adjustments in renal impairment are necessary, but pharmacokinetic data available for patients with impaired renal function is very limited.

The following dose adjustments are suggested as guidance.

Dose reductions are recommended for patients with creatinine clearance < 50 ml/min:

Twice daily dosing is recommended.

Creatinine clearance (ml/min)	Daily dose
< 50- 30	5.5- 7.5 MIU
<30- 10	4.5- 5.5 MIU
<10	3.5 MIU

MIU = million IU

Haemodialysis and continuous haemo(dia)filtration

Colistin appears to be dialyzable through conventional haemodialysis and continuous venovenous haemo(dia)filtration (CVVHF, CVVHDF). There are extremely limited data from population PK studies from very small numbers of patients on renal replacement therapy. Firm dose recommendations cannot be made. The following regimes could be considered.

Haemodialysis

No-HD days: 2.25 MIU/day (2.2-2.3 MIU/day).

HD days: 3 MIU/day on haemodialysis days, to be given after the HD session.

Twice daily dosing is recommended.

CVVHF/ CVVHDF

As in patients with normal renal function. Three times daily dosing is recommended.

Hepatic impairment

There are no data in patients with hepatic impairment. Caution is advised when administering colistimethate sodium in these patients.

Elderly

No dose adjustments in older patients with normal renal function are considered necessary.

Paediatric population

The data supporting the dose regimen in paediatric patients are very limited. Renal maturity should be taken into consideration when selecting the dose. The dose should be based on lean body weight.

Children $\leq$ 40 kg

75,000-150,000 IU/kg/day divided into 3 doses.

For children with a body weight above 40 kg, use of the dosing recommendation for adults should be considered.

The use of doses >150,000 IU/kg/day has been reported in children with cystic fibrosis.

There are no data regarding the use or magnitude of a loading dose in critically ill children.

No dose recommendations have been established in children with impaired renal function.

Intrathecal and Intracerebroventricular administration

Based on limited data, the following dose is recommended in adults:

Intracerebroventricular route

125,000 IU/day

Intrathecally administered doses should not exceed those recommended for Intracerebroventricular use.

No specific dosing recommendation can be made in children for intrathecal and Intracerebroventricular routes of administration.

Method of administration

Colistimethate sodium Accord is administered intravenously as a slow infusion over 30 – 60 minutes.

Patients with a totally implantable venous access device (TIVAD) in place may tolerate a bolus injection of up to 2 million units in 10 ml given over a minimum of 5 minutes (see section 6.6).

Colistimethate sodium undergoes hydrolysis to the active substance colistin in aqueous solution. For dose preparation, particularly where combination of multiple vials is needed, reconstitution of the required dose must be performed using strict aseptic technique (see section 6.6).

Dose conversion table:

In the EU, the dose of colistimethate sodium (CMS) must be prescribed and administered only as IU. The product label states the number of IU per vial.

Confusion and medication errors have occurred because of the different expressions of dose in terms of potency. The dose is expressed in the US, and other parts of the world, as milligrams of colistin base activity (mg CBA).

The following conversion table is prepared for information and the values must be considered nominal and approximate only.

CMS conversion table

Potency		≈ mass of CMS (mg) *
IU	≈ mg CBA	
12 500	0.4	1
150 000	5	12
1 000 000	34	80
4 500 000	150	360
9 000 000	300	720

*Nominal potency of the drug substance = 12500 IU/mg

4.3 Contraindications

Hypersensitivity to the active substance, colistin or to other polymyxins.

4.4 Special warnings and precautions for use

Consideration should be given to co-administering intravenous colistimethate sodium with another antibacterial agent whenever this is possible, taking into account the remaining susceptibilities of the pathogen(s) under treatment. As the development of resistance to intravenous colistin has been reported in particular when it is used as a monotherapy, co-administration with other antibacterial should also be considered in order to prevent the emergence of resistance.

There are limited clinical data on the efficacy and safety of intravenous colistimethate sodium. The recommended doses in all subpopulations are equally based on limited data (clinical and pharmacokinetic/ pharmacodynamics data). In particular there are limited safety data for the use of high doses (> 6 MIU/day) and the use of a loading dose, and for special populations (patients with renal impairment and the paediatric population). Colistimethate sodium should only be used when other, more commonly prescribed antibiotics are not effective or not appropriate.

Renal function monitoring should be performed at the start of treatment and regularly during treatment in all patients. The dose of colistimethate sodium should be adjusted according to creatinine clearance (see section 4.2). Patients who are hypovolaemic or those receiving other potentially nephrotoxic drugs are at increased risk of nephrotoxicity from colistin (see sections 4.5 and 4.8).

Nephrotoxicity has been reported to be associated with cumulative dose and treatment duration in some studies. The benefit of prolonged treatment duration should be balanced against the potentially increased risk of renal toxicity.

Paediatric population

Caution is advised when administering colistimethate sodium to infants < 1 year of age as renal function is not fully mature in this age group. Further, the effect of immature renal and metabolic function on the conversion of colistimethate sodium to colistin is not known.

In case of an allergic reaction, treatment with colistimethate sodium must be discontinued and appropriate measures implemented.

High serum concentrations of colistimethate sodium, which may be associated with overdosage or failure to reduce the dosage in patients with renal impairment, have been reported to lead to neurotoxic effects such as facial paraesthesia, muscle weakness, vertigo, slurred speech, vasomotor instability, visual disturbances, confusion, psychosis and apnoea. Monitoring should be performed for perioral paraesthesia and paraesthesia in the extremities, which are signs of overdose (see section 4.9).

Colistimethate sodium is known to reduce the presynaptic release of acetyl-choline at the neuro-muscular junction and should be used in patients with myasthenia gravis with the greatest caution and only if clearly needed.

Respiratory arrest has been reported following intramuscular administration of colistimethate sodium. Impaired renal function increases the possibility of apnoea and neuromuscular blockade following administration of colistimethate sodium.

Colistimethate sodium should be used with extreme caution in patients with porphyria.

Antibiotic-associated colitis and pseudomembranous colitis have been reported with nearly all anti-bacterial agents and may occur with colistimethate sodium. They may range from mild to life-threatening in severity. It is important to consider this diagnosis in patients who develop diarrhoea during or after the use of colistimethate sodium (see section 4.8). Discontinuation of therapy and the administration of specific treatment for *Clostridioides difficile* should be considered. Medicinal products that inhibit peristalsis should not be given.

Few cases of pseudo-Bartter syndrome have been reported in children and adults with the intravenous use of colistimethate sodium. Monitoring of serum electrolytes should be started in suspected cases and appropriate management should be implemented, however, normalisation of electrolyte imbalance might not be achieved without discontinuation of colistimethate sodium.

Intravenous colistimethate sodium does not cross the blood brain barrier to a clinically relevant extent. The use of intrathecal or Intracerebroventricular administration of colistimethate sodium in the treatment of meningitis was not systematically investigated in clinical trials and is supported by case reports only. Data supporting the posology are very limited. The most commonly observed adverse effect of CMS administration was aseptic meningitis (see section 4.8).

Sodium

This medicinal product contains less than 1 mmol sodium (23 mg) per vial, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

Concomitant use of intravenous colistimethate sodium with other medications that are potentially nephrotoxic or neurotoxic should be undertaken with great caution. Combined administration with drugs presenting renal toxicity increases the risk of nephrotoxicity. If such combination is necessary, renal monitoring has to be strengthened. Such medicines are mainly represented by iodinated contrast agents, aminoglycosides antibiotics such as gentamicin, amikacin, netilmicin and tobramycin, organoplatinum compounds, high-dose methotrexate, some antiviral agents (such as "ciclovirs", foscarnet), pentamidine, ciclosporin or tacrolimus.

There may be an increased risk of nephrotoxicity if given concomitantly with cephalosporin antibiotics.

Caution should be taken with concomitant use with other formulations of colistimethate sodium as there is little experience and there is a possibility of summative toxicity.

No *in vivo* interaction studies have been performed. The mechanism of conversion of colistimethate sodium to the active substance, colistin, is not characterised. The mechanism of colistin clearance, including renal handling, is equally unknown. Colistimethate sodium or colistin did not induce the activity of any P 450 (CYP) enzyme tested (CYP1A2, 2B6, 2C8, 2C9, 2C19 and 3A4/5) in *in vitro* studies in human hepatocytes.

The potential for drug-drug interactions should be borne in mind when colistimethate sodium co- administered with drugs known to inhibit or induce drug metabolising enzymes or drugs known to be substrates for renal carrier mechanisms.

Due to the effects of colistin on the release of acetylcholine, non-depolarising muscle relaxants should be used with caution in patients receiving colistimethate sodium as their effects could be prolonged (see section 4.4).

Co-treatment with colistimethate sodium and macrolides such as azithromycin and clarithromycin, or fluoroquinolones such as norfloxacin and ciprofloxacin should be undertaken with caution in patients with myasthenia gravis (see section 4.4).

Oral anticoagulants

Simultaneous administration of colistimethate with oral anticoagulants may augment its anti-coagulant effects. The risk may vary with the underlying infection, age and general status of the patient so that the contribution of colistimethate to the increase in INR (international normalised ratio) is difficult to assess. The INR should be monitored frequently during and shortly after co-administration of colistimethate with oral anticoagulants.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no adequate data from the use of colistimethate sodium in pregnant women. Single dose studies in human pregnancy show that colistimethate sodium crosses the placental barrier and there may be a risk of fetal toxicity if repeated doses are given to pregnant patients. Animal studies are insufficient with respect to the effect of colistimethate sodium on reproduction and development (see section 5.3, Preclinical safety data). Colistimethate sodium should be used in pregnancy only if the benefit to the mother outweighs the potential risk to the fetus.

Breast-feeding

Colistimethate sodium is secreted in breast milk. Breast feeding is not recommended during therapy.

Fertility

Data on the possible impact of colistimethate sodium on human fertility are not available.

4.7 Effects on ability to drive and use machines

During parenteral treatment with colistimethate sodium neurotoxicity may occur with the possibility of dizziness, confusion or visual disturbance. Patients should be warned not to drive or operate machinery if these effects occur.

4.8 Undesirable effects

The most commonly reported adverse reaction is renal function impairment, and more rarely renal failure, usually following use of higher than recommended doses in patients with normal renal function, or failure to reduce the dosage in patients with renal impairment or when used concomitantly with other nephrotoxic antibiotics. The effect is usually reversible on discontinuation of therapy, but rarely intervention (renal replacement therapy) may be required.

High serum concentrations of colistimethate sodium, which may be associated with overdosage or failure to reduce the dosage in patients with renal impairment, have been reported to lead to neurotoxic effects such as facial paraesthesia, muscle weakness, vertigo, slurred speech, vasomotor instability, visual disturbances, confusion, psychosis and apnoea. Concomitant use with either non-depolarising muscle relaxants or antibiotics with similar neurotoxic effects can also lead to neurotoxicity. Dose reduction of colistimethate sodium may relieve symptoms.

Hypersensitivity reactions such as skin rash and angioedema have been known to occur. In the event such reactions occur, treatment with colistimethate sodium should be withdrawn.

Adverse reactions are tabulated below by system organ class and frequency. Frequencies are defined as Very common ($\geq 1/10$): common ($\geq 1/100$ to $< 1/10$): uncommon ($\geq 1/1,000$ to $< 1/100$): rare ($\geq 1/10,000$ to $< 1/1,000$) and very rare ($< 1/10,000$), not known (cannot be estimated from the available data)

Body system	Frequency	Reported adverse reaction
Immune system disorders	Not known	Hypersensitivity reactions such as skin rash and angioedema
Nervous system disorders	Very Common	Neurotoxicity such as, facial, mouth and peri-oral paraesthesia, headache, and muscle weakness
	Not known	Dizziness Ataxia

Skin and subcutaneous tissue disorders	Very Common	Pruritus
Renal and urinary disorders	Very Common	Renal impairment demonstrated by increased blood creatinine and /or urea and /or decreased creatinine renal clearance
	Rare	Renal failure
Metabolism and nutrition disorders	Not known	Pseudo-Bartter syndrome*
General disorders and administration site conditions	Not known	Injection site reaction

* See section 4.4.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via:

HPRA Pharmacovigilance,

Website: www.hpra.ie.

4.9 Overdose

Overdose can result in neuromuscular blockade that can lead to muscular weakness, apnoea and possible respiratory arrest. Overdose can also cause acute renal failure characterised by decreased urine output and increased serum concentrations of BUN and creatinine.

There is no specific antidote, manage by supportive treatment. Measures to increase the rate of elimination of colistin e.g. mannitol diuresis, prolonged haemodialysis or peritoneal dialysis may be tried, but effectiveness is unknown.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: antibacterials for systemic use, other antibacterials, polymyxins,

ATC Code: J01XB01

Mechanism of action:

Colistin is cyclic polypeptide antibacterial agent belonging to the polymyxins work by damaging the cell membrane and the resulting physiological effects are lethal to bacterium. Polymyxins are selective for aerobic Gram-negative bacteria that have a hydrophobic outer membrane.

Resistance

Resistant bacteria are characterised by modification of the phosphate groups of lipopolysaccharide, which become substituted with ethanolamine or aminoarabinose. Naturally resistant Gram-negative bacteria, such as *Proteus mirabilis* and *Burkholderia cepacia*, show complete substitution of their lipid phosphate by ethanolamine or aminoarabinose.

Cross resistance between colistin (polymyxin E) and polymyxin B is expected. Since the mechanism of action of the polymyxins is different from that of other antibacterial agents, resistance to colistin and polymyxin by the above mechanism alone would not be expected to result in resistance to other drug classes.

PK/PD relationship

Polymyxins have been reported to have a concentration-dependent bactericidal effect on susceptible bacteria. fAUC/ MIC is considered to be correlated with clinical efficacy.

Susceptibility testing breakpoints

MIC (minimum inhibitory concentration) interpretive criteria for susceptibility testing have been established by the European Committee on Antimicrobial Susceptibility Testing (EUCAST) for colistimethate sodium and are listed here:

https://www.ema.europa.eu/documents/other/minimum-inhibitory-concentration-mic-breakpoints_en.xlsx'.

Susceptibility

The prevalence of acquired resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert advice should be sought when the local prevalence of resistance is such that the utility of the agent in at least some types of infections is questionable.

Commonly susceptible species
<i>Acinetobacter baumannii</i> <i>Haemophilus influenzae</i> <i>Klebsiella</i> spp <i>Pseudomonas aeruginosa</i>
Species for which acquired resistance may be a problem
<i>Stenotrophomonas maltophilia</i> <i>Achromobacter xylosoxidans</i> (formerly <i>Alcaligenes xylosoxidans</i>)
Inherently resistant organisms
<i>Burkholderia cepacia</i> and related species. <i>Proteus</i> spp <i>Providencia</i> spp <i>Serratia</i> spp

5.2 Pharmacokinetic properties

The information on the pharmacokinetics of colistimethate sodium (CMS) and colistin is limited. There are indications that pharmacokinetics in critically ill patients differ from those in patients with less severe physiological derangement and from those in healthy volunteers. The following data are based on studies using HPLC to determine CMS/colistin plasma concentrations.

After infusion of colistimethate sodium the inactive pro-drug is converted to the active colistin. Peak plasma concentrations of colistin have been shown to occur with a delay of up to 7 hours after administration of colistimethate sodium in critically ill patients.

Distribution

The volume of distribution of colistin in healthy subjects is low and corresponds approximately to extracellular fluid (ECF). The volume of distribution is relevantly enlarged in critically ill subjects. Protein binding is moderate and decreases at higher concentrations. In the absence of meningeal inflammation, penetration into the cerebrospinal fluid (CSF) is minimal, but increases in the presence of meningeal inflammation.

Both CMS and colistin display linear PK in the clinically relevant dose range.

Elimination

It is estimated that approximately 30% of colistimethate sodium is converted to colistin in healthy subjects, its clearance is dependent on creatinine clearance and as renal function decreases, a greater portion of CMS is converted to colistin. In patients with very poor renal function (creatinine clearance <30 ml/min), the extent of conversion could be as high as 60 to 70%. CMS is eliminated predominantly by the kidneys via glomerular filtration. In healthy subjects, 60% to 70% of CMS is excreted unchanged in the urine within 24 hours.

The elimination of the active colistin is incompletely characterised. Colistin undergoes extensive renal tubular reabsorption and may either be cleared non-renally or undergo renal metabolism with the potential for renal accumulation. Colistin clearance is decreased in renal impairment, possibly due to increased conversion of CMS.

Half-life of colistin in healthy subjects and those with cystic fibrosis is reported to be around 3h and 4h, respectively, with a total clearance of around 3L/h. In critically ill patients, half-life has been reported to be prolonged to around 9-18h.

5.3 Preclinical safety data

Data on potential genotoxicity are limited and carcinogenicity data for colistimethate sodium are lacking. Colistimethate sodium has been shown to induce chromosomal aberrations in human lymphocytes, in vitro. This effect may be related to a reduction in mitotic index, which was also observed.

Reproductive toxicity studies in rats and mice do not indicate teratogenic properties. However, colistimethate sodium given intramuscularly during organogenesis to rabbits at 4.15 and 9.3 mg/kg resulted in talipes varus in 2.6 and 2.9% of fetuses respectively. These doses are 0.5 and 1.2 times the maximum daily human dose. In addition, increased resorption occurred at 9.3 mg/kg.

There are no other preclinical safety data of relevance to the prescriber which are additional to safety data derived from patient exposure and already included in other sections of the SPC.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

None

6.2 Incompatibilities

Mixed infusions and injections involving colistimethate sodium should be avoided.

6.3 Shelf life

Unopened vial:
2 years.

Reconstituted solutions:

Hydrolysis of colistimethate is significantly increased when reconstituted and diluted below its critical micelle concentration of about 80,000 IU per mL. Solutions below this concentration should be used immediately.

For solutions for bolus injection, the chemical and physical in-use stability of reconstituted solution in the original vial, with a concentration $\geq 80,000$ IU/mL, has been demonstrated for 24 hours at 2 to 8°C.

From a microbiological point of view, unless the method of opening/ reconstitution/ dilution precludes the risk of microbial contamination, the product should be used immediately. If not used immediately, in-use storage times and conditions are the responsibility of user.

Solutions for infusion, which have been diluted beyond the original vial volume and/or with a concentration $< 80,000$ IU/mL should be used immediately.

For solutions for intrathecal and Intracerebroventricular administration, the reconstituted product should be used immediately.

6.4 Special precautions for storage

This medicinal product does not require any special storage condition.

For storage condition after reconstitution/dilution of medicinal product, see section 6.3.

6.5 Nature and contents of container

Colistimethate sodium Accord 2 million International Units (IU) powder for solution for injection/infusion

Colistimethate sodium Accord is available in a 10 mL type I glass vial with grey bromobutyl rubber stopper and an aluminium flip-off pink seal.

Packs of 1 vial & 10 vials

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

For bolus injection:

Reconstitute the contents of the vial with not more than 10 ml water for injection or 0.9% sodium chloride.

For infusion:

The contents of the reconstituted vial may be diluted, usually with 50 ml 0.9% sodium chloride.

When the intrathecal and Intracerebroventricular routes of administration are used, the volume administered should not exceed 1 ml (reconstituted concentration 125,000 IU/ml).

Any unused medicinal product or waste material should be disposed of in accordance with local requirements

The appearance of solution after reconstitution should be clear colourless to pale yellow solution free from visible particulate matter. The reconstitution is completed in less than 3 minutes.

Solutions are for single use only and any remaining solution should be discarded.

7 MARKETING AUTHORISATION HOLDER

Accord Healthcare Ireland Ltd.
Euro House
Euro Business Park
Little Island
Cork T45 K857
Ireland

8 MARKETING AUTHORISATION NUMBER

PA2315/283/002

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 26th September 2025

10 DATE OF REVISION OF THE TEXT

December 2025