

## Package leaflet: Information for the patient

### Treatment initiation pack:

**Apremilast axunio Pharma 10 mg + 20 mg + 30 mg film-coated tablets**

### **Apremilast axunio Pharma 30 mg film-coated tablets**

apremilast

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

### **What is in this leaflet**

1. What Apremilast axunio Pharma is and what it is used for
2. What you need to know before you take Apremilast axunio Pharma
3. How to take Apremilast axunio Pharma
4. Possible side effects
5. How to store Apremilast axunio Pharma
6. Contents of the pack and other information

### **1. What Apremilast axunio Pharma is and what it is used for**

#### **What Apremilast axunio Pharma is**

Apremilast axunio Pharma contains the active substance ‘apremilast’. This belongs to a group of medicines called phosphodiesterase 4 inhibitors, which help to reduce inflammation.

#### **What Apremilast axunio Pharma is used for**

Apremilast axunio Pharma is used to treat adults with the following conditions:

- **Active psoriatic arthritis** - if you cannot use another type of medicine called ‘Disease-Modifying Antirheumatic Drugs’ (DMARDs) or when you have tried one of these medicines and it did not work.
- **Moderate to severe chronic plaque psoriasis** - if you cannot use one of the following treatments or when you have tried one of these treatments and it did not work:
  - phototherapy - a treatment where certain areas of skin are exposed to ultraviolet light
  - systemic therapy - a treatment that affects the entire body rather than just one local area, such as ‘ciclosporin’, ‘methotrexate’ or ‘psoralen’
- **Behçet’s disease (BD)** - to treat the mouth ulcers which is a common problem for people with this illness.

#### **What psoriatic arthritis is**

Psoriatic arthritis is an inflammatory disease of the joints, usually accompanied by psoriasis, an inflammatory disease of the skin.

## **What plaque psoriasis is**

Psoriasis is an inflammatory disease of the skin, which can cause red, scaly, thick, itchy, painful patches on your skin and can also affect your scalp and nails.

## **What Behçet's disease is**

Behçet's disease is a rare type of inflammatory disease which affects many parts of the body. The most common problem is mouth ulcers.

## **How Apremilast axunio Pharma works**

Psoriatic arthritis, psoriasis and Behçet's disease are usually lifelong conditions and there is currently no cure. Apremilast axunio Pharma works by reducing the activity of an enzyme in the body called 'phosphodiesterase 4', which is involved in the process of inflammation. By reducing the activity of this enzyme, Apremilast axunio Pharma can help to control the inflammation associated with psoriatic arthritis, psoriasis and Behçet's disease, and thereby reduce the signs and symptoms of these conditions.

In adults with psoriatic arthritis, treatment with Apremilast axunio Pharma results in an improvement in swollen and painful joints and can improve your general physical function.

In adults with psoriasis, treatment with Apremilast axunio Pharma results in a reduction in psoriatic skin plaques and other signs and symptoms of the disease.

In adults with Behçet's disease, treatment with Apremilast axunio Pharma reduces the number of mouth ulcers and can stop them completely. It can also reduce the associated pain.

Apremilast axunio Pharma has also been shown to improve the quality of life in adult patients with psoriasis, adult patients with psoriatic arthritis and adult patients with Behçet's disease. This means that the impact of your condition on daily activities, relationships and other factors should be less than it was before.

## **2. What you need to know before you take Apremilast axunio Pharma**

Do not take Apremilast axunio Pharma:

- if you are allergic to apremilast or any of the other ingredients of this medicine (listed in section 6).
- if you are pregnant or think you may be pregnant.

## **Warnings and precautions**

Talk to your doctor or pharmacist before taking Apremilast axunio Pharma.

## **Depression and suicidal thoughts**

Tell your doctor before starting Apremilast axunio Pharma if you have depression which is getting worse with thoughts of suicide.

You or your caregiver should also tell your doctor straight away of any changes in behaviour or mood, feelings of depression and of any suicidal thoughts you may have after taking Apremilast axunio Pharma.

## **Severe kidney problems**

If you have severe kidney problems, your dose will be different – see section 3.

## **If you are underweight**

Talk to your doctor while taking Apremilast axunio Pharma if you lose weight without meaning to.

## **Gut problems**

If you experience severe diarrhoea, nausea, or vomiting, you should talk to your doctor.

## **Children and adolescents**

Apremilast axunio Pharma has not been studied in children and adolescents, therefore it is not recommended for use in children and adolescents aged 17 years and under.

## **Other medicines and Apremilast axunio Pharma**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes medicines obtained without a prescription and herbal medicines. This is because Apremilast axunio Pharma can affect the way some other medicines work. Also some other medicines can affect the way Apremilast axunio Pharma works.

In particular, tell your doctor or pharmacist before taking Apremilast axunio Pharma if you are taking any of the following medicines:

- rifampicin – an antibiotic used for tuberculosis
- phenytoin, phenobarbital and carbamazepine - medicines used in the treatment of seizures or epilepsy
- St John's Wort – a herbal medicine for mild anxiety and depression.

## **Pregnancy and breast-feeding**

### **Do not take Apremilast axunio Pharma if you are pregnant or think you may be pregnant**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

There is little information about the effects of Apremilast axunio Pharma in pregnancy. You should not become pregnant while taking this medicine and should use an effective method of contraception during treatment with Apremilast axunio Pharma.

It is not known if this medicine passes into human milk. You should not use Apremilast axunio Pharma while breast-feeding.

## **Driving and using machines**

Apremilast axunio Pharma has no effect on the ability to drive and use machines.

## **Apremilast axunio Pharma contains lactose and sodium**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

Apremilast axunio Pharma contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

### 3. How to take Apremilast axunio Pharma

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

#### How much to take

- When you first start taking Apremilast axunio Pharma, you will receive a ‘treatment initiation pack’ which contains all the doses as listed in the table below.
- The ‘treatment initiation pack’ is clearly labelled to make sure you take the correct tablet at the correct time.
- Your treatment will start at a lower dose and will gradually be increased over the first 6 days of treatment.
- The ‘treatment initiation pack’ will also contain enough tablets for another 8 days at the recommended dose (days 7 to 14).
- The recommended dose of Apremilast axunio Pharma is 30 mg twice a day after the titration phase is complete - one 30 mg dose in the morning and one 30 mg dose in the evening, approximately 12 hours apart, with or without food.
- This is a total daily dose of 60 mg. By the end of day 6, you will have reached this recommended dose.
- Once the recommended dose has been reached, you will only get the 30 mg tablet strength in your prescribed packs. You will only ever need to go through this stage of gradually increasing your dose once, even if you re-start treatment.

| Day           | Morning Dose  | Evening Dose              | Total Daily Dose |
|---------------|---------------|---------------------------|------------------|
| Day 1         | 10 mg (pink)  | <b>Do not take a dose</b> | 10 mg            |
| Day 2         | 10 mg (pink)  | 10 mg (pink)              | 20 mg            |
| Day 3         | 10 mg (pink)  | 20 mg (brown)             | 30 mg            |
| Day 4         | 20 mg (brown) | 20 mg (brown)             | 40 mg            |
| Day 5         | 20 mg (brown) | 30 mg (beige)             | 50 mg            |
| Day 6 onwards | 30 mg (beige) | 30 mg (beige)             | 60 mg            |

#### People with severe kidney problems

If you have severe kidney problems, then the recommended dose of Apremilast axunio Pharma is 30 mg **once a day (morning dose)**. Your doctor will talk to you about how to increase your dose when you first start taking Apremilast axunio Pharma.

#### How and when to take Apremilast axunio Pharma

- Apremilast axunio Pharma is for oral use.
- Swallow the tablets whole, preferably with water.
- You can take the tablets either with or without food.
- Take Apremilast axunio Pharma at about the same time each day, one tablet in the morning and one tablet in the evening.

If your condition has not improved after six months of treatment, you should talk to your doctor.

#### If you take more Apremilast axunio Pharma than you should

If you take more Apremilast axunio Pharma than you should, talk to a doctor or go to a hospital straight away. Take the medicine pack and this leaflet with you.

### **If you forget to take Apremilast axunio Pharma**

- If you miss a dose of Apremilast axunio Pharma, take it as soon as you remember. If it is close to the time for your next dose, just skip the missed dose. Take the next dose at your regular time.
- Do not take a double dose to make up for a forgotten dose.

### **If you stop taking Apremilast axunio Pharma**

- You should continue taking Apremilast axunio Pharma until your doctor tells you to stop.
- Do not stop taking Apremilast axunio Pharma without talking to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

### **Serious side effects – depression and suicidal thoughts**

Tell your doctor straight away about any changes in behaviour or mood, feelings of depression, thoughts of suicide or suicidal behaviour (this is uncommon).

#### **Very common side effects** (may affect more than 1 in 10 people)

- diarrhoea
- nausea
- headache
- upper respiratory tract infections such as cold, runny nose, sinus infection

#### **Common side effects** (may affect up to 1 in 10 people)

- cough
- back pain
- vomiting
- feeling tired
- stomach pain
- loss of appetite
- frequent bowel movements
- difficulty sleeping (insomnia)
- indigestion or heartburn
- inflammation and swelling of the tubes in your lungs (bronchitis)
- common cold (nasopharyngitis)
- depression
- migraine
- tension headache

#### **Uncommon side effects** (may affect up to 1 in 100 people)

- rash
- hives (urticaria)
- weight loss
- allergic reaction
- bleeding in the bowel or in the stomach
- suicidal ideation or behaviour

#### **Not known side effects** (frequency cannot be estimated from the available data):

- severe allergic reaction (may include swelling of the face, lips, mouth, tongue, or throat that

may lead to difficulty breathing or swallowing)

If you are 65 years of age or older, you might have a higher risk of complications of severe diarrhoea, nausea and vomiting. If your gut problems become severe, you should talk to your doctor.

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance Website: [www.hpra.ie](http://www.hpra.ie).

By reporting side effects, you can help provide more information on the safety of this medicine.

## **5. How to store Apremilast axunio Pharma**

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the blister or on the wallet or on the carton after EXP. The expiry date refers to the last day of that month.
- This medicine does not require any special storage conditions.
- Do not use this medicine if you notice any damage or signs of tampering to the medicine packaging.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Apremilast axunio Pharma contains**

The active substance is apremilast.

#### Treatment initiation pack:

Each film-coated tablet contains 10 mg, 20 mg or 30 mg of apremilast.

#### Apremilast axunio Pharma 30 mg film-coated tablets

Each film-coated tablet contains 30 mg apremilast.

The other ingredients in the tablet core are lactose monohydrate, cellulose microcrystalline, croscarmellose sodium, magnesium stearate and in the film-coating hypromellose, titanium dioxide (E 171), macrogol, iron oxide red (E 172), iron oxide yellow (E172) and iron oxide black (E 172) (30 mg only).

### **What Apremilast axunio Pharma looks like and contents of the pack**

Apremilast axunio Pharma 10 mg are pink, oval, biconvex film-coated tablets with '10' engraved on one side.

Apremilast axunio Pharma 20 mg are brown, oval, biconvex film-coated tablets with '20' engraved on one side.

Apremilast axunio Pharma 30 mg are beige, oval, biconvex film-coated tablets with '30' engraved on one side.

#### Pack sizes

#### 10 mg, 20 mg and 30 mg film-coated tablets (initiation pack)

The treatment initiation pack is a wallet containing 27 film-coated tablets: 4 x 10 mg film-coated

tablets, 4 x 20 mg film-coated tablets and 19 x 30 mg film-coated tablets.

#### 30 mg film-coated tablets

The one-month standard pack contains blisters of 56 film-coated tablets.

The one-month standard pack contains perforated unit-dose blisters of 56 x 1 film-coated tablet.

The three-month standard pack contains blisters of 168 film-coated tablets.

The three-month standard pack contains perforated unit-dose blisters of 168 x 1 film-coated tablet.

#### **Marketing Authorisation Holder**

axunio Pharma GmbH  
Van-der-Smissen-Str. 1  
22767 Hamburg  
Germany

#### **Manufacturer**

Delorbis Pharmaceuticals Ltd.  
17 Athinon Str., Ergates Industrial Area  
2643 Ergates, Lefkosia  
Cyprus

**This medicine is authorised in the Member States of the European Economic Area and in the United Kingdom (Northern Ireland) under the following names:**

|         |                                                                                                                             |
|---------|-----------------------------------------------------------------------------------------------------------------------------|
| Ireland | Apremilast axunio Pharma 10 mg + 20 mg + 30 mg film-coated tablets<br>Apremilast axunio Pharma 30 mg film-coated tablets    |
| Germany | Apremilast axunio Pharma Starterpackung 10 mg + 20 mg + 30 mg Filmtabletten<br>Apremilast axunio Pharma 30 mg Filmtabletten |

**This leaflet was last revised in September 2025.**