

# Summary of Product Characteristics

## 1 NAME OF THE MEDICINAL PRODUCT

Eptifibatide Jed 0.75 mg/ml solution for infusion

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml of solution for infusion contains 0.75 mg of eptifibatide.

One vial of 100 ml of solution for infusion contains 75 mg of eptifibatide.

For the full list of excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Solution for infusion. Clear, colourless or almost colourless solution.

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Eptifibatide Jed is intended for use with acetylsalicylic acid and unfractionated heparin.

Eptifibatide Jed is indicated for the prevention of early myocardial infarction in adults presenting with unstable angina or non-Q-wave myocardial infarction, with the last episode of chest pain occurring within 24 hours and with electrocardiogram (ECG) changes and/or elevated cardiac enzymes.

Patients most likely to benefit from Eptifibatide Jed treatment are those at high risk of developing myocardial infarction within the first 3-4 days after onset of acute angina symptoms including for instance those that are likely to undergo an early PTCA (Percutaneous Transluminal Coronary Angioplasty) (see section 5.1).

### 4.2 Posology and method of administration

This product is for hospital use only. It should be administered by specialist physicians experienced in the management of acute coronary syndromes.

Eptifibatide Jed solution for infusion must be used in conjunction with Eptifibatide Jed solution for injection.

Concurrent administration of heparin is recommended unless this is contraindicated for reasons such as a history of thrombocytopenia associated with use of heparin (see 'Heparin administration', section 4.4). Eptifibatide Jed is also intended for concurrent use with acetylsalicylic acid, as it is part of standard management of patients with acute coronary syndromes, unless its use is contraindicated.

#### Posology

*Adults (≥ 18 years of age) presenting with unstable angina (UA) or non-Q-wave myocardial infarction (NQMI)*

The recommended dosage is an intravenous bolus of 180 microgram/kg administered as soon as possible following diagnosis, followed by a continuous infusion of 2 microgram/kg/min for up to 72 hours, until initiation of coronary artery bypass graft (CABG) surgery, or until discharge from the hospital (whichever occurs first). If Percutaneous Coronary Intervention (PCI) is performed during eptifibatide therapy, continue the infusion for 20-24 hours post-PCI for an overall maximum duration of therapy of 96 hours.

#### *Emergency or semi-elective surgery*

If the patient requires emergency or urgent cardiac surgery during the course of eptifibatide therapy, terminate the infusion immediately. If the patient requires semi-elective surgery, stop the eptifibatide infusion at an appropriate time to allow time for platelet function to return towards normal.

*Hepatic impairment*

Experience in patients with hepatic impairment is very limited. Administer with caution to patients with hepatic impairment in whom coagulation could be affected (see section 4.3, prothrombin time). It is contraindicated in patients with clinically significant hepatic impairment.

*Renal impairment*

In patients with moderate renal impairment (creatinine clearance  $\geq 30$  -  $< 50$  ml/min), an intravenous bolus of 180 microgram/kg should be administered followed by a continuous infusion dose of 1.0 microgram/kg/min for the duration of therapy. This recommendation is based on pharmacodynamic and pharmacokinetic data. The available clinical evidence cannot however confirm that this dose modification results in a preserved benefit (see section 5.1). Use in patients with more severe renal impairment is contraindicated (see section 4.3).

*Paediatric population*

It is not recommended for use in children and adolescents below 18 years of age, due to a lack of data on safety and efficacy.

**4.3 Contraindications**

Eptifibatide Jed must not be used to treat patients with:

- hypersensitivity to the active substance or to any of the excipients listed in section 6.1
- evidence of gastrointestinal bleeding, gross genitourinary bleeding or other active abnormal bleeding within the previous 30 days of treatment
- history of stroke within 30 days or any history of haemorrhagic stroke
- known history of intracranial disease (neoplasm, arteriovenous malformation, aneurysm)
- major surgery or severe trauma within past 6 weeks
- a history of bleeding diathesis
- thrombocytopenia ( $< 100,000$  cells/mm<sup>3</sup>)
- prothrombin time  $> 1.2$  times control, or International Normalized Ratio (INR)  $\geq 2.0$
- severe hypertension (systolic blood pressure  $> 200$  mm Hg or diastolic blood pressure  $> 110$  mm Hg on antihypertensive therapy)
- severe renal impairment (creatinine clearance  $< 30$  ml/min) or dependency on renal dialysis
- clinically significant hepatic impairment
- concomitant or planned administration of another parenteral glycoprotein (GP) IIb/IIIa inhibitor

**4.4 Special warnings and precautions for use***Bleeding*

Eptifibatide Jed is an antithrombotic agent that acts by inhibition of platelet aggregation; therefore the patient must be observed carefully for indications of bleeding during treatment (see section 4.8). Women, the elderly, patients with low body weight or with moderate renal impairment (creatinine clearance  $\geq 30$  -  $< 50$  ml/min) may have an increased risk of bleeding. Monitor these patients closely with regard to bleeding.

An increased risk of bleeding may also be observed in patients who receive early administration of eptifibatide (e.g. upon diagnosis) compared to receiving it immediately prior to PCI, as seen in the Early ACS trial. Unlike the approved posology in the EU, all patients in this trial were administered a double bolus before the infusion (see section 5.1).

Bleeding is most common at the arterial access site in patients undergoing percutaneous arterial procedures. All potential bleeding sites, (e.g., catheter insertion sites; arterial, venous, or needle puncture sites; cutdown sites; gastrointestinal and genitourinary tracts) must be observed carefully. Other potential bleeding sites such as central and peripheral nervous system and retroperitoneal sites, must be carefully considered too.

Because Eptifibatide Jed inhibits platelet aggregation, caution must be employed when it is used with other medicinal products that affect haemostasis, including ticlopidine, clopidogrel, thrombolytics, oral anticoagulants, dextran solutions, adenosine, sulfapyrazone, prostacyclin, non-steroidal anti-inflammatory agents, or dipyridamole (see section 4.5).

There is no experience with Eptifibatide Jed and low molecular weight heparins.

There is limited therapeutic experience with Eptifibatide Jed in patients for whom thrombolytic therapy is generally indicated (e.g. acute transmural myocardial infarction with new pathological Q-waves or elevated ST-segments or left bundle branch block in the ECG). Consequently the use of Eptifibatide Jed is not recommended in these circumstances (see section 4.5).

Eptifibatide Jed infusion should be stopped immediately if circumstances arise that necessitate thrombolytic therapy or if the patient must undergo an emergency CABG surgery or requires an intraortic balloon pump.

If serious bleeding occurs that is not controllable with pressure, the Eptifibatide Jed infusion should be stopped immediately and any unfractionated heparin that is given concomitantly.

#### *Arterial procedures*

During treatment with eptifibatide there is a significant increase in bleeding rates, especially in the femoral artery area, where the catheter sheath is introduced. Take care to ensure that only the anterior wall of the femoral artery is punctured. Arterial sheaths may be removed when coagulation has returned to normal (e.g. when activated clotting time (ACT) is less than 180 seconds (usually 2- 6 hours after discontinuation of heparin). After removal of the introducer sheath, careful haemostasis must be ensured under close observation.

#### *Thrombocytopenia and Immunogenicity related to GPIIb/IIIa inhibitors*

Eptifibatide Jed inhibits platelet aggregation, but does not appear to affect the viability of platelets. As demonstrated in clinical trials, the incidence of thrombocytopenia was low, and similar in patients treated with eptifibatide or placebo.

Thrombocytopenia, including acute profound thrombocytopenia, has been observed with eptifibatide administration post-marketing (see section 4.8)

The mechanism, whether immune- and/or non-immune-mediated, by which eptifibatide may induce thrombocytopenia is not fully understood. However, treatment with eptifibatide was associated with antibodies that recognise GPIIb/IIIa occupied by eptifibatide, suggesting an immune-mediated mechanism. Thrombocytopenia occurring after first exposure to a GPIIb/IIIa inhibitor may be explained by the fact that antibodies are naturally present in some normal individuals.

Since either repeat exposure with any GP IIb/IIIa ligand-mimetic agent (like abciximab or eptifibatide) or first-time exposure to a GP IIb/IIIa inhibitor may be associated with immune-mediated thrombocytopenic responses, monitoring is required, i.e. platelet counts should be monitored prior to treatment, within 6 hours of administration, and at least once daily thereafter while on therapy and immediately at clinical signs of unexpected bleeding tendency.

If either a confirmed platelet decrease to  $< 100,000/\text{mm}^3$  or acute profound thrombocytopenia is observed, discontinuation of each treatment medication having known or suspected thrombocytopenic effects, including eptifibatide, heparin and clopidogrel, should be considered immediately. The decision to use platelet transfusions should be based upon clinical judgment on an individual basis.

In patients with previous immune-mediated thrombocytopenia from other parenteral GP IIb/IIIa inhibitors, there are no data with the use of Eptifibatide Jed. Therefore, it is not recommended to administer eptifibatide in patients who have previously experienced immune mediated thrombocytopenia with GP IIb/IIIa inhibitors, including eptifibatide.

#### *Heparin administration*

Heparin administration is recommended unless a contraindication (such as a history of thrombocytopenia associated with use of heparin) is present.

UA/NQMI: For a patient who weighs  $\geq 70$  kg, it is recommended that a bolus dose of 5,000 units is given, followed by a constant intravenous infusion of 1,000 units/hr. If the patient weighs  $< 70$  kg, a bolus dose of 60 units/kg is recommended, followed by an infusion of 12 units/kg/hr. The activated partial thromboplastin time (aPTT) must be monitored in order to maintain a value between 50 and 70 seconds, above 70 seconds there may be an increased risk of bleeding.

If PCI is to be performed in the setting of UA/NQMI, monitor the activated clotting time (ACT) to maintain a value between 300-350 seconds. Stop heparin administration if the ACT exceeds 300 seconds; do not administer until the ACT falls below 300 seconds.

#### *Monitoring of laboratory values*

Before infusion of Eptifibatide Jed, the following laboratory tests are recommended to identify pre-existing haemostatic abnormalities: prothrombin time (PT) and aPTT, serum creatinine, platelet count,

haemoglobin and haematocrit levels. Haemoglobin, haematocrit, and platelet count are to be monitored as well within 6 hours after start of therapy and at least once daily thereafter while on therapy (or more often if there is evidence of a marked decrease). If the platelet count falls below 100,000/mm<sup>3</sup>, further platelet counts are required to rule out pseudothrombocytopenia. Discontinue unfractionated heparin. In patients undergoing PCI, measure the ACT also.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

##### *Warfarin and dipyridamole*

Eptifibatide Jed did not appear to increase the risk of major and minor bleeding associated with concomitant use of warfarin and dipyridamole. Eptifibatide Jed-treated patients who had a prothrombin time (PT) > 14.5 seconds and received warfarin concomitantly did not appear to be at an increased risk of bleeding.

##### *Eptifibatide Jed and thrombolytic agents*

Data are limited on the use of Eptifibatide Jed in patients receiving thrombolytic agents. There was no consistent evidence that eptifibatide increased the risk of major or minor bleeding associated with tissue plasminogen activator in either a PCI or an acute myocardial infarction study. Eptifibatide appeared to increase the risk of bleeding when administered with streptokinase in an acute myocardial infarction study. The combination of reduced dose tenecteplase and eptifibatide compared to placebo and eptifibatide significantly increased the risk of both major and minor bleeding when administered concomitantly in an acute ST-elevation myocardial infarction study.

In an acute myocardial infarction study involving 181 patients, eptifibatide (in regimens up to a bolus injection of 180 microgram/kg, followed by an infusion up to 2 microgram/kg/min for up to 72 hours) was administered concomitantly with streptokinase (1.5 million units over 60 minutes). At the highest infusion rates (1.3 microgram/kg/min and 2.0 microgram/kg/min) studied, eptifibatide was associated with an increased incidence of bleeding and transfusions compared to the incidence seen when streptokinase was given alone.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

There are no adequate data from the use of eptifibatide in pregnant women.

Animal studies are insufficient with respect to effects on pregnancy, embryonal/foetal development, parturition or postnatal development (see section 5.3). The potential risk for humans is unknown. Eptifibatide Jed should not be used during pregnancy unless clearly necessary.

##### Breast-feeding

It is not known whether eptifibatide is excreted in human milk. Interruption of breast-feeding during the treatment period is recommended.

#### **4.7 Effects on ability to drive and use machines**

Not relevant, as Eptifibatide Jed is intended for use only in hospitalised patients.

#### **4.8 Undesirable effects**

The majority of adverse reactions experienced by patients treated with eptifibatide were generally related to bleeding or to cardiovascular events that occur frequently in this patient population.

##### *Clinical Trials*

The data sources used to determine adverse reaction frequency descriptors included two phase III clinical studies (PURSUIT and ESPRIT). These trials are briefly described below.

**PURSUIT:** This was a randomised, double-blind evaluation of the efficacy and safety of eptifibatide versus placebo for reducing mortality and myocardial (re)infarction in patients with unstable angina or non-Q-wave myocardial infarction.

ESPRIT: This was a double-blind, multicentre, randomised, parallel-group, placebo-controlled trial evaluating the safety and efficacy of eptifibatide therapy in patients scheduled to undergo non-emergent percutaneous coronary intervention (PCI) with stent implantation.

In PURSUIT, bleeding and non-bleeding events were collected from hospital discharge to the 30 day visit. In ESPRIT, bleeding events were reported at 48 hours, and non-bleeding events were reported at 30 days. While Thrombolysis in Myocardial Infarction TIMI bleeding criteria were used to categorize the incidence of major and minor bleeding in both the PURSUIT and the ESPRIT trials, PURSUIT data was collected within 30 days while ESPRIT data was limited to events within 48 hours or discharge, whichever came first.

The undesirable effects are listed by body system and frequency. Frequencies are defined as very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ). These are absolute reporting frequencies without taking into account placebo rates. For a particular adverse reaction, if data was available from both PURSUIT and ESPRIT, then the highest reported incidence was used to assign adverse reaction frequency.

Note that causality has not been determined for all adverse reactions.

<b>Blood and Lymphatic System Disorder</b>	
Very common	Bleeding (major and minor bleeding including femoral artery access, CABG-related, gastrointestinal, genitourinary, retroperitoneal, intracranial, haematemesis, haematuria, oral/oropharyngeal, haemoglobin/haematocrit decreased and other).
Uncommon	Thrombocytopenia.
<b>Nervous System disorders</b>	
Uncommon	Cerebral ischaemia.
<b>Cardiac Disorders</b>	
Common	Cardiac arrest, ventricular fibrillation, ventricular tachycardia, congestive heart failure, atrioventricular block, atrial fibrillation.
<b>Vascular Disorders</b>	
Common	Shock, hypotension, phlebitis.

Cardiac arrest, congestive heart failure, atrial fibrillation, hypotension, and shock, which are commonly reported events from the PURSUIT trial, were events related to the underlying disease.

Administration of eptifibatide is associated with an increase in major and minor bleeding as classified by the criteria of the TIMI study group. At the recommended therapeutic dose, as administered in the PURSUIT trial involving nearly 11,000 patients, bleeding was the most common complication encountered during eptifibatide therapy. The most common bleeding complications were associated with cardiac invasive procedures (coronary artery bypass grafting (CABG)-related or at femoral artery access site).

Minor bleeding was defined in the PURSUIT trial as spontaneous gross haematuria, spontaneous haematemesis, observed blood loss with a haemoglobin decrease of more than 3 g/dl, or a haemoglobin decrease of more than 4 g/dl in the absence of an observed bleeding site. During treatment with eptifibatide in this study, minor bleeding was a very common complication ( $> 1/10$ , or 13.1% for eptifibatide versus 7.6% for placebo). Bleeding events were more frequent in patients receiving concurrent heparin while undergoing PCI, when ACT exceeded 350 seconds (see section 4.4, heparin use).

Major bleeding was defined in the PURSUIT trial as either an intracranial haemorrhage or a decrease in haemoglobin concentrations of more than 5 g/dl. Major bleeding was also very common and reported more frequently with eptifibatide than with placebo in the PURSUIT study ( $\geq 1/10$  or 10.8% versus 9.3%), but it was infrequent in the vast majority of patients who did not undergo CABG within 30 days of inclusion in the study. In patients undergoing CABG, the incidence of bleeding was not increased by eptifibatide compared to the patients treated with placebo. In the subgroup of patients undergoing PCI, major bleeding was observed commonly, in 9.7 % of eptifibatide-treated patients vs. 4.6 % of placebo-treated patients.

The incidence of severe or life threatening bleeding events with eptifibatide was 1.9% compared to 1.1% with placebo. The need for blood transfusions was increased modestly by eptifibatide treatment (11.8% versus 9.3% for placebo).

Changes during eptifibatide treatment result from its known pharmacological action, i.e., inhibition of platelet aggregation. Thus, changes in laboratory parameters associated with bleeding (e.g. bleeding time) are common and expected. No apparent differences were observed between patients treated with eptifibatide or with placebo in values for liver function (SGOT/AST, SGPT/ALT, bilirubin, alkaline phosphatase) or renal function (serum creatinine, blood urea nitrogen).

*Post-marketing experience*

<b>Blood and lymphatic system disorders</b>	
Very rare	Fatal bleeding (the majority involved central and peripheral nervous system disorders: cerebral or intracranial haemorrhages); pulmonary haemorrhage, acute profound thrombocytopenia, haematoma.
<b>Immune system disorders</b>	
Very rare	Anaphylactic reactions.
<b>Skin and subcutaneous tissue disorders</b>	
Very rare	Rash, application site disorders such as urticaria.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRAs Pharmacovigilance Website: [www.hpra.ie](http://www.hpra.ie)

**4.9 Overdose**

The experience in humans with overdose of eptifibatide is extremely limited. There was no indication of severe adverse reactions associated with administration of accidental large bolus doses, rapid infusion reported as overdose or large cumulative doses. In the PURSUIT trial, there were 9 patients who received bolus and/or infusion doses more than double the recommended dose, or who were identified by the investigator as having received an overdose. There was no excessive bleeding in any of these patients, although one patient undergoing CABG surgery was reported as having had a moderate bleed. Specifically, no patients experienced an intracranial bleed.

Potentially, an overdose of eptifibatide could result in bleeding. Because of its short half-life and rapid clearance, the activity of eptifibatide may be halted readily by discontinuing the infusion. Thus, although eptifibatide can be dialysed, the need for dialysis is unlikely.

**5 PHARMACOLOGICAL PROPERTIES****5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Antithrombotic agent (platelet aggregation inhibitors excl. heparin),  
ATC code: B01AC16

Mechanism of action

Eptifibatide, a synthetic cyclic heptapeptide containing six amino acids, including one cysteine amide and one mercaptopropionyl (desamino cysteinyl) residue, is an inhibitor of platelet aggregation and belongs to the class of RGD (arginine-glycine-aspartate)-mimetics.

Eptifibatide reversibly inhibits platelet aggregation by preventing the binding of fibrinogen, von Willebrand factor and other adhesive ligands to the glycoprotein (GP) IIb/IIIa receptors.

Pharmacodynamic effects

Eptifibatide inhibits platelet aggregation in a dose- and concentration-dependent manner as demonstrated by *ex vivo* platelet aggregation using adenosine diphosphate (ADP) and other agonists to induce platelet aggregation. The effect of eptifibatide is observed immediately after administration of a 180 microgram/kg intravenous bolus. When followed by a 2.0 microgram/kg/min continuous infusion, this regimen produces a > 80 % inhibition of ADP-induced *ex vivo* platelet aggregation, at physiologic calcium concentrations, in more than 80 % of patients. Platelet inhibition was readily reversed, with a return of platelet function towards baseline (> 50 % platelet aggregation) 4 hours after stopping a continuous infusion of 2.0 microgram/kg/min. Measurements of ADP-induced *ex vivo* platelet aggregation at physiologic calcium concentrations (D-phenylalanyl-L-prolyl-L-arginine chloromethyl ketone anticoagulant) in patients presenting with unstable angina and Non Q-Wave Myocardial Infarction showed a concentration-dependent inhibition with an IC<sub>50</sub> (50% inhibitory concentration) of approximately 550 ng/ml and an IC<sub>80</sub> (80 % inhibitory concentration) of approximately 1,100 ng/ml.

There is limited data with regards to platelet inhibition in patients with renal impairment. In patients with moderate renal impairment, (creatinine clearance 30 – 50mL/min) 100% inhibition was achieved at 24 hours following administration of 2 microgram/kg/min. In patients with severe renal impairment (creatinine clearance <30mL/min) administered 1microgram/kg/min , 80% inhibition was achieved in more than 80% of patients at 24 hours.

Clinical efficacy and safety

*PURSUIT trial*

The pivotal clinical trial for Unstable Angina (UA)/Non-Q Wave Myocardial Infarction (NQMI) was PURSUIT. This study was a 726-center, 27-country, double-blind, randomised, placebo-controlled study in 10,948 patients presenting with UA or NQMI. Patients could be enrolled only if they had experienced cardiac ischemia at rest ( $\geq$  10 minutes) within the previous 24 hours and had:

- either ST-segment changes: ST depression > 0.5 mm of less than 30 minutes or persistent ST elevation > 0.5 mm not requiring reperfusion therapy or thrombolytic agents, T-wave inversion (> 1 mm),
- or increased CK-MB.

Patients were randomised to either placebo, eptifibatide 180 microgram/kg bolus followed by a 2.0 microgram/kg/min infusion (180/2.0), or eptifibatide 180 microgram/kg bolus followed by a 1.3 microgram/kg/min infusion (180/1.3).

The infusion was continued until hospital discharge, until the time of coronary artery bypass grafting (CABG) or for up to 72 hours, whichever occurred first. If PCI was performed, the eptifibatide infusion was continued for 24 hours after the procedure, allowing for a duration of infusion up to 96 hours.

The 180/1.3 arm was stopped after an interim analysis, as prespecified in the protocol, when the two active-treatment arms appeared to have a similar incidence of bleeding.

Patients were managed according to the usual standards of the investigational site; frequencies of angiography, PCI and CABG therefore differed widely from site to site and from country to country. Of the patients in PURSUIT, 13% were managed with PCI during eptifibatide infusion, of whom approximately 50 % received intracoronary stents; 87 % were managed medically (without PCI during eptifibatide infusion).

The vast majority of patients received acetylsalicylic acid (75-325 mg once daily). Unfractionated heparin was administered intravenously or subcutaneously at the physician's discretion, most commonly as an intravenous bolus of 5,000 U followed by a continuous infusion of 1,000 U/h. A target aPTT of 50-70 seconds was recommended. A total of 1,250 patients underwent PCI within 72 hours after randomisation, in which case they received intravenous unfractionated heparin to maintain an activated clotting time (ACT) of 300-350 seconds.

The primary endpoint of the study was the occurrence of death from any cause or new myocardial infarction (MI) (evaluated by a blinded Clinical Events Committee) within 30 days of randomisation. The component MI could be defined as asymptomatic with enzymatic elevation of CK-MB or new Q wave.

Compared to placebo, eptifibatide administered as 180/2.0 significantly reduced the incidence of the primary endpoint events (table 1): this represents around 15 events avoided for 1,000 patients treated:

<b>Table 1 Incidence of Death/CEC-Assessed MI («Treated as Randomised» Population)</b>			
Time	Placebo	Eptifibatide	p-Value
30 days	743/4,697 (15.8 %)	667/4,680 (14.3 %)	0.034 <sup>a</sup>
a: Pearson's chi-square test of difference between placebo and eptifibatide.			

Results on the primary endpoint were principally attributed to the occurrence of myocardial infarction. The reduction in the incidence of endpoint events in patients receiving eptifibatide appeared early during treatment (within the first 72-96 hours) and this reduction was maintained through 6 months, without any significant effect on mortality.

Patients most likely to benefit from eptifibatide treatment are those at high risk of developing myocardial infarction within the first 3-4 days after onset of acute angina.

According to epidemiological findings, a higher incidence of cardiovascular events has been associated with certain indicators, for instance:

- age
- elevated heart rate or blood pressure
- persistent or recurrent ischemic cardiac pain
- marked ECG changes (in particular ST-segment abnormalities)
- raised cardiac enzymes or markers (e.g. CK-MB, troponins) and
- heart failure

PURSUIT was conducted at a time when the standard of care of managing acute coronary syndromes was different from that of present times in terms of thienopyridine use and the routine use of intracoronary stents.

#### *ESPRIT trial*

ESPRIT (Enhanced Suppression of the Platelet IIb/IIIa Receptor with eptifibatide Therapy) was a double-blind, randomised, placebo-controlled trial (n= 2,064) for nonurgent PCI with intracoronary stenting.

All patients received routine standard of care and were randomised to either placebo or eptifibatide (2 bolus doses of 180 microgram/kg and a continuous infusion until discharge from hospital or a maximum of 18-24 hours).

The first bolus and the infusion were started simultaneously, immediately before the PCI procedure and were followed by a second bolus 10 minutes after the first. The rate of infusion was 2.0 microgram/kg/min for patients with serum creatinine  $\leq$  175 micromols/l or 1.0 microgram/kg/min for serum creatinine  $>$  175 up to 350 micromols/l.

In the eptifibatide arm of the trial, virtually all patients received aspirin (99.7 %), and 98.1 % received a thienopyridine, (clopidogrel in 95.4 % and ticlopidine in 2.7 %). On the day of PCI, prior to catheterization, 53.2 received a thienopyridine (clopidogrel 52.7 %; ticlopidine 0.5 %) – mostly as a loading dose (300 mg or more). The placebo arm was comparable (aspirin 99.7 %, clopidogrel 95.9 %, ticlopidin 2.6 %).

The ESPRIT trial used a simplified regimen of heparin during PCI that consisted of an initial bolus of 60 units/kg, with a target ACT of 200 - 300 seconds. The primary endpoint of the trial was death (D), MI, urgent target vessel revascularisation (UTVR), and acute antithrombotic rescue with GP IIb/IIIa inhibitor therapy (RT) within 48 hours of randomisation.

MI was identified per the CK-MB core laboratory criteria. For this diagnosis, within 24 hours after the index PCI procedure, there had to be at least two CK-MB values  $\geq$  3 x the upper limit of normal; thus, validation by the CEC was not required. MI could also be reported following CEC adjudication of an investigator report.

The primary endpoint analysis [quadruple composite of death, MI, urgent target vessel revascularisation (UTVR) and thrombolytic bail-out (TBO) at 48 hours] showed a 37 % relative and 3.9 % absolute reduction in the eptifibatide group (6.6 % events versus 10.5 %,  $p = 0.0015$ ). Results on the primary endpoint were mainly attributed to the reduction of enzymatic MI occurrence, identified as the occurrence of early elevation of cardiac enzymes after PCI (80 out of 92 MIs in the placebo group vs. 47 out of 56 MIs in the eptifibatide group). The clinical relevance of such enzymatic MIs is still controversial.

Similar results were also obtained for the 2 secondary endpoints assessed at 30 days: a triple composite of death, MI and UTVR, and the more robust combination of death and MI.

The reduction in the incidence of endpoint events in patients receiving eptifibatide appeared early during treatment. There was no increased benefit thereafter, up to 1 year.

#### *Prolongation of bleeding time*

Administration of eptifibatide by intravenous bolus and infusion causes up to a 5-fold increase in bleeding time. This increase is readily reversible upon discontinuation of the infusion with bleeding times returning towards baseline in approximately 6 (2-8) hours. When administered alone,

eptifibatide has no measurable effect on prothrombin time (PT) or activated partial thromboplastin time (aPTT).

### *EARLY-ACS trial*

EARLY ACS (Early Glycoprotein IIb/IIIa Inhibition in Non-ST-segment Elevation Acute Coronary Syndrome) was a study of early routine eptifibatide versus placebo (with delayed provisional use of eptifibatide in the catheterization laboratory) used in combination with antithrombotic therapies (ASA, UFH, bivalirudin, fondaparinux or low molecular weight heparin), in subjects with high-risk NSTEMI

ACS. Patients were to undergo an invasive strategy for further management after receiving study drug for 12 to 96 hours. Patients could be medically managed, proceed to coronary artery bypass graft (CABG), or undergo percutaneous coronary intervention (PCI). Unlike the approved posology in the EU, the study used a double bolus of study drug (separated by 10 minutes) before the infusion.

Early routine eptifibatide in this high-risk NSTEMI-ACS optimally-treated population who were managed with an invasive strategy did not result in a statistically significant reduction in the composite primary endpoint of rate of death, MI, RI-UR, and TBO within 96 hours compared with a regimen of delayed provisional eptifibatide (9.3% in early eptifibatide patients vs. 10.0% in patients assigned to delayed provisional eptifibatide; odds ratio=0.920; 95% CI=0.802-1.055; p=0.234). GUSTO severe/life threatening bleeding was uncommon and comparable in both treatment groups (0.8%). GUSTO moderate or severe/life threatening bleeding occurred significantly more often with early routine eptifibatide (7.4% vs. 5.0% in delayed provisional eptifibatide group; p <0.001). Similar differences were noted for TIMI major haemorrhage (118 [2.5%] in early routine use vs. 83 [1.8%] in delayed provisional use; p=0.016).

No statistically significant benefit of early routine eptifibatide strategy was demonstrated in the subgroup of patients who were managed medically or during the medical management periods prior to PCI or CABG.

In a post hoc analysis of the EARLY ACS trial the risk benefit of dose reduction in patients with moderate renal impairment is inconclusive. The primary endpoint event rate was 11.9 % in patients who received a reduced dose (1 microgram/kg/min) vs 11.2% in patients who received the standard dose (2 microgram/kg/min) when eptifibatide was administered in the early routine fashion (p=0.81). With delayed provisional eptifibatide administration, the event rates were 10% vs 11.5% in patients who received reduced dose and standard dose respectively (p=0.61). TIMI major bleeding occurred in 2.7 % of patients who received a reduced dose (1 microgram/kg/min) vs 4.2% of patients who received the standard dose (2 microgram/kg/min) when eptifibatide was administered in the early routine fashion (p=0.36). With delayed provisional eptifibatide administration, the TIMI major events were 1.4% vs 2.0% in patients who received reduced dose and standard dose respectively (p=0.54). There were no notable differences observed with GUSTO severe bleeding rates.

## **5.2 Pharmacokinetic properties**

The pharmacokinetics of eptifibatide are linear and dose proportional for bolus doses ranging from 90 to 250 microgram/kg and infusion rates from 0.5 to 3.0 microgram/kg/min. For a 2.0 microgram/kg/min infusion, mean steady-state plasma eptifibatide concentrations range from 1.5 to 2.2 microgram/ml in patients with coronary artery disease. These plasma concentrations are achieved rapidly when the infusion is preceded by a 180 microgram/kg bolus. The extent of eptifibatide binding to human plasma protein is about 25 %. In the same population, plasma elimination half-life is approximately 2.5 hours, plasma clearance 55 to 80 ml/kg/hr and volume of distribution of approximately 185 to 260 ml/kg.

In healthy subjects, renal excretion accounted for approximately 50 % of total body clearance; approximately 50 % of the amount cleared is excreted unchanged. In patients with moderate to severe renal insufficiency (creatinine clearance < 50ml/min), the clearance of eptifibatide is reduced by approximately 50% and steady-state plasma levels are approximately doubled.

No formal pharmacokinetic interaction studies have been conducted. However, in a population pharmacokinetic study there was no evidence of a pharmacokinetic interaction between eptifibatide and the following concomitant medicinal products: amlodipine, atenolol, atropine, captopril, cefazolin, diazepam, digoxin, diltiazem, diphenhydramine, enalapril, fentanyl, furosemide, heparin, lidocaine, lisinopril, metoprolol, midazolam, morphine, nitrates, nifedipine, and warfarin.

## **5.3 Preclinical safety data**

Toxicology studies conducted with eptifibatide include single and repeated dose studies in the rat, rabbit and monkey, reproduction studies in the rat and rabbit, *in vitro* and *in vivo* genetic toxicity studies, and irritation, hypersensitivity and

antigenicity studies. No unexpected toxic effects for an agent with this pharmacologic profile were observed and findings were predictive of clinical experience, with bleeding effects being the principal adverse event. No genotoxic effects were observed with eptifibatide.

Teratology studies have been performed by continuous intravenous infusion of eptifibatide in pregnant rats at total daily doses of up to 72 mg/kg/day (about 4 times the recommended maximum daily human dose on a body surface area basis) and in pregnant rabbits at total daily doses of up to 36 mg/kg/day (about 4 times the recommended maximum daily human dose on a body surface area basis). These studies revealed no evidence of impaired fertility or harm to the foetus due to eptifibatide. Reproduction studies in animal species where eptifibatide shows a similar pharmacologic activity as in humans are not available. Consequently these studies are not suitable to evaluate the toxicity of eptifibatide on reproductive function (see section 4.6).

The carcinogenic potential of eptifibatide has not been evaluated in long-term studies.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Citric acid monohydrate  
Sodium hydroxide  
Water for injections

### **6.2 Incompatibilities**

Eptifibatide Jed is not compatible with furosemide.

In the absence of compatibility studies, Eptifibatide Jed must not be mixed with other medicinal products except those mentioned in 6.6.

### **6.3 Shelf life**

3 years

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C).  
Store in the original package in order to protect from light.

### **6.5 Nature and contents of container**

One 100 ml Type I glass vial, closed with a chlorinated butyl stopper and sealed with a crimped aluminium-plastic flip-off seal.

### **6.6 Special precautions for disposal and other handling**

Physical and chemical compatibility testing indicate that Eptifibatide Jed may be administered through an intravenous line with atropine sulfate, dobutamine, heparin, lidocaine, meperidine, metoprolol, midazolam, morphine, nitroglycerin, tissue plasminogen activator, or verapamil. Eptifibatide Jed is compatible with 0.9 % sodium chloride solution for infusion and with 0.9 % sodium chloride solution for infusion in 5 % dextrose with or without potassium chloride.

Before using, inspect the vial contents. Do not use if particulate matter or discolouration is present. Protection of Eptifibatide Jed solution from light is not necessary during administration.

Discard any unused medicinal product after opening.

## **7 MARKETING AUTHORISATION HOLDER**

JED Pharma Limited  
Questum Business Park

South Ballingarrane  
Clonmel  
Co Tipperary  
E91 V329  
Ireland

**8 MARKETING AUTHORISATION NUMBER**

PA23183/009/001

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 29<sup>th</sup> August 2025

**10 DATE OF REVISION OF THE TEXT**