

Package leaflet: Information for the patient

Sitagliptin Rowex 25 mg film-coated tablets
Sitagliptin Rowex 50 mg film-coated tablets
Sitagliptin Rowex 100 mg film-coated tablets

Sitagliptin

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Sitagliptin Rowex is and what it is used for
2. What you need to know before you take Sitagliptin Rowex
3. How to take Sitagliptin Rowex
4. Possible side effects
5. How to store Sitagliptin Rowex
6. Contents of the pack and other information

1. What Sitagliptin Rowex is and what it is used for

Sitagliptin Rowex contains the active substance sitagliptin which is a member of a class of medicines called DPP-4 inhibitors (dipeptidyl peptidase-4 inhibitors) that lowers blood sugar levels in adult patients with type 2 diabetes mellitus.

This medicine helps to increase the levels of insulin produced after a meal and decreases the amount of sugar made by the body.

Your doctor has prescribed this medicine to help lower your blood sugar, which is too high because of your type 2 diabetes. This medicine can be used alone or in combination with certain other medicines (insulin, metformin, sulphonylureas, or glitazones) that lower blood sugar, which you may already be taking for your diabetes together with a food and exercise plan.

What is type 2 diabetes?

Type 2 diabetes is a condition in which your body does not make enough insulin, and the insulin that your body produces does not work as well as it should. Your body can also make too much sugar. When this happens, sugar (glucose) builds up in the blood. This can lead to serious medical problems like heart disease, kidney disease, blindness, and amputation.

2. What you need to know before you take Sitagliptin Rowex

Do not take Sitagliptin Rowex

- if you are allergic to sitagliptin or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor or pharmacist before taking Sitagliptin Rowex.

Cases of inflammation of the pancreas (pancreatitis) have been reported in patients receiving Sitagliptin Rowex (see section 4).

If you encounter blistering of the skin it may be a sign for a condition called bullous pemphigoid. Your doctor may ask you to stop Sitagliptin Rowex.

Tell your doctor if you have or have had:

- a disease of the pancreas (such as pancreatitis)
- gallstones, alcohol dependence or very high levels of triglycerides (a form of fat) in your blood
These medical conditions can increase your chance of getting pancreatitis (see section 4).
- type 1 diabetes
- diabetic ketoacidosis (a complication of diabetes with high blood sugar, rapid weight loss, nausea or vomiting)
- any past or present kidney problems
- an allergic reaction to Sitagliptin Rowex (see section 4).

This medicine is unlikely to cause low blood sugar because it does not work when your blood sugar is low. However, when this medicine is used in combination with a sulphonylurea medicine or with insulin, low blood sugar (hypoglycaemia) can occur. Your doctor may reduce the dose of your sulphonylurea or insulin medicine.

Children and adolescents

Children and adolescents below 18 years should not use this medicine. It is not effective in children and adolescents between the ages of 10 and 17 years. It is not known if this medicine is safe and effective when used in children younger than 10 years.

Other medicines and Sitagliptin Rowex

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

In particular, tell your doctor if you are taking digoxin (a medicine used to treat irregular heartbeat and other heart problems). The level of digoxin in your blood may need to be checked if taking with Sitagliptin Rowex.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. You should not take this medicine during pregnancy.

It is not known if this medicine passes into breast milk. You should not take this medicine if you are breast-feeding or plan to breast-feed.

Driving and using machines

This medicine has no or negligible influence on the ability to drive and use machines. However, dizziness and drowsiness have been reported, which may affect your ability to drive or use machines. Taking this medicine in combination with medicines called sulphonylureas or with insulin can cause low blood sugar (hypoglycaemia), which may affect your ability to drive and use machines or work without safe foothold.

Sitagliptin Rowex contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially ‘sodium free’.

3. How to take Sitagliptin Rowex

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The usual recommended dose is:

- one 100 mg film-coated tablet
- once a day
- by mouth.

If you have kidney problems, your doctor may prescribe lower doses (such as 25 mg or 50 mg). You can take this medicine with or without food and drink.

Your doctor may prescribe this medicine alone or with certain other medicines that lower blood sugar. Diet and exercise can help your body use its blood sugar better. It is important to stay on the diet and exercise recommended by your doctor while taking Sitagliptin Rowex.

If you take more Sitagliptin Rowex than you should

If you take more than the prescribed dose of this medicine, contact your doctor immediately.

If you forget to take Sitagliptin Rowex

If you miss a dose, take it as soon as you remember. If you do not remember until it is time for your next dose, skip the missed dose and go back to your regular schedule. Do not take a double dose to make up for a forgotten dose.

If you stop taking Sitagliptin Rowex

Continue to take this medicine as long as your doctor prescribes it so you can continue to help control your blood sugar. You should not stop taking this medicine without talking to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

STOP taking Sitagliptin Rowex and contact a doctor immediately if you notice any of the following serious side effects:

- Severe and persistent pain in the abdomen (stomach area) which might reach through to your back with or without nausea and vomiting, as these could be signs of an inflamed pancreas (pancreatitis).

If you have a serious allergic reaction (frequency not known), including rash, hives, blisters on the skin/peeling skin and swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing, stop taking this medicine and call your doctor right away. Your doctor may prescribe a medicine to treat your allergic reaction and a different medicine for your diabetes.

Some patients have experienced the following side effects after adding sitagliptin to metformin:

Common (may affect up to 1 in 10 people): low blood sugar, nausea, flatulence, vomiting

Uncommon (may affect up to 1 in 100 people): stomachache, diarrhoea, constipation, drowsiness

Some patients have experienced different types of stomach discomfort when starting the combination of sitagliptin and metformin together (frequency is common).

Some patients have experienced the following side effects while taking sitagliptin in combination with a sulphonylurea and metformin:

Very common (may affect more than 1 in 10 people): low blood sugar

Common: constipation

Some patients have experienced the following side effects while taking sitagliptin and pioglitazone:

Common: flatulence, swelling of the hands or legs

Some patients have experienced the following side effects while taking sitagliptin in combination with pioglitazone and metformin:

Common: swelling of the hands or legs

Some patients have experienced the following side effects while taking sitagliptin in combination with insulin (with or without metformin):

Common: flu

Uncommon: dry mouth

Some patients have experienced the following side effects while taking sitagliptin alone in clinical studies, or during post-approval use alone and/or with other diabetes medicines:

Common: low blood sugar, headache, upper respiratory infection, stuffy or runny nose and sore throat, osteoarthritis, arm or leg pain

Uncommon: dizziness, constipation, itching

Rare: reduced number of platelets

Frequency not known: kidney problems (sometimes requiring dialysis), vomiting, joint pain, muscle pain, back pain, inflammation of the lungs (interstitial lung disease), bullous pemphigoid (a type of skin blister)

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system: HPRA Pharmacovigilance; website: www.hpra.ie. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Sitagliptin Rowex

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister foil after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Open the blister just before taking the tablets.

Do not use this medicine if any pack is damaged or shows signs of tampering.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Sitagliptin Rowex contains

- The active substance is sitagliptin.

Sitagliptin Rowex 25 mg tablets:

- Each film-coated tablet contains sitagliptin hydrochloride monohydrate equivalent to 25 mg sitagliptin.

Sitagliptin Rowex 50 mg tablets:

- Each film-coated tablet contains sitagliptin hydrochloride monohydrate equivalent to 50 mg sitagliptin.

Sitagliptin Rowex 100 mg tablets:

- Each film-coated tablet contains sitagliptin hydrochloride monohydrate equivalent to 100 mg sitagliptin.

The other ingredients are calcium hydrogen phosphate (E341), microcrystalline cellulose (E460), croscarmellose sodium (E468), magnesium stearate (E470b), sodium stearyl fumarate, hypromellose (E464), hydroxypropylcellulose (E463), macrogol (E1521), titanium dioxide (E171), yellow iron oxide (E172), red iron oxide (E172), black iron oxide (E172, only applicable for the 100 mg strength) and talc (E553b).

What Sitagliptin Rowex looks like and contents of the pack

Sitagliptin Rowex 25 mg tablets are pink, round, biconvex film-coated tablets debossed with “ST 25” on one side: diameter 5.7 – 6.6 mm.

Sitagliptin Rowex 50 mg tablets are light pink, round, biconvex film-coated tablets debossed with “ST 50” on one side: diameter 7.7 – 8.6 mm.

Sitagliptin Rowex 100 mg tablets are light brown, round, biconvex film-coated tablets debossed with “ST 100” on one side: diameter 9.7 – 10.6 mm.

Sitagliptin Rowex is available in

OPA/Al/PVC//Al and transparent PVC/PE/PVDC//Al blister packs containing 28, 30, 50, 56, 60, 84, 90, 98, 100, 105, 108, 110, 112, 120, 168, 180, 196, 198 or 200 film-coated tablets.

Pack of 28 x 1, 30 x 1, 50 x 1, 56 x 1, 84 x 1, 90 x 1, 98 x 1 film-coated tablets in perforated unit dose blisters.

HDPE bottle with a child resistant screw cap and polypropylene closure containing 30, 100 or 120 film-coated tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturers

Marketing Authorisation Holder

Rowex Ltd., Bantry, Co. Cork, Ireland.

Manufacturers

Lek Pharmaceuticals d.d., Verovškova Ulica 57, 1526 Ljubljana, Slovenia.

Lek Pharmaceuticals d.d., Trimlini 2D, 9220 Lendava, Slovenia.

Lek S.A., ul. Podlipie 16, Strykow, 95-010, Poland.

This medicinal product is authorised in the Member States of the EEA and in the United Kingdom (Northern Ireland) under the following names:

AT	Sitagliptin Sandoz 25 mg – Filmtabletten
	Sitagliptin Sandoz 50 mg – Filmtabletten
	Sitagliptin Sandoz 100 mg – Filmtabletten
BE	Sitagliptin Sandoz 25 mg filmomhulde tabletten
	Sitagliptin Sandoz 50 mg filmomhulde tabletten
	Sitagliptin Sandoz 100 mg filmomhulde tabletten
CZ	Sitagliptin Sandoz
DK	Sitagliptin Sandoz
ES	Sitagliptina Sandoz 25 mg comprimidos recubiertos con película EFG
	Sitagliptina Sandoz 50 mg comprimidos recubiertos con película EFG
	Sitagliptina Sandoz 100 mg comprimidos recubiertos con película EFG
FI	Sitagliptin Sandoz 25 mg kalvopäällysteiset tabletit
	Sitagliptin Sandoz 50 mg kalvopäällysteiset tabletit
	Sitagliptin Sandoz 100 mg kalvopäällysteiset tabletit
FR	SITAGLIPTINE SANDOZ 50 mg, comprimé pelliculé

	SITAGLIPTINE SANDOZ 100 mg, comprimé pelliculé
HU	Sitagliptin Sandoz 25 mg, 50 mg, 100 mg filmtabletta
IE	Sitagliptin Rowex 25 mg film-coated tablets Sitagliptin Rowex 50 mg film-coated tablets Sitagliptin Rowex 100 mg film-coated tablets
IS	Sitagliptin Sandoz 25 mg filmuhúðaðar töflur Sitagliptin Sandoz 50 mg filmuhúðaðar töflur Sitagliptin Sandoz 100 mg filmuhúðaðar töflur
IT	Sitagliptin Sandoz
NL	Sitagliptine Sandoz 25 mg, filmomhulde tabletten Sitagliptine Sandoz 50 mg, filmomhulde tabletten Sitagliptine Sandoz 100 mg, filmomhulde tabletten
NO	Sitagliptin Sandoz 25 mg filmdrasjerte tabletter Sitagliptin Sandoz 50 mg filmdrasjerte tabletter Sitagliptin Sandoz 100 mg filmdrasjerte tabletter
PT	Sitagliptina Sandoz Farmacêutica
SE	Sitagliptin Sandoz 25 mg filmdragerade tabletter Sitagliptin Sandoz 50 mg filmdragerade tabletter Sitagliptin Sandoz 100 mg filmdragerade tabletter
UK	Sitagliptin Sandoz 25mg film-coated tablets
(Northern Ireland)	Sitagliptin Sandoz 50mg film-coated tablets Sitagliptin Sandoz 100mg film-coated tablets

This leaflet was last revised in 01/2026.